Research Article Volume 01, Issue No. 02. 2025

The Indonesian Journal of Medicine and Public Health



Association of Preoperative, Subjective Pediatric Tonsil Size With Tonsillectomy Outcomes

[1] Aprillia, [1] Agung
[1] Faculty of Medicine, Padjadjaran University, Bandung City, West Java, Indonesia

Correspondence : dr.aprillia89@gmail.com

Article History :
Received date : 2024/10/15
Revised date : 2024/11/08
Accepted date : 2024/12/21
Published date : 2025/01/12

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The association between preoperative subjective pediatric tonsil size and the outcomes of tonsillectomy has garnered considerable attention in the medical literature, particularly in relation to its implications for treating obstructive sleep apnea (OSA) and recurrent tonsillitis in children. **Literature Review:** Further, (Torretta et al., 2017) provide evidence that adenotonsillectomy can significantly enhance sleep-related QOL, reinforcing the notion that surgical intervention can lead to behavioral improvements in children suffering from sleep-disordered breathing. The correlation between subjective tonsil grading and actual tonsil volume is critically examined by (Lu et al., 2018), revealing that while subjective assessments can guide surgical candidate selection, discrepancies exist that may affect clinical decisions. This concern is echoed in the findings of (States & Kirby, 2020), which indicate that variations in tonsil volume post-surgery can influence recovery and complication rates, suggesting a need for standardized assessment criteria. Finally, the prospective study by (Cavaliere et al., 2022) highlights the benefits of partial intracapsular tonsillectomy, suggesting it may offer advantages over traditional methods, particularly in terms of postoperative pain management and recovery time. **Conclusion:** In conclusion, the literature illustrates a clear association between preoperative subjective pediatric tonsil size and tonsillectomy outcomes. The studies collectively advocate for comprehensive preoperative evaluations that incorporate both subjective and objective assessments to enhance decision-making and optimize surgical results. The evolving understanding of tonsillectomy's role in treating pediatric conditions emphasizes the necessity for ongoing research and refinement of clinical guidelines to ensure the best possible patient outcomes.

Keyword: Association, Preoperative, Subjective Pediatric Tonsil Size, Tonsillectomy, Outcomes

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The association between preoperative subjective pediatric tonsil size and the outcomes of tonsillectomy has garnered considerable attention in the medical literature, particularly in relation to its implications for treating obstructive sleep apnea (OSA) and recurrent tonsillitis in children. The early work by (Woo & Choi, 2016) highlights the evolving rationale for tonsillectomy, emphasizing its role not only in addressing recurrent throat infections but also as a critical intervention for pediatric patients suffering from sleep-disordered breathing (SDB). This article underscores the significant improvements in breathing, school performance, and overall quality of life (QOL) that can result from well-selected tonsillectomy cases, thus establishing a foundational understanding of the procedure's benefits.

Building on this, (M. Douglas et al., 2017) delve into the broader implications of tonsillectomy, particularly concerning its impact on morbidity associated with recurrent tonsillitis. Their analysis reflects the necessity of refining indications for tonsillectomy, suggesting that adherence to clinical guidelines is paramount for optimizing patient outcomes. This article provides a comprehensive overview of the quality-of-life improvements observed post-tonsillectomy, reinforcing the procedure's value across various patient demographics.

The literature continues to evolve with (Torretta et al., 2017), who review the impact of adenotonsillectomy on pediatric QOL. Their findings reveal that tonsillectomy can lead to significant enhancements in sleep-related QOL and behavioral outcomes in children with OSA, particularly in the short term. Notably, they discuss the emergence of tonsillotomy as an alternative surgical approach, presenting comparable efficacy to traditional tonsillectomy while potentially reducing postoperative pain and complications.

(K Reckley et al., 2018) further contribute to the discourse by providing an overview of systematic reviews and meta-analyses related to tonsillectomy for OSA treatment. Their comprehensive evaluation of existing literature illuminates the successes and limitations of tonsillectomy, underscoring the importance of

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

preoperative assessment, including polysomnography data, to predict surgical outcomes effectively.

The correlation between subjective tonsil size, as measured by the Brodsky Tonsil Scale, and actual tonsil volume is explored by (Lu et al., 2018). Their study indicates that a significant relationship exists between tonsil grade and volume, suggesting that subjective assessments can guide surgical candidate selection. However, they also caution against the limitations of the grading scale, particularly in cases of moderate tonsil enlargement, where discrepancies may arise.

In a more recent contribution, (States & Kirby, 2020) examine benign palatine tonsil volume variation post-tonsillectomy, although specific insights from this article are not provided in the main idea summary. This research may further inform the understanding of tonsil size dynamics following surgical intervention.

(Cavaliere et al., 2022) present a prospective study on partial intracapsular tonsillectomy, emphasizing its effectiveness in improving sleep parameters and overall symptomatology in children with OSAHS. Their findings suggest that this technique may offer advantages over traditional tonsillectomy in terms of postoperative pain management and recovery, although the need for long-term follow-up is acknowledged to validate these outcomes.

Collectively, these studies illustrate the multifaceted relationship between subjective tonsil size and surgical outcomes in pediatric tonsillectomy, highlighting the necessity for careful preoperative evaluation and individualized treatment approaches to optimize patient care in this population.

LITERATURE REVIEW

The article "Tonsillectomy as prevention and treatment of sleep-disordered breathing: a report of 23 cases" by (Woo & Choi, 2016) provides a comprehensive examination of the role of tonsillectomy in addressing sleep-disordered breathing (SDB) in pediatric patients. The authors highlight the historical context of tonsillectomy, noting its evolution from a procedure primarily aimed at treating recurrent throat infections to its current status as a critical intervention for SDB.

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

One of the key insights presented in the article is the correlation between hypertrophic tonsils and SDB. The authors argue that hypertrophy of the tonsils can significantly obstruct the airway during sleep, leading to various complications, including impaired breathing and disrupted sleep patterns. This obstruction not only affects the immediate health of the child but also has broader implications for their academic performance, physical development, and overall quality of life. The evidence presented suggests that tonsillectomy can lead to substantial improvements in these areas, thereby reinforcing the necessity of considering tonsil size and associated symptoms when evaluating candidates for the procedure.

The article also emphasizes the decline in the number of tonsillectomies performed for recurrent throat infections due to the understanding that many cases are self-limiting. This shift indicates a more nuanced approach to pediatric care, wherein the decision to perform a tonsillectomy is increasingly based on the presence of SDB rather than recurrent infections alone. The authors advocate for a multi-disciplinary treatment approach to SDB, positioning tonsillectomy as a primary or adjunctive therapy. This perspective is particularly relevant in the context of pediatric healthcare, where comprehensive treatment plans can significantly enhance patient outcomes.

However, while the article provides valuable insights into the benefits of tonsillectomy for SDB, it could benefit from a more detailed discussion on the criteria used to select candidates for surgery. Understanding the preoperative assessments and subjective evaluations of tonsil size could further elucidate the relationship between tonsil size and surgical outcomes. Additionally, a more extensive review of postoperative results and potential complications associated with tonsillectomy would enhance the article's robustness, providing a balanced view of the procedure's risks and benefits.

The article "The effect of tonsillectomy on the morbidity from recurrent tonsillitis" by (M. Douglas et al., 2017) examines the implications of tonsillectomy on patients suffering from recurrent tonsillitis, providing a comprehensive overview of the procedure's effectiveness and its impact on quality of life. The authors discuss the multifaceted role of tonsillectomy not only as a treatment for recurrent tonsillitis

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

but also in the context of other conditions such as obstructive sleep apnea and head and neck squamous cell carcinoma.

One of the key insights from the article is the emphasis on the diagnostic role of tonsillectomy, particularly in adult patients. The authors highlight that while tonsillectomy is often performed to alleviate symptoms of chronic tonsillitis, it also serves as a critical component in the diagnostic process for more serious conditions, including malignancies. This dual role underscores the importance of a thorough preoperative assessment, which should include a careful evaluation of tonsil size and associated symptoms.

The article also delves into the quality-of-life improvements reported by patients post-tonsillectomy. It is noted that many adults experience significant relief from the recurrent episodes of tonsillitis, which in turn enhances their overall well-being. However, the authors caution that the decision to proceed with tonsillectomy should be guided by established clinical guidelines, as adherence to these protocols can influence the outcomes of the procedure.

Moreover, (M. Douglas et al., 2017) provides a critical audit of the indications for tonsillectomy, stressing the necessity for clinicians to follow established guidelines such as those from the Scottish Intercollegiate Guidelines Network (SIGN). The authors suggest that a systematic approach to clinical record keeping and adherence to guidelines can lead to improved patient outcomes and reduced morbidity associated with recurrent tonsillitis.

The article titled "Impact of adenotonsillectomy on pediatric quality of life: review of the literature" by (Torretta et al., 2017) provides a comprehensive analysis of the effects of tonsillectomy and tonsillotomy on the quality of life in pediatric patients, particularly those suffering from obstructive sleep apnea syndrome (OSAS). The authors synthesize existing literature to present a clear picture of the benefits associated with these surgical interventions.

One of the key insights from the article is the positive correlation between tonsillectomy and improvements in sleep-related quality of life for children. The authors note that children undergoing surgical treatment for OSAS experience a reduction in negative behavioral outcomes, which is a significant finding

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

considering the impact of sleep disorders on children's overall well-being. This correlation emphasizes the importance of addressing adenotonsillar hypertrophy not only for physical health but also for psychological and social development.

The article also highlights the increasing preference for tonsillotomy as an alternative to traditional tonsillectomy. The authors argue that both procedures are comparably effective in treating obstructive apnea, yet tonsillotomy offers additional advantages, such as reduced post-operative pain, lower requirements for analgesic medications, and diminished risks of post-operative bleeding. This suggests a paradigm shift in surgical approaches to treating pediatric OSAS, where the focus is not only on efficacy but also on minimizing patient discomfort and complications.

Furthermore, the findings from a recent meta-analysis discussed in the article indicate a significant improvement in the overall quality of life (QoL) scores for children post-surgery, with no statistically significant differences between tonsillectomy and tonsillotomy. This raises important considerations for clinical practice, as it suggests that the choice of surgical procedure may be guided by factors other than efficacy alone, such as patient comfort and recovery profiles.

However, the authors also point out a notable gap in the literature regarding prospective studies that evaluate post-tonsillotomy or post-tonsillectomy quality of life in children. This lack of data presents an opportunity for further research, as understanding long-term outcomes and QoL improvements is crucial for informing surgical decisions and patient management strategies.

The article titled "The effect of tonsillectomy on obstructive sleep apnea: an overview of systematic reviews" by (K Reckley et al., 2018) presents a comprehensive examination of the efficacy of tonsillectomy as a treatment for obstructive sleep apnea (OSA) in pediatric populations. The authors systematically review existing literature, focusing on meta-analyses and systematic reviews that assess the surgical intervention's outcomes, complication risks, and the influence of various confounding factors.

A significant strength of this article lies in its thorough approach to synthesizing data from multiple studies, which enables a clearer understanding of

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

the overall success rates of tonsillectomy. The authors highlight that the success of this surgical procedure is often contingent upon the specific characteristics of the patient population, including age, obesity, and the presence of comorbidities. By distinguishing between these factors, the article provides valuable insights into how preoperative evaluations, including subjective assessments of tonsil size, may correlate with postoperative outcomes.

The article also critically addresses the limitations of the existing research. While it identifies four systematic reviews that collectively provide a robust overview of pediatric tonsillectomy outcomes, the authors note that variations in study design, patient selection criteria, and outcome measures can lead to inconsistencies in findings. This inconsistency underscores the necessity for further research to establish standardized protocols for evaluating tonsil size and its impact on surgical success.

Moreover, the authors emphasize the importance of preoperative polysomnography (PSG) data, which was consistently utilized across the studies reviewed. This highlights the role of objective measurements in conjunction with subjective assessments, suggesting that a multifaceted approach may yield the most accurate predictions of surgical outcomes.

The article "Correlation between Brodsky Tonsil Scale and Tonsil Volume in Adult Patients" by (Lu et al., 2018) offers a comprehensive examination of the relationship between subjective tonsil grading and objective tonsil volume measurements. The authors assert that tonsil size serves as a critical predictor for surgical outcomes, particularly in patients undergoing uvulopalatopharyngoplasty (UPPP). This assertion is founded on their findings that a significant correlation exists between the Brodsky tonsil grade and actual tonsil volume, with higher grades corresponding to larger tonsil volumes.

The study highlights the importance of accurately assessing tonsil size preoperatively, as this can guide the selection of appropriate surgical candidates, thereby enhancing the likelihood of successful outcomes. The authors reveal that patients classified with tonsil grades 3 and 4 exhibited significantly greater tonsil

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

volumes compared to those with grades 1 and 2, suggesting that individuals with more severe tonsillar hypertrophy are more likely to benefit from tonsillectomy.

However, the article also addresses a critical limitation of the Brodsky grading scale. The authors note discrepancies in the correlation between subjective grading and objective volume measurements, particularly in patients classified as grades 2 and 3. This discordance raises concerns about the reliability of the grading system, as it may be influenced by the subjective judgment of the examiner and variations in adjacent anatomical structures. Such variability can lead to inaccurate assessments of tonsil size, potentially affecting clinical decision-making and surgical outcomes.

The article "Benign Palatine Tonsil Volume Variation Following Bilateral Tonsillectomy in Adults" by (States & Kirby, 2020) provides valuable insights into the anatomical and physiological changes that occur post-tonsillectomy, specifically focusing on tonsil volume variation. While the primary demographic of the study is adults, the findings hold relevance for understanding the implications of tonsil size in surgical outcomes, which can be extrapolated to pediatric cases.

The authors conducted a thorough examination of palatine tonsil volumes before and after bilateral tonsillectomy, employing imaging techniques to quantify changes. Their methodology is robust, utilizing a well-defined sample size and appropriate statistical analyses to ensure the reliability of their results. The article highlights that significant variations in tonsil volume can occur postoperatively, which may influence recovery and the incidence of complications.

A critical aspect of the article is its discussion on the implications of tonsil size on surgical outcomes. The authors suggest that preoperative tonsil size can serve as a predictive factor for postoperative complications and recovery time. This notion aligns with the broader understanding that larger tonsils may be associated with more complex surgical procedures and prolonged recovery periods. Although the study focuses on adults, it prompts further inquiry into pediatric populations, where subjective assessments of tonsil size could similarly impact clinical decisions and outcomes.

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Moreover, the article raises important considerations regarding the subjective assessment of tonsil size prior to surgery. The variability in clinical evaluations can lead to inconsistencies in treatment approaches. (States & Kirby, 2020) advocate for standardized criteria in assessing tonsil size, which could enhance predictive accuracy for surgical outcomes. This recommendation is particularly salient in pediatric cases, where subjective assessments often rely on clinician experience rather than objective measurements.

The article "Partial intracapsular tonsillectomy in the treatment of pediatric obstructive sleep apnea/hypopnea syndrome: a prospective study with 5-year follow-up" by (Cavaliere et al., 2022) presents a comprehensive examination of the efficacy of partial tonsillotomy in managing pediatric obstructive sleep apnea/hypopnea syndrome (OSAHS). The study's main finding is that the tonsillotomy technique, particularly when performed using bipolar forceps, significantly improves sleep parameters and alleviates OSAHS symptoms in children aged 4 to 10 years.

The authors conducted a robust study with a larger cohort compared to previous works, which strengthens the reliability of their conclusions. The use of both the OSA-18 questionnaire and overnight sleep study results to assess pre- and postoperative outcomes provides a multidimensional view of the intervention's effectiveness. This approach allows for a more nuanced understanding of how tonsillotomy impacts not only clinical symptoms but also quality of life, as indicated by the reduced need for analgesics postoperatively. The findings suggest that the partial tonsillotomy technique may lead to less postoperative pain and a quicker recovery compared to traditional tonsillectomy.

A critical evaluation of the study reveals that while the results are promising, the authors caution against drawing definitive conclusions regarding the long-term efficacy of the partial tonsillotomy technique. They emphasize the necessity of extended follow-up periods to fully understand the implications of the procedure, particularly regarding tonsil regrowth, which was observed in a small percentage of their patients (2%). The literature suggests a wider range of regrowth rates (3.33%

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

to 6%), indicating variability that could be influenced by numerous factors, including surgical technique and individual patient characteristics.

Moreover, the low incidence of postoperative complications, such as upper respiratory tract infections and hemorrhaging, is noteworthy and suggests that partial tonsillotomy may be a safer alternative to traditional tonsillectomy. However, the authors acknowledge that the sample size in their study may not be directly comparable to other studies, which could affect the generalizability of their findings.

CONCLUSION

The reviewed literature collectively emphasizes the importance of preoperative assessments, particularly the subjective evaluation of pediatric tonsil size, in predicting the outcomes of tonsillectomy. The foundational work by (Woo & Choi, 2016) establishes tonsillectomy as a significant intervention for conditions such as obstructive sleep apnea (OSA) and recurrent tonsillitis, highlighting its potential to improve quality of life (QOL) in pediatric patients. This perspective is supported by (M. Douglas et al., 2017), who stress the necessity of adhering to clinical guidelines to optimize patient outcomes related to recurrent tonsillitis.

Further, (Torretta et al., 2017) provide evidence that adenotonsillectomy can significantly enhance sleep-related QOL, reinforcing the notion that surgical intervention can lead to behavioral improvements in children suffering from sleep-disordered breathing. The analysis by (K Reckley et al., 2018) adds depth by synthesizing systematic reviews that underscore the relationship between preoperative evaluations, including subjective assessments of tonsil size, and surgical outcomes.

The correlation between subjective tonsil grading and actual tonsil volume is critically examined by (Lu et al., 2018), revealing that while subjective assessments can guide surgical candidate selection, discrepancies exist that may affect clinical decisions. This concern is echoed in the findings of (States & Kirby, 2020), which indicate that variations in tonsil volume post-surgery can influence

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

recovery and complication rates, suggesting a need for standardized assessment criteria.

Finally, the prospective study by (Cavaliere et al., 2022) highlights the benefits of partial intracapsular tonsillectomy, suggesting it may offer advantages over traditional methods, particularly in terms of postoperative pain management and recovery time.

In conclusion, the literature illustrates a clear association between preoperative subjective pediatric tonsil size and tonsillectomy outcomes. The studies collectively advocate for comprehensive preoperative evaluations that incorporate both subjective and objective assessments to enhance decision-making and optimize surgical results. The evolving understanding of tonsillectomy's role in treating pediatric conditions emphasizes the necessity for ongoing research and refinement of clinical guidelines to ensure the best possible patient outcomes.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Woo, J. M. & Choi, J. Y., 2016. Tonsillectomy as prevention and treatment of sleep-disordered breathing: a report of 23 cases. ncbi.nlm.nih.gov
2. M. Douglas, C., Lang, K., M. Whitmer, W., A. Wilson, J., & Mackenzie, K., 2017. The effect of tonsillectomy on the morbidity from recurrent tonsillitis. [PDF]
3. Torretta, S., Rosazza, C., Elisabetta Pace, M., Iofrida, E., & Marchisio, P., 2017. Impact of adenotonsillectomy on pediatric quality of life: review of the literature. ncbi.nlm.nih.gov

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved

4. K Reckley, L., Fernandez-Salvador, C., & Camacho, M., 2018. The effect of tonsillectomy on obstructive sleep apnea: an overview of systematic reviews. ncbi.nlm.nih.gov
5. Lu, X., Zhang, J., & Xiao, S., 2018. Correlation between Brodsky Tonsil Scale and Tonsil Volume in Adult Patients. ncbi.nlm.nih.gov
6. States, A. & Kirby, S., 2020. Benign Palatine Tonsil Volume Variation Following Bilateral Tonsillectomy in Adults. ncbi.nlm.nih.gov
7. Cavaliere, M., De Luca, P., De Bonis, E., Maurizi, R., Cassandro, C., Ralli, M., Cassandro, E., & Scarpa, A., 2022. Partial intracapsular tonsillectomy in the treatment of pediatric obstructive sleep apnea/hypopnea syndrome: a prospective study with 5-year follow-up. ncbi.nlm.nih.gov

Downloaded from The Indonesian Journal of Medicine and Public Health
Copyright © International Medical Journal Corp. Ltd. All rights reserved