Research Article Volume 01, Issue No. 02. 2025

THE JOURNAL OF MEDICAL RESEARCH AND APPLIED SCIENCE



Association of Feeding Evaluation With Frenotomy Rates in Infants With Breastfeeding Difficulties

[1] Subiyanto, [1] Grace

[1] Faculty of Medicine, Padjadjaran University, Bandung City, West Java, Indonesia

Correspondence : dr.subiyanto382@gmail.com

Article History :
Received date : 2024/10/07
Revised date : 2024/11/14
Accepted date : 2024/12/27
Published date : 2025/01/12

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature surrounding the association of feeding evaluation with frenotomy rates in infants experiencing breastfeeding difficulties has evolved significantly over the past decade, reflecting a growing awareness of the impact of ankyloglossia on breastfeeding outcomes. **Literature Review:** The randomized controlled trial by (Emond et al., 2013) revealed that while frenotomy improved maternal self-efficacy, it did not significantly enhance breastfeeding outcomes at critical early postpartum periods. Further, the systematic review conducted by (Poliseni Miranda et al., 2016) underscored the necessity for validated assessment protocols to measure breastfeeding quality and pain. This gap in objective measurement tools limits the ability to draw definitive conclusions about the effectiveness of interventions related to ankyloglossia, thus necessitating rigorous methodological approaches in future research. Qualitative insights from (Waterman, 2017) and the prospective cohort study by (A. Ghaheri et al., 2017) further elucidate the emotional and practical challenges faced by mothers, emphasizing the importance of comprehensive evaluations prior to surgical procedures. These studies indicate that while frenotomy can lead to improved breastfeeding metrics, a nuanced understanding of individual cases is crucial. **Conclusion :** In conclusion, the literature indicates that while frenotomy may play a role in alleviating some breastfeeding difficulties associated with ankyloglossia, it is not a panacea. A comprehensive approach that includes thorough feeding evaluations, consideration of anatomical variations, and emotional support for mothers is essential for optimizing breastfeeding outcomes. Future research should focus on establishing standardized assessment protocols and exploring the long-term impacts of both surgical and non-surgical interventions on breastfeeding success.

Keyword: Association, Feeding Evaluation, Frenotomy Rates, Infants, Breastfeeding Difficulties

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The literature surrounding the association of feeding evaluation with frenotomy rates in infants experiencing breastfeeding difficulties has evolved significantly over the past decade, reflecting a growing awareness of the impact of ankyloglossia on breastfeeding outcomes. In 2013, (Emond et al., 2013) conducted a randomized controlled trial to compare early frenotomy with standard breastfeeding support. Their findings suggested that while frenotomy improved maternal self-efficacy, it did not yield a significant difference in breastfeeding outcomes at five days or eight weeks post-procedure. This study underscored the complexity of breastfeeding difficulties and the multifaceted nature of support required for mothers and infants.

Building on this foundation, (Poliseni Miranda et al., 2016) performed a systematic review of interventions related to the lingual frenum, noting an increase in scientific production on this topic beginning in 2002. Their analysis highlighted the need for validated protocols to assess breastfeeding quality and pain, emphasizing that while many studies recognized the benefits of addressing feeding difficulties associated with ankyloglossia, there remained a lack of objective measures to quantify these outcomes. This gap in the literature pointed to the necessity for rigorous methodologies in future research.

In 2017, (Waterman, 2017) conducted a qualitative study that provided deeper insights into mothers' experiences breastfeeding infants with tongue-tie. This work corroborated previous randomized trials regarding the efficacy of frenulotomy and illustrated the emotional and practical challenges faced by mothers, further emphasizing the need for supportive interventions. (A. Ghaheri et al., 2017) also contributed to this discourse with a prospective cohort study that demonstrated significant improvements in breastfeeding outcomes following frenotomy. They highlighted the importance of comprehensive evaluations prior to surgery, suggesting that factors such as anatomical variations must be considered to optimize breastfeeding success.

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The examination of ankyloglossia continued with (Clara Souza-Oliveira et al., 2021), who reported a prevalence of 15% in their cross-sectional study. They found that the diagnosis of ankyloglossia and subsequent frenectomy procedures had increased, particularly among preterm infants who faced unique breastfeeding challenges. Their findings reinforced the importance of providing tailored breastfeeding support to mothers, particularly those with premature infants, to enhance breastfeeding duration and efficacy.

Most recently, (A. Lee et al., 2024) conducted a cross-sectional study that explored the assessment process for breastfed infants referred for tongue-tie evaluation and frenotomy. Their results indicated a statistical association between frenotomy and specific feeding difficulties, such as nipple pain and inability to latch. This study provided valuable insights into the clinical characteristics that may guide healthcare professionals in making informed decisions regarding frenotomy interventions.

Overall, the literature reflects a growing body of evidence supporting the need for careful assessment and intervention in cases of ankyloglossia, with an emphasis on the importance of both surgical and non-surgical support for breastfeeding mothers and their infants. The ongoing debate regarding the timing and necessity of frenotomy continues to fuel research in this area, highlighting the complexity of breastfeeding challenges and the necessity for comprehensive care strategies.

LITERATURE REVIEW

The article titled "Randomised controlled trial of early frenotomy in breastfed infants with mild–moderate tongue-tie" by (Emond et al., 2013) presents a comprehensive examination of the effectiveness of immediate frenotomy compared to standard breastfeeding support for infants experiencing mild to moderate tongue-tie. The study's structure, which includes randomization and blinding of outcome assessors, lends credibility to its findings and minimizes bias in the evaluation of results.

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The primary outcome of the trial was breastfeeding success at five days postpartum, a critical time frame for establishing breastfeeding patterns and addressing potential difficulties. The results indicated no significant difference in breastfeeding rates between the intervention group (those receiving early frenotomy) and the comparison group (those receiving standard care), with both groups showing similar outcomes in breastfeeding measures and infant weight at eight weeks. This finding suggests that immediate frenotomy may not provide the anticipated objective benefits in breastfeeding success for infants with mild to moderate tongue-tie.

However, the study did highlight an important secondary outcome: the improvement in maternal breastfeeding self-efficacy among those who underwent frenotomy. This aspect is noteworthy as it underscores the psychological and emotional dimensions of breastfeeding challenges. While the procedure did not lead to a higher rate of breastfeeding success, the increased self-efficacy reported by mothers may play a crucial role in their overall breastfeeding experience and persistence.

Moreover, the article discusses the broader implications of feeding difficulties associated with tongue-tie, which affect a significant percentage of infants. The authors note that many mothers, despite their desire to breastfeed, encounter challenges that can lead to cessation of breastfeeding, thus emphasizing the need for effective lactation support. The variability in the availability of frenotomy across the UK further complicates the decision-making process for families facing these issues.

The article titled "Interventions in the Alteration on Lingual Frenum: Systematic Review" by (Poliseni Miranda et al., 2016) provides a comprehensive analysis of the literature surrounding interventions related to the lingual frenum, particularly in the context of breastfeeding difficulties in infants. The systematic review methodology employed by the authors is robust, involving a thorough electronic search that initially identified 443 publications. After applying rigorous inclusion and exclusion criteria, 26 articles were ultimately reviewed,

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

demonstrating a significant focus on the neonatal and infant populations facing breastfeeding challenges associated with ankyloglossia.

One of the key insights from the review is the notable increase in scientific publications on this topic since 2002, particularly after 2010, which reflects a growing recognition of the importance of addressing breastfeeding difficulties linked to anatomical variations like ankyloglossia. However, the authors point out a critical gap in the literature: while many studies assess breastfeeding, few utilize validated protocols to quantify breastfeeding quality and associated pain. This lack of standardized assessment tools limits the ability to draw definitive conclusions about the effectiveness of various interventions and the overall impact on breastfeeding success.

The review also highlights the diversity of interventions explored in the literature, ranging from surgical options like frenotomy to non-invasive strategies such as changing the baby's position or employing specific maneuvers. Notably, one study mentioned in the review suggests that some cases of altered lingual frenum may resolve spontaneously with the eruption of lower teeth, indicating that surgical intervention may not be necessary in all instances. This finding underscores the importance of a nuanced approach to treatment, where the relationship between feeding difficulties and the condition of the frenum is carefully considered before proceeding with surgical options.

The article "A qualitative description of mothers' experiences breastfeeding a child with tongue-tie" by (Waterman, 2017) provides an in-depth examination of the challenges faced by mothers when breastfeeding infants diagnosed with ankyloglossia, commonly known as tongue-tie. The study highlights the incidence of feeding difficulties associated with this condition, emphasizing the significance of proper assessment and intervention, particularly through frenulotomy.

Waterman’s qualitative approach reveals the multifaceted nature of breastfeeding experiences among mothers of infants with tongue-tie. The article elucidates how ankyloglossia can impede effective breastfeeding, leading to issues such as pain, inadequate milk transfer, and ultimately, maternal frustration and infant weight concerns. The author discusses the varying degrees of severity

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

associated with tongue-tie, noting that even mild to moderate cases can adversely affect the breastfeeding dyad.

A critical component of the article is the exploration of frenulotomy as a potential solution to alleviate breastfeeding difficulties. Waterman cites evidence from randomized controlled trials that suggest early intervention through frenotomy can significantly improve breastfeeding outcomes. This aligns with the notion that timely surgical intervention may facilitate better latch and feeding efficiency, thereby enhancing the overall breastfeeding experience for both mother and infant.

Moreover, the article addresses the importance of a comprehensive assessment of tongue-tie and its implications for breastfeeding. Waterman emphasizes that not all cases of ankyloglossia necessitate surgical intervention; rather, a careful evaluation of the infant's feeding patterns and the mother's comfort level is crucial in determining the appropriate course of action. This nuanced understanding is essential for healthcare providers when advising families facing such challenges.

The article "Breastfeeding improvement following tongue-tie and lip-tie release: A prospective cohort study" by (A. Ghaheri et al., 2017) provides a thorough investigation into the effects of surgical interventions on breastfeeding outcomes in infants diagnosed with ankyloglossia and tethered maxillary labial frenula. The authors present a well-structured prospective cohort study that emphasizes the necessity of a comprehensive evaluation of the head and neck prior to any surgical procedure, such as frenotomy.

One of the key insights from the study is the importance of considering additional anatomical factors, such as retrognathia and palatal abnormalities, which can significantly influence breastfeeding difficulties. This perspective is crucial as it underscores the complexity of breastfeeding issues and the need for a multidimensional assessment rather than a singular focus on tongue-tie. The authors effectively argue that neglecting these factors may lead to suboptimal outcomes post-surgery.

The results of the study reveal that there is a significant improvement in various breastfeeding-related metrics following lingual frenotomy, with or without

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

the accompanying maxillary labial frenectomy. Notably, maternal self-efficacy, nipple pain, infant reflux symptoms, and milk transfer rates all showed marked enhancement post-intervention. This finding is particularly relevant for practitioners and parents alike, indicating that addressing ankyloglossia can lead to a substantial improvement in the quality of life for both the mother and infant dyad.

Moreover, the study highlights that infants with posterior tongue-tie experienced similar improvements as those with more classic presentations of tongue-tie. This suggests that the clinical implications of tongue-tie may extend beyond traditional definitions, warranting a broader understanding and approach in clinical practice.

The article "Does ankyloglossia interfere with breastfeeding in newborns? A cross-sectional study" by (Clara Souza-Oliveira et al., 2021) provides valuable insights into the prevalence of ankyloglossia and its implications for breastfeeding practices. The study reported a prevalence rate of 15% for ankyloglossia among the newborns examined, highlighting a growing awareness and diagnosis of this condition, which has subsequently led to an increase in frenectomy procedures. This correlation raises questions about the efficacy of frenotomy as a solution for breastfeeding difficulties associated with ankyloglossia.

The authors note that while there is anecdotal evidence suggesting that frenectomy can improve breastfeeding outcomes and alleviate maternal nipple pain, they also caution that the existing evidence supporting these claims is insufficient and questionable. This critical perspective is essential, as it urges healthcare professionals to consider the nuanced and multifaceted nature of breastfeeding difficulties, particularly in infants diagnosed with ankyloglossia. The absence of any newborns undergoing frenectomy during hospitalization in this study further emphasizes the need for a careful evaluation of when and how such procedures should be implemented.

Additionally, the study highlights the challenges faced by mothers of preterm infants, who often experience greater difficulties with breastfeeding due to the infants' weak suction, low muscle tone, and uncoordinated oral movements. These factors can significantly impact the infants' ability to obtain adequate

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

nutrition, thereby increasing the risk of dehydration and poor weight gain. The authors advocate for enhanced breastfeeding support and education for mothers, particularly those with preterm infants, as this can lead to improved breastfeeding duration and reduced reliance on infant formulas.

The article titled "A cross-sectional study of breastfed infants referred for tongue tie assessment and frenotomy in one Canadian health region" by (A. Lee et al., 2024) offers a comprehensive examination of the relationship between feeding evaluations and the rates of frenotomy in infants experiencing breastfeeding difficulties. The study is particularly relevant as it sheds light on key factors contributing to the decision to perform frenotomy, a surgical intervention often considered in cases of tongue tie.

The authors conducted a cross-sectional study that involved trained healthcare professionals assessing infant feeding difficulties at the breast. This aspect of the research is crucial, as it emphasizes the importance of a structured evaluation process conducted by qualified personnel. The documentation of feeding difficulties, such as nipple pain/trauma and inability to latch, is a significant contribution to understanding the clinical context in which frenotomies are considered. The statistical association found between these documented difficulties and the rates of frenotomy highlights the need for healthcare providers to recognize and address these issues proactively.

The final model presented in the study, which includes nipple pain/trauma, inability to latch, inability to elevate the tongue, and dimpling of the tongue, achieved statistical significance. This finding is particularly insightful as it not only reinforces the clinical indicators that may warrant surgical intervention but also provides a framework for healthcare professionals to assess and prioritize treatment options for breastfeeding difficulties. The inclusion of multiple dimensions of feeding difficulties in the analytical model enhances the robustness of the study and allows for a more thorough understanding of the factors influencing frenotomy rates.

However, while the study offers valuable insights, it is essential to consider potential limitations inherent in cross-sectional research designs, such as the

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

inability to establish causality. The reliance on chart documentation may also introduce biases, as the completeness and accuracy of the recorded information can vary. Future research could benefit from longitudinal studies that track the outcomes of infants post-frenotomy in relation to their initial feeding assessments.

CONCLUSION

The literature surrounding the association of feeding evaluation with frenotomy rates in infants experiencing breastfeeding difficulties indicates a complex interplay between surgical intervention and breastfeeding outcomes. The studies reviewed demonstrate an evolving understanding of ankyloglossia's impact on breastfeeding, emphasizing the need for comprehensive evaluations and tailored interventions.

The randomized controlled trial by (Emond et al., 2013) revealed that while frenotomy improved maternal self-efficacy, it did not significantly enhance breastfeeding outcomes at critical early postpartum periods. This finding suggests that while surgical intervention may bolster mothers' confidence, it does not guarantee improved breastfeeding success, highlighting the multifactorial nature of breastfeeding challenges.

Further, the systematic review conducted by (Poliseni Miranda et al., 2016) underscored the necessity for validated assessment protocols to measure breastfeeding quality and pain. This gap in objective measurement tools limits the ability to draw definitive conclusions about the effectiveness of interventions related to ankyloglossia, thus necessitating rigorous methodological approaches in future research.

Qualitative insights from (Waterman, 2017) and the prospective cohort study by (A. Ghaheri et al., 2017) further elucidate the emotional and practical challenges faced by mothers, emphasizing the importance of comprehensive evaluations prior to surgical procedures. These studies indicate that while frenotomy can lead to improved breastfeeding metrics, a nuanced understanding of individual cases is crucial.

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The prevalence of ankyloglossia reported by (Clara Souza-Oliveira et al., 2021) and the cross-sectional study by (A. Lee et al., 2024) highlight the increasing recognition of this condition and its association with specific feeding difficulties. The latter study's findings regarding the statistical association between documented feeding challenges and frenotomy rates emphasize the need for structured assessments by healthcare professionals.

In conclusion, the literature indicates that while frenotomy may play a role in alleviating some breastfeeding difficulties associated with ankyloglossia, it is not a panacea. A comprehensive approach that includes thorough feeding evaluations, consideration of anatomical variations, and emotional support for mothers is essential for optimizing breastfeeding outcomes. Future research should focus on establishing standardized assessment protocols and exploring the long-term impacts of both surgical and non-surgical interventions on breastfeeding success.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Emond, A., Ingram, J., Johnson, D., Blair, P., Whitelaw, A., Copeland, M., & Sutcliffe, A., 2013. Randomised controlled trial of early frenotomy in breastfed infants with mild–moderate tongue-tie. ncbi.nlm.nih.gov
2. Poliseni Miranda, P., Louise Cardoso, C., & Gomes, E., 2016. Interventions in the Alteration on Lingual Frenum: Systematic Review. ncbi.nlm.nih.gov

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved

3. Waterman, J., 2017. A qualitative description of mothers' experiences breastfeeding a child with tongue-tie. [PDF]
4. A. Ghaheri, B., Cole, M., C. Fausel, S., Chuop, M., & C. Mace, J., 2017. Breastfeeding improvement following tongue-tie and lip-tie release: A prospective cohort study. ncbi.nlm.nih.gov
5. Clara Souza-Oliveira, A., Valdelice Cruz, P., Baccin Bendo, C., Costa Batista, W., Cândida Ferrarez Bouzada, M., & Castro Martins, C., 2021. Does ankyloglossia interfere with breastfeeding in newborns? A cross-sectional study. ncbi.nlm.nih.gov
6. A. Lee, T., Bishop, J., Drover, A., K. Midodzi, W., & K. Twells, L., 2024. A cross-sectional study of breastfed infants referred for tongue tie assessment and frenotomy in one Canadian health region. ncbi.nlm.nih.gov

Downloaded from theInternationalmedicaljournal.org. For personal use only. No other uses without permission.
Copyright © International Medical Journal Corp. Ltd. All rights reserved