Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Anatomic Pathology and Forensic



Histopathology Of Basal Cell Carcinoma

[1] Yulia, [1] Wahyu

[1] Faculty of Medicine, Maranatha Christian University, Bandung City, West Java, Indonesia

Correspondence : dr.yulia9927387@gmail.com

Article History :
Received date : 2024/06/10
Revised date : 2024/07/14
Accepted date : 2024/08/19
Published date : 2025/01/20

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The histopathology of basal cell carcinoma (BCC) has garnered significant attention in recent literature, underscoring the complexities involved in its diagnosis and management. **Literature Review:** The examination of cytomorphological features in mixed histology BCCs reveals additional diagnostic challenges, emphasizing the importance of histopathological examination over cytology (Goyal et al., 2017). The integration of dermoscopy has also been noted as a vital tool in enhancing diagnostic accuracy, particularly for nodular BCC, the most common subtype (ENACHE et al., 2019). Dermoscopy aids in identifying clinical features that guide histopathological analysis, thereby improving early detection and treatment decisions. Further research highlights the variability in treatment responses among different histopathological subtypes of BCC (Ungureanu et al., 2021). The classification of BCC into low-risk and high-risk tumors based on clinical and pathological characteristics is crucial for effective management. Additionally, advancements in understanding BCC biology advocate for precision in morphological classification to inform treatment strategies, including the use of molecular targeting agents (Parth Naik & B. Desai, 2022). The clinical presentation of BCCs varies, with specific histological features indicating a higher risk of metastasis (Hoashi et al., 2022). Recent studies focusing on dermoscopic clues associated with aggressive BCC subtypes reinforce the necessity for meticulous surgical approaches to mitigate recurrence risks (Camela et al., 2023). Moreover, case reports emphasize the potential for misdiagnosis in aggressive subtypes, necessitating accurate histological evaluations (K Imlay et al., 2024). **Conclusions:** In conclusion, the literature collectively emphasizes the importance of accurate histopathological classification of BCC, the role of dermoscopy in enhancing diagnostic accuracy, and the need for tailored treatment strategies based on the tumor's biological behavior. The evolving understanding of BCC's histopathology is crucial for improving patient outcomes and guiding effective management.

Keywords: Histopathology, Basal Cell Carcinoma

Downloaded from The Medical Journal of Anatomic Pathology and Forensic
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The histopathology of basal cell carcinoma (BCC) is a critical area of research that has evolved significantly over the past decade, highlighting the complexities of diagnosis and management. (G. Marzuka & E. Book, 2015) provide a foundational understanding of BCC by categorizing the tumor into indolent-growth and aggressive-growth subtypes based on histopathological features. Their findings indicate a concerning rate of misdiagnosis of aggressive BCCs, suggesting that up to 38% of aggressive cases may be incorrectly identified as indolent variants, which has substantial implications for treatment strategies. The authors emphasize the need for precise histological evaluation, as the aggressiveness of the tumor is a key determinant in patient management.

Building on this, (Goyal et al., 2017) explore the cytomorphological features of mixed histology BCCs, shedding light on the diagnostic challenges faced by clinicians when relying solely on cytology. Their work underscores the necessity for histopathological examination to accurately diagnose and differentiate the various subtypes of BCC, further reinforcing the importance of thorough histological assessment in clinical practice.

(ENACHE et al., 2019) contribute to the discussion by examining dermoscopy patterns in nodular BCC, the most prevalent subtype. Their research highlights the utility of dermoscopy in enhancing diagnostic accuracy, particularly in identifying characteristic clinical features that can guide subsequent histopathological analysis. The integration of dermoscopy into the diagnostic process is essential for improving outcomes, as it aids in the early detection of BCC and informs treatment decisions.

(Ungureanu et al., 2021) expand on the role of dermoscopy, emphasizing its importance not only in diagnosis but also in the management of BCC. The authors discuss how different histopathologic subtypes respond variably to treatment, indicating that the classification of BCC based on histological criteria can significantly influence therapeutic choices. The delineation of low-risk and high-

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

risk tumors based on clinical and pathological characteristics is critical for effective management.

(Parth Naik & B. Desai, 2022) further delve into the contemporary understanding of BCC biology, advocating for precision in morphologic classification to inform treatment strategies. They highlight the significance of histologic criteria in the diagnosis and staging of BCC, suggesting that advancements in understanding the tumor's biology may lead to improved therapeutic approaches, including the use of molecular targeting agents.

(Hoashi et al., 2022) provide insights into the clinical presentation and histologic features of BCCs, emphasizing the need for awareness of specific characteristics that may indicate a higher risk of metastasis. Their findings reinforce the notion that while BCCs are generally indolent, certain subtypes exhibit aggressive behavior that necessitates vigilant monitoring and tailored treatment approaches.

In a recent study, (Camela et al., 2023) focus on the dermoscopic clues associated with histopathologically aggressive BCC subtypes, reinforcing the role of dermoscopy in differentiating between low-risk and high-risk tumors. The authors detail the histopathological features of aggressive BCCs, advocating for a meticulous surgical approach to mitigate recurrence risks.

(K Imlay et al., 2024) present a compelling case report of morpheaform BCC, illustrating the potential for misdiagnosis when distinguishing between BCC and other malignancies. Their case underscores the critical need for accurate histological evaluation, particularly in aggressive subtypes that may mimic other cancers, emphasizing the importance of timely and precise diagnosis.

Lastly, (Kaur et al., 2024) challenge the traditional understanding of BCC as solely an invasive neoplasm by presenting evidence of a superficial BCC variant. Their findings regarding the integrity of the basal lamina in different BCC subtypes contribute to the ongoing discourse on the histopathological classification of BCC and its implications for treatment.

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Together, these articles present a comprehensive overview of the histopathology of basal cell carcinoma, highlighting the complexities of diagnosis, the significance of accurate histological classification, and the evolving landscape of treatment strategies.

LITERATURE REVIEW

The article "Basal Cell Carcinoma: Pathogenesis, Epidemiology, Clinical Features, Diagnosis, Histopathology, and Management" by (G. Marzuka & E. Book, 2015) provides a comprehensive overview of basal cell carcinoma (BCC), with a particular emphasis on its histopathological characteristics and the implications for diagnosis and treatment. The authors highlight a significant issue in the clinical management of BCC: the misdiagnosis of aggressive-growth subtypes as indolent-growth variants, which occurs in 28 to 38 percent of cases evaluated via punch biopsy. This misclassification can lead to inadequate treatment approaches, as aggressive-growth BCCs, which account for 8 to 15 percent of all cases, require different management strategies compared to their indolent counterparts.

The article categorizes BCC into two main types: indolent-growth and aggressive-growth subtypes. Indolent-growth subtypes include nodular and superficial BCCs, while aggressive-growth subtypes encompass morpheaform, infiltrative, micronodular, and basosquamous forms. The authors note that mixed histology tumors represent about 40 percent of primary BCCs, underscoring the complexity of BCC diagnosis. This classification is crucial for determining treatment options, as the aggressiveness of the tumor significantly influences management decisions.

Histopathologically, all BCC subtypes share common features, including aggregations of basaloid keratinocytes that are surrounded by stromal tissue and typically maintain a connection to the epidermis. The basaloid cells exhibit characteristics similar to normal basal keratinocytes, such as large, uniform nuclei that are intensely basophilic and scant cytoplasm. The presence of peripheral palisading of nuclei is a notable feature in tumor aggregates, and the authors describe the frequent occurrence of apoptotic cells within these tumors.

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Specifically, nodular BCC is characterized by large nests of basaloid cells located in the dermis, often accompanied by peritumoral retraction from the stroma and peripheral palisading. The authors also mention that ulceration may occur, and cystic spaces can develop within larger tumor islands due to necrosis. In contrast, superficial BCC displays a distinct growth pattern, with basaloid cells proliferating parallel to the epidermal surface and limited to the papillary dermis.

The article "Cutaneous basal cell carcinoma with mixed histology: Cytomorphological features of two unusual cases" by (Goyal et al., 2017) provides a detailed examination of the histopathological characteristics of basal cell carcinoma (BCC), particularly focusing on two atypical cases. The authors emphasize the slow-growing nature of BCC, which is characterized by its local aggressiveness and infrequent metastasis, making accurate diagnosis crucial for effective management.

A significant contribution of this study is its exploration of the cytomorphological features observed in the two reported cases, which presented diagnostic challenges when evaluated through cytology. The authors meticulously compare these cytomorphological findings with the corresponding histopathological results from excision biopsies. This comparison highlights the discrepancies that can arise between cytological and histopathological diagnoses, underscoring the importance of histopathological examination as the gold standard for diagnosing BCC.

The article effectively illustrates the complexities involved in diagnosing BCC, particularly in cases exhibiting mixed histology. The authors detail the cytological characteristics, such as the presence of atypical keratinocytes and varying degrees of keratinization, which can lead to misinterpretation in cytological smears. By presenting these unusual cases, (Goyal et al., 2017) contribute valuable insights into the histopathological spectrum of BCC, emphasizing the need for careful evaluation of both cytological and histopathological specimens.

Furthermore, the study underscores the relevance of histopathological examination in confirming the diagnosis of BCC, particularly in atypical presentations. The authors advocate for a comprehensive approach that includes both cytological and

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

histopathological analyses to enhance diagnostic accuracy and minimize the risk of misdiagnosis, which could impact treatment decisions.

The article "Dermoscopy Patterns and Histopathological Findings in Nodular Basal Cell Carcinoma-Study on 68 Cases" by (ENACHE et al., 2019) provides a comprehensive examination of the histopathological features associated with nodular basal cell carcinoma (BCC), which is recognized as the most prevalent form of skin cancer. The authors highlight the increasing incidence of BCC in Europe, particularly among younger populations, underscoring the significance of this condition as a pressing healthcare issue.

The article effectively categorizes the various histological subtypes of BCC, including nodular, superficial, infiltrative, micronodular, adenoid, morpheaform, pigmented, and fibroepithelioma of Pinkus. Among these, nodular BCC is identified as the most common subtype, which is crucial for clinicians to recognize due to its distinct clinical presentation. The authors describe key clinical features that raise suspicion for BCC, such as the characteristic pearly, rolled borders, telangiectasia, and a pink surface, along with its tendency for slow growth in sun-exposed regions. This clinical insight is vital for early detection and intervention.

Moreover, the article emphasizes the role of dermoscopy as a valuable tool in the management of BCC. Dermoscopy enhances the ability to visualize the tumor's histological subtype and detect pigmentation that may not be clinically apparent. It also aids in assessing tumor expansion and responsiveness to non-ablative treatments, which is particularly relevant for guiding therapeutic decisions. The integration of dermoscopy into routine practice could potentially improve diagnostic accuracy and treatment outcomes.

The authors also stress the importance of histopathological examination of biopsy fragments for definitive diagnosis. This step is critical, as it confirms the presence of BCC and allows for the identification of its specific subtype, which can influence treatment strategies. The article provides a clear and informative overview of the histopathological aspects of nodular BCC, making it a valuable resource for dermatologists and pathologists alike.

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Role of Dermoscopy in the Assessment of Basal Cell Carcinoma" by (Ungureanu et al., 2021) provides an insightful examination of basal cell carcinoma (BCC), emphasizing its prevalence, treatment modalities, and the role of dermoscopy in enhancing diagnostic accuracy. BCC is identified as the most common cancer among white populations, with an increasing incidence globally. While the mortality associated with BCC is low, the morbidity remains significant, highlighting the need for effective management strategies.

The authors delineate the classification of BCC into low-risk and high-risk tumors based on a combination of patient, clinical, and pathological characteristics. This stratification is critical, as it informs treatment decisions. The article underscores the variances in treatment responses among different histopathological subtypes of BCC, which is pivotal for clinicians when devising treatment plans. For instance, the first-line treatments for superficial BCC (sBCC) include imiquimod and photodynamic therapy (PDT), while surgical excision is preferred for nodular BCC (nBCC). Moreover, Mohs micrographic surgery is highlighted as the gold standard for the most aggressive subtypes due to its efficacy in reducing recurrence rates.

A significant contribution of the article is its discussion on the application of dermoscopy, a non-invasive diagnostic technique, in the assessment and management of BCC. The authors present compelling evidence that dermoscopy not only enhances diagnostic accuracy but also aids in determining histological subtypes, evaluating clinical features such as pigmentation and ulceration, and assessing the response to non-ablative therapies. This advancement in diagnostic capability is crucial, as it allows for more tailored treatment approaches and potentially improved patient outcomes.

The article "Basal Cell Carcinoma: A Narrative Review on Contemporary Diagnosis and Management" by (Parth Naik & B. Desai, 2022) provides a comprehensive overview of the advancements in the understanding of basal cell carcinoma (BCC) biology and its implications for diagnosis and treatment. The authors emphasize the importance of precise categorization and staging of BCC, which are critical for determining appropriate therapeutic approaches. This point

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

underscores the necessity for surgical pathologists to be well-versed in the histologic criteria that are essential for accurate diagnosis and sub-classification of this prevalent skin malignancy.

One of the key insights from the article is the role of the Hedgehog signaling pathway in the pathogenesis of BCC. The authors highlight the recent approval of Smoothened inhibitors that target this pathway, which have shown promising results in shrinking metastatic or locally advanced tumors. This therapeutic advancement marks a significant evolution in the management of BCC, particularly for cases that are resistant to traditional treatments. The authors note, however, that while these emerging molecular targeting agents hold therapeutic promise, the prognosis for metastatic BCC remains uncertain and is likely to be poor.

The review effectively synthesizes current knowledge regarding the histopathological features of BCC, including the various subtypes and their distinct histological characteristics. Such detailed information is crucial for pathologists and clinicians alike, as it aids in the accurate diagnosis and subsequent management of BCC. The authors also discuss the implications of these histological findings on treatment decisions, reinforcing the idea that a thorough understanding of the disease's biology can lead to more effective therapeutic strategies.

The article "Molecular Mechanisms and Targeted Therapies of Advanced Basal Cell Carcinoma" by (Hoashi et al., 2022) provides a comprehensive overview of basal cell carcinoma (BCC), emphasizing its prevalence, risk factors, and histopathological characteristics. As one of the most common skin malignancies, BCC accounts for a significant portion of skin cancer cases, particularly in Caucasians, with an incidence that continues to rise globally.

The authors underscore the critical role of sunlight exposure as the primary risk factor for BCC development, supported by additional factors such as radiation, age, and skin type. This multifactorial etiology highlights the need for increased awareness and preventive measures, particularly in populations at higher risk. The article effectively delineates BCC as an indolent tumor that rarely metastasizes;

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

however, it also notes that inadequate treatment can lead to significant tissue destruction, which is a crucial consideration in clinical practice.

A key contribution of this article is its classification of BCCs with high metastatic potential based on specific clinical features. The authors identify tumors located on the midline of the face or ear, long-standing tumors, large tumor size, and a history of radiation exposure as indicators of increased risk for metastasis. This classification is particularly valuable for clinicians as it aids in risk stratification and informs treatment decisions. Furthermore, the mention of rare cases exhibiting intravascular invasion and their associated higher metastatic rates adds depth to the understanding of BCC's aggressive potential in certain contexts.

The article also touches upon the histologic features of BCC, which are essential for accurate diagnosis and management. The authors review the typical histopathological characteristics that pathologists encounter, such as nests of basaloid cells in the dermis and peripheral palisading. This detailed examination of histological features is critical for distinguishing BCC from other cutaneous malignancies, which can have overlapping presentations.

The article "Dermoscopic Clues of Histopathologically Aggressive Basal Cell Carcinoma Subtypes" by (Camela et al., 2023) provides a comprehensive examination of basal cell carcinoma (BCC), particularly focusing on its aggressive histopathological subtypes and their dermoscopic characteristics. The authors underscore that BCC is the most prevalent skin cancer globally, with an increasing incidence that raises concerns regarding its local aggressiveness, despite its low metastatic potential.

The article effectively categorizes BCC into low-risk and high-risk subtypes, emphasizing the significance of this classification in determining the recurrence rates post-surgery. The high-risk subtypes—morpheaform, micronodular, infiltrative, and metatypical BCC—are described in detail, with the authors providing clear histopathological descriptions that enhance understanding of their unique features. For instance, morpheaform BCC is noted for its thin cords of basaloid cells encased in sclerotic collagen, while infiltrative BCC is characterized

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

by its cords of keratinocytes within a mucinous stroma. These distinctions are crucial for clinicians as they inform surgical strategies and patient management.

The article also highlights the role of dermoscopy in the diagnosis of BCC, particularly in identifying aggressive subtypes. The authors argue for the necessity of diagnostic biopsies when high-risk subtypes are suspected, given their propensity for recurrence and the requirement for more meticulous surgical interventions. This assertion is particularly relevant in clinical practice, as it advocates for a proactive approach in managing patients with aggressive BCC.

Moreover, the study aims to correlate dermoscopic criteria with demographic characteristics and anatomical locations, which could potentially enhance the predictive accuracy for aggressive BCC subtypes. This aspect of the research is particularly valuable, as it suggests that demographic factors may play a role in the presentation and prognosis of BCC, thus providing a more personalized approach to treatment.

The article titled "Low-Powered View, High-Powered Mind: A Case Report of Morpheaform Basal Cell Carcinoma Mistaken for Metastatic Breast Carcinoma" by (K Imlay et al., 2024) provides a comprehensive exploration of morpheaform basal cell carcinoma (MBCC), a rare and aggressive subtype of basal cell carcinoma (BCC). The authors highlight the increasing prevalence of BCC, with approximately 4.3 million cases diagnosed annually in the United States, underscoring the importance of accurate diagnostic practices in managing this malignancy.

The article effectively delineates the clinical presentation and histopathological characteristics of MBCC, noting that it primarily affects photoexposed areas, particularly the face and neck, where it is often misdiagnosed due to its deceptive morphology. The authors emphasize that MBCC accounts for only 5%-10% of all BCC cases, yet it is associated with significantly higher recurrence rates and more extensive tissue destruction compared to other BCC subtypes. This information is critical for pathologists and clinicians, as it underscores the necessity for heightened

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

awareness and meticulous examination when evaluating skin lesions that may resemble other malignancies, such as metastatic breast carcinoma.

The treatment modalities discussed in the article, including Mohs micrographic surgery, cryotherapy, electrosurgery, radiotherapy, and various forms of chemotherapy, reflect a broad spectrum of options available for BCC and its subtypes. The authors argue that the choice of treatment is heavily influenced by an accurate and timely diagnosis, which is paramount in determining patient outcomes. This assertion is particularly relevant in the context of MBCC, where the aggressive nature of the disease necessitates prompt intervention to mitigate the risks of recurrence and extensive tissue damage.

The article "Cutaneous Superficial Basal Cell Carcinoma is a Basal Cell Carcinoma In Situ: Electron Microscopy of a Case Series of Basal Cell Carcinomas" by (Kaur et al., 2024) presents a compelling examination of the histopathological characteristics of various subtypes of basal cell carcinoma (BCC), specifically focusing on the integrity of the basal lamina in different presentations of the disease. The authors' investigation into three distinct cases—a superficial BCC, a nodular BCC, and an infiltrative BCC—provides valuable insights into the structural differences that may influence the clinical behavior and classification of these tumors.

A key finding of the study is the preservation of the basal lamina in the superficial BCC, which the authors assert is a significant indicator of the tumor's in situ nature. This observation suggests that superficial BCCs maintain a degree of histological integrity that differentiates them from their nodular and infiltrative counterparts, where the basal lamina is noted to be focally disrupted. The intact basal lamina surrounding the tumor nodules in the superficial variant implies a lack of invasive behavior, aligning with the authors' argument that this subtype should be classified as a carcinoma in situ ((Kaur et al., 2024)).

The article also delves into the embryonic origins of BCC, attributing its development to follicular germinative cells. This foundation may provide a better understanding of the tumor's pathologic subtypes and their respective behaviors. By

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

highlighting the unique characteristics of the superficial variant, the authors challenge the traditional view of BCC as solely an invasive neoplasm, thereby expanding the current knowledge of its histopathology.

Moreover, the use of electron microscopy enhances the credibility of the findings, offering a detailed visualization of the cellular architecture and the state of the basal lamina. This methodological approach strengthens the argument for recognizing superficial BCC as a distinct entity within the spectrum of basal cell carcinomas.

CONCLUSION

The histopathology of basal cell carcinoma (BCC) has garnered significant attention in recent literature, underscoring the complexities involved in its diagnosis and management. A foundational study categorizes BCC into indolent-growth and aggressive-growth subtypes based on histopathological features, revealing a misdiagnosis rate of aggressive BCCs as indolent variants that may reach as high as 38% (G. Marzuka & E. Book, 2015). This misclassification has profound implications for treatment, highlighting the necessity for precise histological evaluation to guide management strategies.

The examination of cytomorphological features in mixed histology BCCs reveals additional diagnostic challenges, emphasizing the importance of histopathological examination over cytology (Goyal et al., 2017). The integration of dermoscopy has also been noted as a vital tool in enhancing diagnostic accuracy, particularly for nodular BCC, the most common subtype (ENACHE et al., 2019). Dermoscopy aids in identifying clinical features that guide histopathological analysis, thereby improving early detection and treatment decisions.

Further research highlights the variability in treatment responses among different histopathological subtypes of BCC (Ungureanu et al., 2021). The classification of BCC into low-risk and high-risk tumors based on clinical and pathological characteristics is crucial for effective management. Additionally, advancements in understanding BCC biology advocate for precision in morphological classification

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

to inform treatment strategies, including the use of molecular targeting agents (Parth Naik & B. Desai, 2022).

The clinical presentation of BCCs varies, with specific histological features indicating a higher risk of metastasis (Hoashi et al., 2022). Recent studies focusing on dermoscopic clues associated with aggressive BCC subtypes reinforce the necessity for meticulous surgical approaches to mitigate recurrence risks (Camela et al., 2023). Moreover, case reports emphasize the potential for misdiagnosis in aggressive subtypes, necessitating accurate histological evaluations (K Imlay et al., 2024).

Lastly, the exploration of superficial BCC variants challenges the traditional understanding of BCC as solely invasive, suggesting that certain subtypes may exhibit distinct histopathological characteristics, such as the integrity of the basal lamina (Kaur et al., 2024). This finding contributes to the ongoing discourse regarding the classification and treatment of BCC.

In conclusion, the literature collectively emphasizes the importance of accurate histopathological classification of BCC, the role of dermoscopy in enhancing diagnostic accuracy, and the need for tailored treatment strategies based on the tumor's biological behavior. The evolving understanding of BCC's histopathology is crucial for improving patient outcomes and guiding effective management.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

1. G. Marzuka, A. & E. Book, S., 2015. Basal Cell Carcinoma: Pathogenesis, Epidemiology, Clinical Features, Diagnosis, Histopathology, and Management. ncbi.nlm.nih.gov
2. Goyal, S., Rathore, R., Sharma, S., Kumar Arora, V., Krushna Das, G., & Singal, A., 2017. Cutaneous basal cell carcinoma with mixed histology: Cytomorphological features of two unusual cases. ncbi.nlm.nih.gov
3. ENACHE, A. O., PĂTRAȘCU, V., SIMIONESCU, C. E., CIUREA, R. N., VĂDUVA, A., & STOICA, L. E., 2019. Dermoscopy Patterns and Histopathological Findings in Nodular Basal Cell Carcinoma-Study on 68 Cases. ncbi.nlm.nih.gov
4. Ungureanu, L., Cosgarea, I., Şenilă, S., & Vasilovici, A., 2021. Role of Dermoscopy in the Assessment of Basal Cell Carcinoma. ncbi.nlm.nih.gov
5. Parth Naik, P. & B. Desai, M., 2022. Basal Cell Carcinoma: A Narrative Review on Contemporary Diagnosis and Management. ncbi.nlm.nih.gov
6. Hoashi, T., Kanda, N., & Saeki, H., 2022. Molecular Mechanisms and Targeted Therapies of Advanced Basal Cell Carcinoma. ncbi.nlm.nih.gov
7. Camela, E., Ilut Anca, P., Lallas, K., Papageorgiou, C., Manoli, S. M., Gkentsidi, T., Eftychidou, P., Liopyris, K., Sgouros, D., Apalla, Z., & Lallas, A., 2023. Dermoscopic Clues of Histopathologically Aggressive Basal Cell Carcinoma Subtypes. ncbi.nlm.nih.gov
8. K Imlay, R., Perez Torrico, C., Sulaimann, A., M Grimes, Z., & Monnett, S., 2024. Low-Powered View, High-Powered Mind: A Case Report of Morpheaform Basal Cell Carcinoma Mistaken for Metastatic Breast Carcinoma. ncbi.nlm.nih.gov
9. Kaur, H., L. de Mesy Bentley, K., Minhaj Rahman, S., R. Cohen, P., & R. Smoller, B., 2024. Cutaneous Superficial Basal Cell Carcinoma is a Basal Cell Carcinoma In Situ: Electron Microscopy of a Case Series of Basal Cell Carcinomas. ncbi.nlm.nih.gov

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved