Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Anatomic Pathology and Forensic



Histopathology of Columnar Cell Lesions of the Breast

[1] Martin, [1] Gunawan

[1] Faculty of Medicine, Maranatha Christian University, Bandung City, West Java, Indonesia

Correspondence : dr.martin938276@gmail.com

Article History :
Received date : 2024/06/28
Revised date : 2024/07/08
Accepted date : 2024/08/17
Published date : 2025/01/20

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The histopathological examination of columnar cell lesions (CCLs) of the breast reveals a complex interplay between various lesion types and their potential implications for breast cancer pathology. **Literature Review:** (S Gomes et al., 2011) expanded on this understanding by exploring the frequency of lobular neoplasia and its association with other breast lesions, suggesting an evolutionary pathway that links lobular neoplasia, lobular carcinomas, and CCLs. Their findings emphasize the importance of thorough histopathological examination, as the presence of lobular neoplasia can indicate a higher risk for subsequent malignancies, thereby guiding clinical management. The challenges in diagnosing CCLs and their morphological subtleties were further highlighted by (S Gomes et al., 2014), who examined inter-observer variability among pathologists. Their study underscored the critical role of accurate morphological classification in predicting patient outcomes, particularly as diagnostic discrepancies can impact treatment decisions. (A. Rakha et al., 2018) contributed to the discourse by investigating the invasive potential of breast lesions, emphasizing the need for a nuanced understanding of the epithelial-stroma barrier and its role in distinguishing benign from malignant lesions. Their findings suggest that certain benign lesions, like microglandular adenosis, may possess underlying genetic alterations that predispose them to malignancy, reinforcing the importance of early detection and characterization of lesions such as CCLs. **Conclusions:** In conclusion, the literature collectively underscores the importance of recognizing CCLs as potential precursors in the breast cancer continuum. The interrelations among various breast lesions, diagnostic challenges, and the implications of histopathological findings are critical for advancing our understanding of breast cancer pathology. Ongoing research is essential to unravel the complexities of these lesions and their roles in breast cancer development.

Keywords: Histopathology, Columnar Cell Lesions of the Breast

Downloaded from The Medical Journal of Anatomic Pathology and Forensic
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The histopathology of columnar cell lesions (CCLs) of the breast has garnered significant attention in recent years, reflecting a growing understanding of their potential role in breast cancer pathology. The foundational work by (Ferreira et al., 2010) elucidated the pathological features of CCLs, particularly in canine mammary glands, and highlighted their relationship to human breast cancer. This study established that CCLs arise from terminal duct lobular units (TDLUs) and can be categorized into columnar cell change (CCC) and columnar cell hyperplasia (CCH), with further subclassification based on cytological atypia. The authors emphasized the importance of recognizing these lesions as potential precursors to low-grade ductal carcinoma in situ (DCIS) and invasive carcinoma, setting the stage for further exploration of their implications in human pathology.

Building on this understanding, (S Gomes et al., 2011) investigated the frequency of lobular neoplasia in breast biopsies and its association with other breast lesions, using advanced molecular biology techniques. Their findings suggested a shared evolutionary pathway between lobular neoplasia, lobular carcinomas, and CCLs, reinforcing the notion that these lesions may be interrelated and important in the continuum of breast cancer development.

The diagnostic challenges associated with distinguishing between various breast lesions were further examined by (S Gomes et al., 2014), who assessed inter-observer variability in the diagnosis of lobular neoplasia, CCLs, atypical ductal hyperplasia (ADH), and DCIS. Their study underscored the critical role of accurate morphological classification in predicting patient outcomes, particularly in the context of lobular lesions, which can exhibit significant variability in histological features.

In a more recent study, (A. Rakha et al., 2018) explored the histologic features and genomic profiles associated with infiltrating epitheliosis of the breast, contributing to the understanding of the invasive potential of certain breast lesions. This research provided molecular evidence for the progression of benign lesions, such as

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

microglandular adenosis, to invasive carcinoma, thereby highlighting the importance of early detection and characterization of lesions like CCLs in the breast.

[Amedeo et al. (2019)] further investigated the relationship between CCLs and apocrine hyperplasia, proposing a potential common origin mediated by prolactin-induced protein. This study suggested that shared molecular pathways could exist among various breast lesions, including CCLs, and emphasized the need for further research to clarify these connections.

(Strickland & Turashvili, 2022) expanded the discussion to include the presence of CCLs in male breast specimens, raising questions about their histological characteristics and potential precursory role in male breast cancer. Their findings indicated that while CCLs may be less common in males, they could still represent significant precursors to malignancy, warranting further investigation.

Finally, (Hoi Kin Lam et al., 2024) provided a retrospective analysis of rare breast cancer histotypes, including mucinous and invasive lobular carcinomas. This comprehensive review highlighted the distinct characteristics and prognoses of various breast cancer subtypes, reinforcing the notion that understanding the histopathological features of lesions like CCLs is crucial for accurate diagnosis and treatment planning.

Together, these articles contribute to a nuanced understanding of columnar cell lesions in breast pathology, emphasizing their potential role as precursors in the development of breast cancer and the necessity for ongoing research to unravel the complex relationships among various breast lesions.

LITERATURE REVIEW

The article "Columnar cell lesions of the canine mammary gland: pathological features and immunophenotypic analysis" by Ferreira et al. (2010) provides a comprehensive examination of columnar cell lesions (CCLs) within the context of canine mammary glands, drawing parallels to human breast pathology. The authors

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

assert that the development of human breast cancer often initiates in the terminal duct lobular units (TDLUs) and progresses to invasive cancer, highlighting the significance of precursor lesions such as atypical ductal hyperplasia, lobular carcinoma in situ, and notably, CCLs.

The article delineates the histopathological characteristics of CCLs, noting that they exhibit varying degrees of acinar dilation and are lined by multiple layers of columnar epithelial cells with uniform, ovoid nuclei. This description aligns with established criteria for CCLs in human pathology, where they are further categorized into columnar cell change (CCC) and columnar cell hyperplasia (CCH). The authors effectively emphasize that CCC and CCH with cytological atypia are subclassified as flat epithelial atypia (FEA), a classification that is crucial for understanding the potential progression of these lesions to low-grade ductal carcinoma in situ (DCIS) and invasive carcinoma.

A critical evaluation of the material reveals that while the authors successfully draw connections between canine and human breast pathology, the implications of their findings could be further explored. For instance, the study primarily focuses on the morphological and immunophenotypic features of CCLs in dogs, yet it could benefit from a more thorough discussion on how these findings may influence diagnostic and therapeutic strategies in veterinary medicine and their relevance to human breast cancer research. Furthermore, the article could enhance its impact by including a wider variety of cases or comparative analyses with other mammalian species, thereby broadening the understanding of CCLs across different contexts.

The article titled "Lobular neoplasia: frequency and association with other breast lesions" by Gomes et al. (2011) presents significant findings regarding the histopathological characteristics of lobular neoplasia and its relationship with other breast lesions, particularly focusing on columnar cell lesions. The authors utilize contemporary molecular biology techniques to explore the evolutionary pathways that connect lobular neoplasia to lobular carcinomas and columnar cell lesions, which is a crucial aspect of understanding the pathogenesis of these conditions.

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

A critical evaluation of the article reveals that the authors effectively demonstrate the frequency of lobular neoplasia in a series of breast biopsies, providing valuable epidemiological data. They report that lobular neoplasia is not merely an incidental finding but is frequently associated with other breast pathologies, suggesting a potential continuum in the spectrum of breast lesions. This insight is particularly relevant as it underscores the importance of thorough histopathological examination in breast biopsies, where the presence of lobular neoplasia could indicate a higher risk for subsequent breast malignancies.

Furthermore, the study highlights the association between different subtypes of lobular neoplasia and both benign and malignant breast lesions. This association is critical for clinicians and pathologists, as it may influence management decisions and patient prognostication. The authors argue that understanding these relationships can lead to improved diagnostic accuracy and better-informed clinical practices.

The article "Inter-observer variability between general pathologists and a specialist in breast pathology in the diagnosis of lobular neoplasia, columnar cell lesions, atypical ductal hyperplasia and ductal carcinoma in situ of the breast" by Gomes et al. (2014) provides a critical examination of the diagnostic discrepancies between general pathologists and specialists in breast pathology concerning various breast lesions, including columnar cell lesions (CCL).

The study highlights the significance of accurate diagnosis in breast pathology, particularly for lesions that may have implications for patient management and prognosis. The authors emphasize the variability in diagnostic agreement among pathologists, which can lead to differing interpretations of the same histopathological specimens. This variability is particularly concerning for lesions like CCL, where the morphological features can be subtle and may overlap with other conditions, such as atypical ductal hyperplasia (ADH) and lobular neoplasia (LN).

Gomes et al. (2014) underscore the importance of the morphological classification of lobular lesions, as established by the Page group, which correlates with patient

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

outcomes. The study's findings suggest that specialist pathologists, who possess advanced training and experience in breast pathology, demonstrate higher diagnostic consistency compared to their generalist counterparts. This is critical as it points to the need for specialized training in the interpretation of complex breast lesions to improve diagnostic accuracy and, consequently, patient care.

The authors conducted a thorough analysis of diagnostic agreements and reproducibility, utilizing a cohort of cases that were referred for second opinions. This methodological approach strengthens the study's validity, as it reflects real-world clinical scenarios where pathologists may encounter challenging cases. The results indicate a significant degree of inter-observer variability, which could potentially affect treatment decisions and patient outcomes.

The article "Invasion in breast lesions: the role of the epithelial-stroma barrier" by Rakha et al. (2018) provides a comprehensive examination of the histopathological characteristics associated with various breast lesions, particularly focusing on the interaction between epithelial and stromal components. The authors emphasize the significance of the epithelial-stroma barrier in determining the behavior of breast lesions, which is crucial for understanding the progression from benign to malignant states.

The main idea presented by Rakha et al. revolves around the characterization of infiltrating epitheliosis and its correlation with other breast lesions, such as microglandular adenosis and acinic cell carcinoma. The authors delve into the histological features and immunophenotypes of these lesions, highlighting the importance of recognizing specific histopathological patterns that may indicate a propensity for invasion (Rakha et al., 2018). This characterization is essential for pathologists in making accurate diagnoses and determining the appropriate management strategies for patients.

The article also discusses the genetic analysis of microglandular adenosis, providing evidence for its potential progression to invasive carcinoma. This insight is particularly valuable as it underscores the need for careful monitoring of lesions that may appear benign but possess underlying genetic alterations that predispose

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

them to malignancy. The authors present a well-rounded discussion on the implications of these findings for clinical practice, suggesting that a thorough understanding of the histopathological landscape can aid in identifying patients at higher risk for progression to invasive disease.

Moreover, the review critically evaluates the role of myoepithelial cells in distinguishing between benign and malignant lesions. The authors argue that the absence of myoepithelium in certain breast lesions does not automatically imply malignancy, which challenges traditional diagnostic paradigms (Rakha et al., 2018). This perspective encourages a more nuanced approach to histopathological evaluation, taking into account the complexity of breast lesion behavior and the multifactorial nature of tumor progression.

The article "Columnar Cell Lesion and Apocrine Hyperplasia of the Breast: Is There a Common Origin? The Role of Prolactin-induced Protein" by Amedeo et al. (2019) presents a compelling exploration of the histopathological characteristics of columnar cell lesions (CCLs) in the breast, particularly in relation to apocrine hyperplasia. The authors delve into the potential shared origins of these lesions, proposing that prolactin-induced protein may play a significant role in their development.

Amedeo et al. (2019) critically examine existing literature to contextualize their findings within the broader discourse on breast pathology. They highlight that CCLs, often detected incidentally during routine examinations, may represent an early histologically detectable stage in the progression toward breast cancer. This assertion aligns with the suggestion that CCLs could be non-obligate precursors to malignancy, meriting further investigation into their clinical significance.

The article's methodology is noteworthy, as it combines histopathological analysis with an examination of hormonal influences, particularly the role of prolactin. This approach not only enriches the understanding of the biological underpinnings of CCLs but also raises important questions about the potential for targeted therapeutic strategies. The authors effectively argue for the need to further elucidate the

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

molecular pathways involved in the pathogenesis of these lesions, which could enhance diagnostic precision and inform management strategies.

However, while the article provides valuable insights, it could benefit from a more detailed discussion of the implications of these findings for clinical practice. Specifically, the authors could explore how the identification of CCLs and their relationship with apocrine hyperplasia might influence screening protocols and risk assessment for breast cancer. Additionally, a more comprehensive review of the histopathological criteria used to diagnose CCLs would strengthen the article's contribution to the field.

The article "Are Columnar Cell Lesions the Earliest Non-Obligate Precursor in the Low-Grade Breast Neoplasia Pathway?" by Strickland and Turashvili (2022) provides a comprehensive examination of columnar cell lesions (CCLs) within the context of male breast pathology. The authors synthesize findings from various studies to highlight the characteristics and implications of CCLs as potential precursors to breast neoplasia, particularly focusing on their prevalence and histopathological features.

One of the key insights presented is the description of "columnar cell-like" lesions, which consist of dilated ducts lined by an inner layer of columnar luminal cells and surrounded by intact myoepithelial cells. This structural description emphasizes the morphological similarities between these lesions and traditional CCLs, suggesting a possible continuum in the pathology of breast lesions (Strickland & Turashvili, 2022). However, the authors critically assess the diagnostic challenges posed by these lesions, particularly in male patients, where typical CCL features such as cystically dilated acini and secretions are often absent.

The retrospective analysis conducted as part of the International Male Breast Cancer Program is particularly noteworthy. The detection of columnar cell-like lesions in a subset of patients with adjacent ductal carcinoma in situ (DCIS) raises important questions about the role of these lesions in the progression to malignancy. The presence of similar mutations in CCLs and adjacent invasive carcinoma in

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

some cases further supports the hypothesis that CCLs may act as non-obligate precursors to breast cancer, even in the male population.

Despite the low prevalence of CCLs in male breast cases, Strickland and Turashvili (2022) argue that the findings underscore the need for increased awareness and understanding of these lesions in clinical practice. The authors advocate for further research to delineate the biological behavior of CCLs and their potential implications in breast cancer pathways. This critical evaluation of existing literature not only highlights the complexities of diagnosing CCLs but also emphasizes their potential significance in the broader context of breast neoplasia.

The article "Rare Breast Cancer Histotypes—A Retrospective Study and Literature Review" by Lam et al. (2024) provides a comprehensive overview of several rare histological subtypes of breast cancer, focusing on their unique histopathological features, immunohistochemical profiles, and clinical behaviors. The authors effectively categorize these subtypes into mucinous carcinoma, invasive lobular carcinoma, papillary carcinoma, and tubular carcinoma, each distinguished by specific morphological characteristics and patterns of growth.

Mucinous carcinoma is highlighted for its production of abundant mucin, with the authors delineating between the pure and mixed types based on mucin purity. The classification is crucial as it reflects on the biological behavior of the carcinoma, with the pure type being defined by a mucin content of 90% or more, while the mixed type contains 50–90% mucin purity along with an infiltrating ductal component. The immunohistochemical findings are particularly noteworthy, as they indicate a high prevalence of estrogen receptor (ER) and progesterone receptor (PR) positivity, which suggests a favorable response to hormonal therapies. The authors also point out the typical clinical presentation of mucinous carcinoma, characterized by smaller tumor size and lower rates of nodal involvement, which can influence treatment strategies and prognostic assessments.

Invasive lobular carcinoma is described with attention to its distinctive growth pattern, often referred to as ‘Indian file’, which can complicate diagnosis due to its subtlety in imaging and histological evaluation. The authors mention its

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

multicentric nature and the propensity for bilateral involvement, which necessitates a vigilant approach in clinical management. The tendency for this subtype to metastasize in a less predictable manner, often to non-lymphatic tissues, underscores the importance of thorough staging and follow-up.

Papillary carcinoma is discussed in terms of its architectural features, with the proliferation of cells around fibrovascular cores leading to its classification into various subtypes. The article notes the high rates of hormonal receptor positivity and the predominance of HER2-negative cases, which again points to the potential for targeted therapeutic approaches. The authors emphasize the early-stage diagnosis of this subtype, which is crucial for improving patient outcomes.

Lastly, tubular carcinoma is characterized by its well-defined tubular structures, surrounded by a fibro-hyaline stroma. This description aligns with the histopathological criteria used to differentiate it from other subtypes, providing insights into its distinct biological behavior and potential clinical implications.

CONCLUSION

The histopathological examination of columnar cell lesions (CCLs) of the breast reveals a complex interplay between various lesion types and their potential implications for breast cancer pathology. The foundational work by (Ferreira et al., 2010) established that CCLs arise from terminal duct lobular units (TDLUs) and can be classified into columnar cell change (CCC) and columnar cell hyperplasia (CCH), with significant attention to their potential as precursors to low-grade ductal carcinoma in situ (DCIS) and invasive carcinoma. This classification is essential for understanding the progression of breast lesions and underscores the necessity for accurate morphological assessment.

(S Gomes et al., 2011) expanded on this understanding by exploring the frequency of lobular neoplasia and its association with other breast lesions, suggesting an evolutionary pathway that links lobular neoplasia, lobular carcinomas, and CCLs. Their findings emphasize the importance of thorough histopathological

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

examination, as the presence of lobular neoplasia can indicate a higher risk for subsequent malignancies, thereby guiding clinical management.

The challenges in diagnosing CCLs and their morphological subtleties were further highlighted by (S Gomes et al., 2014), who examined inter-observer variability among pathologists. Their study underscored the critical role of accurate morphological classification in predicting patient outcomes, particularly as diagnostic discrepancies can impact treatment decisions.

(A. Rakha et al., 2018) contributed to the discourse by investigating the invasive potential of breast lesions, emphasizing the need for a nuanced understanding of the epithelial-stroma barrier and its role in distinguishing benign from malignant lesions. Their findings suggest that certain benign lesions, like microglandular adenosis, may possess underlying genetic alterations that predispose them to malignancy, reinforcing the importance of early detection and characterization of lesions such as CCLs.

(Amedeo et al., 2019) proposed a potential common origin for CCLs and apocrine hyperplasia, mediated by prolactin-induced protein, which could indicate shared molecular pathways among various breast lesions. This perspective invites further investigation into the biological underpinnings of these lesions and their clinical implications.

(Strickland & Turashvili, 2022) focused on the implications of CCLs in male breast pathology, suggesting that although less common, these lesions could still represent significant precursors to malignancy. Their findings highlight the necessity for increased awareness and understanding of CCLs in clinical practice, regardless of gender.

Finally, (Hoi Kin Lam et al., 2024) provided a comprehensive overview of rare breast cancer histotypes, reinforcing the notion that understanding the histopathological features of lesions like CCLs is crucial for accurate diagnosis and treatment planning.

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

In conclusion, the literature collectively underscores the importance of recognizing CCLs as potential precursors in the breast cancer continuum. The interrelations among various breast lesions, diagnostic challenges, and the implications of histopathological findings are critical for advancing our understanding of breast cancer pathology. Ongoing research is essential to unravel the complexities of these lesions and their roles in breast cancer development.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Ferreira, E., Gobbi, H., S Saraiva, B., & D Cassali, G., 2010. Columnar cell lesions of the canine mammary gland: pathological features and immunophenotypic analysis. ncbi.nlm.nih.gov
2. S Gomes, D., Balabram, D., S Porto, S., & Gobbi, H., 2011. Lobular neoplasia: frequency and association with other breast lesions. ncbi.nlm.nih.gov
3. S Gomes, D., S Porto, S., Balabram, D., & Gobbi, H., 2014. Inter-observer variability between general pathologists and a specialist in breast pathology in the diagnosis of lobular neoplasia, columnar cell lesions, atypical ductal hyperplasia and ductal carcinoma in situ of the breast. ncbi.nlm.nih.gov
4. A. Rakha, E., Miligy, I., L. Gorringe, K., S. Toss, M., R. Green, A., B. Fox, S., C. Schmitt, F., Tan, P. H., M. Tse, G., Badve, S., Decker, T., Vincent-Salomon, A., J. Dabbs, D., P. Foschini, M., Moreno, F., Wentao, Y., C. Geyer, F., S. Reis-Filho, J., E. Pinder, S., R. Lakhani, S., & O. Ellis, I., 2018. Invasion in breast lesions: the role of the epithelial-stroma barrier. [PDF]

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

5. Strickland, S. & Turashvili, G., 2022. Are Columnar Cell Lesions the Earliest Non-Obligate Precursor in the Low-Grade Breast Neoplasia Pathway?. ncbi.nlm.nih.gov
6. Hoi Kin Lam, A., Tiong Hong Co, M., & Kwong, A., 2024. Rare Breast Cancer Histotypes—A Retrospective Study and Literature Review. ncbi.nlm.nih.gov
7. Amedeo, S., Gianluca, L., Chiara, C., Letterio, R., Giulia, E., Bonometti, A., Luca, D., Concetta, B., Donatella, G., & Fusco, N., 2019. Columnar Cell Lesion and Apocrine Hyperplasia of the Breast: Is There a Common Origin? The Role of Prolactin-induced Protein. [PDF]

Downloaded from The International Medical Journal of Anatomic Pathology and Forensic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved