Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Clinical Nutrition and Family Medicine



Effect of Vitamin D Supplementation on Blood Pressure

[1] Rahmawati, [1] Andi

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.rahmawati93847@gmail.com

Article History :

Received date : 2024/10/12
Revised date : 2024/11/27
Accepted date : 2024/12/15
Published date : 2025/01/08

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The relationship between vitamin D supplementation and blood pressure has garnered significant attention in the academic literature, reflecting a growing interest in the potential role of vitamin D beyond its established benefits for bone health. **Literature Review:** Several articles contribute to the discourse, indicating that while observational studies have established a correlation between vitamin D levels and blood pressure, interventional studies yield mixed results. For instance, high-dose vitamin D supplementation showed no significant effect on blood pressure in resistant hypertension patients (Pawlicka et al., 2018), while systematic reviews indicate potential blood pressure reduction in vitamin D-deficient populations (Farapti et al., 2020). The nuanced relationship is further complicated by the mechanisms proposed for vitamin D's influence on blood pressure, such as its effects on the renin-angiotensin-aldosterone system (RAAS) and endothelial function [ref: 288f4604-dcd6-4d1e-9a52-dea38e60821a; ref: 7b5cd7dd-0262-472f-bf1b-7a0aba5e1b28]. Moreover, the NHANES study data suggest that optimal vitamin D levels may mitigate age-related increases in systolic blood pressure, although vitamin D levels do not predict future hypertension (Che et al., 2023). This highlights the importance of individual patient factors, such as baseline vitamin D status and demographic variables, in understanding the effects of supplementation on blood pressure regulation. **Conclusions:** In conclusion, the reviewed literature indicates a need for more rigorous, well-designed randomized controlled trials to establish causality and clarify the mechanisms through which vitamin D may influence blood pressure. While there is evidence suggesting that vitamin D supplementation may lower blood pressure, particularly in deficient populations, inconsistencies in study outcomes necessitate further investigation to develop definitive guidelines for vitamin D supplementation in hypertension management.

Keywords: Vitamin D Supplementation ,Blood Pressure

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The relationship between vitamin D supplementation and blood pressure has garnered significant attention in the academic literature, reflecting a growing interest in the potential role of vitamin D beyond its established benefits for bone health. (Kienreich et al., 2013) highlight that vitamin D deficiency is prevalent and linked to various chronic diseases, including cardiovascular and renal conditions. Their review emphasizes the need for further investigation into whether vitamin D deficiency is a causative factor or merely a consequence of these diseases, particularly arterial hypertension and other cardiovascular risk factors.

In the same vein, (Kienreich et al., 2013) further explore the association between vitamin D deficiency and elevated blood pressure levels. They note that while observational studies suggest a correlation, the evidence is complicated by potential residual confounding. Their findings indicate that vitamin D may influence blood pressure through mechanisms involving the renin-angiotensin-aldosterone system (RAAS) and secondary hyperparathyroidism, suggesting a nuanced relationship that warrants further exploration through randomized controlled trials (RCTs).

(Yun Jeong et al., 2017) expand upon these insights by examining the pleiotropic effects of vitamin D across various organ systems. They point out that despite a significant body of evidence linking vitamin D deficiency with hypertension, the results from interventional studies remain inconsistent. This inconsistency underscores the importance of critically assessing the existing literature to clarify the role of vitamin D in blood pressure regulation.

(Pawlicka et al., 2018) contribute to this discourse by reporting on high-dose vitamin D supplementation in patients with resistant hypertension, finding no significant impact on blood pressure or other cardiovascular markers. Their results challenge the notion of vitamin D as a straightforward remedy for hypertension,

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

emphasizing the complexity of the relationship and the need for further rigorous investigation.

(Farapti et al., 2020) conduct a systematic review and meta-analysis, concluding that vitamin D supplementation may indeed lower blood pressure, particularly in vitamin D-deficient populations. They highlight the necessity for placebo-controlled RCTs to establish causality, as previous observational studies suggest a correlation but do not definitively prove a cause-and-effect relationship.

Finally, (Che et al., 2023) analyze data from the NHANES study, revealing that optimal vitamin D levels may mitigate age-related increases in systolic blood pressure, particularly among white Americans. Their findings further reinforce the association between vitamin D status and blood pressure regulation, while also acknowledging that vitamin D levels do not predict future hypertension.

Together, these articles illustrate a complex interplay between vitamin D supplementation and blood pressure, with evidence supporting both beneficial effects and inconsistencies across different populations and study designs. The literature calls for more targeted research to clarify these relationships and establish definitive guidelines for vitamin D supplementation in the context of hypertension management.

LITERATURE REVIEW

The article titled "Vitamin D, arterial hypertension & cerebrovascular disease" by (Kienreich et al., 2013) provides a comprehensive overview of the role of vitamin D in relation to arterial hypertension and its potential implications for cardiovascular health. The authors begin by highlighting the well-established benefits of vitamin D supplementation in preventing rickets in children and treating osteoporosis, noting its effectiveness in reducing fractures and falls. However, they pivot to the growing body of research that suggests a more complex relationship between vitamin D levels and extra-skeletal diseases, particularly cardiovascular and renal conditions.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

One of the critical insights from the article is the discussion of epidemiological studies that have identified associations between vitamin D deficiency and an increased risk of cardiovascular diseases, including arterial hypertension. The authors emphasize that while these studies suggest a correlation, the nature of the relationship—whether vitamin D deficiency is a causative factor or a consequence of these diseases—remains ambiguous. This point is crucial as it underscores the need for further research to clarify the underlying mechanisms that may link vitamin D status with hypertension and other cardiovascular risk factors.

(Kienreich et al., 2013) also summarize various studies that have explored the potential mechanisms by which vitamin D may influence blood pressure regulation. They discuss the role of vitamin D in the renin-angiotensin system, which is integral to blood pressure control, and how vitamin D's anti-inflammatory properties might contribute to cardiovascular health. However, the authors note that while some interventional studies have shown promising results regarding the effects of vitamin D supplementation on blood pressure, the evidence is not yet conclusive.

The review calls attention to the need for well-designed clinical trials to establish a causal relationship between vitamin D supplementation and blood pressure reduction. The authors argue that understanding this relationship is vital, as it could have significant implications for public health strategies aimed at preventing hypertension and its associated complications.

The article "Vitamin D and Cardiovascular Disease" by (Kienreich et al., 2013) provides a comprehensive examination of the relationship between vitamin D levels and blood pressure, particularly in the context of cardiovascular health. The authors highlight that vitamin D deficiency is associated with elevated blood pressure levels, suggesting that adequate vitamin D status may play a protective role against hypertension.

A critical evaluation of the article reveals several key insights into the mechanisms underlying this association. The authors discuss the role of the renin-

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

angiotensin-aldosterone system (RAAS), noting that vitamin D may inversely affect RAAS activity. This is significant because an overactive RAAS is a well-known contributor to hypertension. Furthermore, the article emphasizes the improvement of endothelial function as another potential mechanism, suggesting that vitamin D may enhance vascular health and thereby lower blood pressure.

The article also addresses the impact of parathyroid hormone (PTH) levels, which are known to rise in conditions of vitamin D deficiency. Elevated PTH is associated with myocardial hypertrophy and increased blood pressure, indicating that vitamin D's role in regulating PTH levels could be crucial for maintaining optimal blood pressure. The authors present data from a meta-analysis that demonstrates a significant inverse relationship between baseline serum vitamin D levels and the risk of developing hypertension, quantifying this relationship with a reported 12% decrease in hypertension risk for every 10 ng/mL increase in 25(OH)D.

Despite these findings, the article notes variability in the results of randomized controlled trials (RCTs) assessing vitamin D supplementation and blood pressure outcomes. Some studies reported non-significant reductions in blood pressure among hypertensive patients receiving vitamin D, while others indicated significant reductions in blood pressure specifically among those with vitamin D insufficiency or deficiency. For instance, one trial highlighted that black participants experienced a decrease in systolic blood pressure correlating with increases in vitamin D levels, underscoring the potential for demographic factors to influence the efficacy of supplementation.

The article "Vitamin D and Hypertension" by (Yun Jeong et al., 2017) provides a comprehensive examination of the relationship between vitamin D levels and hypertension, highlighting the multifaceted roles of vitamin D beyond its traditional association with bone health. The authors emphasize the importance of understanding vitamin D's pleiotropic effects, notably its role in cardiovascular health, which is increasingly recognized in contemporary research.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article begins by outlining the fundamental physiological functions of vitamin D, particularly its impact on the absorption of essential minerals such as calcium, magnesium, and phosphate. This foundational knowledge sets the stage for discussing vitamin D's broader implications for various organ systems, including the cardiovascular system. The identification of vitamin D receptors in most human tissues underscores its systemic significance, suggesting that vitamin D may influence blood pressure regulation through multiple pathways.

The authors discuss the alarming prevalence of hypertension, often referred to as a "silent killer" due to its asymptomatic nature. They reference a range of observational studies that have established a correlation between vitamin D deficiency and increased risk of cardiovascular diseases, particularly hypertension. This correlation is critical, as it suggests that maintaining adequate vitamin D levels may be a preventive strategy against hypertension.

Despite the compelling observational data, (Yun Jeong et al., 2017) note the inconsistencies in the results of interventional studies examining vitamin D supplementation's effect on blood pressure. This inconsistency raises questions about the efficacy of vitamin D supplementation as a therapeutic approach for hypertension. The authors argue for a nuanced understanding of the existing literature, advocating for further research to elucidate the mechanisms by which vitamin D may influence blood pressure. They highlight that while some studies have reported beneficial effects of vitamin D supplementation on blood pressure, others have found no significant impact, indicating a need for more robust clinical trials with larger sample sizes and standardized methodologies.

In evaluating the mechanisms proposed for vitamin D's role in blood pressure regulation, the article discusses potential pathways, including the modulation of renin-angiotensin system activity and the promotion of vascular health. These insights are crucial for understanding how vitamin D may exert its effects on hypertension and for guiding future research directions.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Vitamin D and hypertension" by (Pawlicka et al., 2018) provides a comprehensive examination of the relationship between vitamin D supplementation and blood pressure, particularly focusing on patients with resistant hypertension and older individuals with isolated systolic hypertension. The authors conducted a series of studies that highlighted the ineffectiveness of high-dose oral vitamin D supplementation in reducing blood pressure levels.

One of the key findings is that high-dose vitamin D supplementation, administered at 100,000 IU every three months over a year, did not yield significant improvements in blood pressure or orthostatic hypertension among older patients ((Pawlicka et al., 2018)). This lack of efficacy raises critical questions about the role of vitamin D in hypertension management, particularly in populations considered at risk for cardiovascular complications. The study also noted that ergocalciferol supplementation had no measurable effect on blood pressure, plasma renin activity, or angiotensin II levels, suggesting that vitamin D may not play a direct role in the regulation of these important cardiovascular markers.

Additionally, the research indicated that patients with resistant hypertension had lower vitamin D levels compared to those with white coat hypertension and normotensive individuals ((Pawlicka et al., 2018)). This observation points to a potential correlation between vitamin D deficiency and the severity of hypertension, yet it does not imply causation or therapeutic benefit from supplementation. The authors also reported no significant improvement in endothelial function following vitamin D supplementation, which is a critical factor in vascular health and blood pressure regulation.

The article further critiques the existing literature by highlighting the inconsistency of results from various studies, particularly those conducted on animal models, which may not adequately translate to human physiology ((Pawlicka et al., 2018)). This inconsistency underscores the need for more targeted research to clarify the mechanisms through which vitamin D might

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

influence blood pressure and to determine the appropriate contexts in which supplementation might be beneficial.

The article "Effects of vitamin D supplementation on 25(OH)D concentrations and blood pressure in the elderly: a systematic review and meta-analysis" by (Farapti et al., 2020) presents a comprehensive examination of the relationship between vitamin D supplementation and blood pressure regulation, particularly in elderly populations. The authors systematically reviewed existing literature and conducted a meta-analysis to assess the efficacy of vitamin D in lowering blood pressure among various cohorts, with a particular focus on individuals who are vitamin D deficient.

The findings reported in the article indicate that vitamin D supplementation can have a beneficial effect on lowering blood pressure, especially in patients who are vitamin D deficient and suffer from hypertension. This aligns with previous studies that have suggested a correlation between low vitamin D levels and increased blood pressure, highlighting the importance of maintaining adequate vitamin D status for cardiovascular health ((Farapti et al., 2020)). The authors emphasize that while several meta-analyses support the association between vitamin D deficiency and the risk of developing hypertension, it is crucial to note that correlation does not equate to causation. This distinction is vital as it underscores the need for further research to establish a direct causal relationship.

Moreover, the article points out that significant reductions in blood pressure were primarily observed in hypertensive patients receiving vitamin D supplementation, whereas normotensive individuals did not experience a notable decrease in blood pressure. This observation raises important questions regarding the potential differential effects of vitamin D based on baseline blood pressure status, suggesting that vitamin D may play a more critical role in managing hypertension rather than serving as a universal treatment for all blood pressure conditions ((Farapti et al., 2020)).

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The authors also highlight the necessity for placebo-controlled randomized controlled trials (RCTs) to further elucidate the effects of vitamin D on blood pressure. Such studies would provide more definitive evidence regarding the therapeutic potential of vitamin D supplementation in hypertensive patients and help clarify the mechanisms through which vitamin D influences blood pressure regulation.

The article "Relationship between serum 25-hydroxyvitamin D concentrations and blood pressure among US adults without a previous diagnosis of hypertension: evidence from NHANES 2005–2018" by (Che et al., 2023) presents a comprehensive analysis of the relationship between vitamin D levels and blood pressure among a specific cohort of adults. The study utilizes data from the National Health and Nutrition Examination Survey (NHANES), spanning from 2005 to 2018, to explore the implications of serum 25-hydroxyvitamin D concentrations on blood pressure regulation.

One of the key insights from the article is the demonstrated association between optimal vitamin D status and the attenuation of age-related increases in systolic blood pressure, particularly in white Americans. This finding is crucial as it suggests that maintaining adequate vitamin D levels may play a protective role against hypertension as individuals age. The authors highlight that while serum 25-hydroxyvitamin D levels are significantly correlated with systolic blood pressure, they do not predict future hypertension, indicating a complex relationship that warrants further investigation.

The systematic reviews and meta-analyses referenced within the article underscore the multifaceted nature of vitamin D supplementation and its effects on blood pressure. For instance, the article discusses various studies that have examined the effects of calcium and vitamin D supplementation on blood pressure, revealing a nuanced interaction where these nutrients may jointly influence cardiovascular health. The authors emphasize the importance of considering individual patient data, as responses to vitamin D supplementation can vary based on demographic factors and baseline vitamin D status.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Moreover, the article critically evaluates the role of vitamin D in the renin-angiotensin system, positing that 1,25-dihydroxyvitamin D3 acts as a negative endocrine regulator. This mechanistic insight provides a biological basis for understanding how vitamin D may influence blood pressure regulation and highlights the need for further research to elucidate the underlying pathways..

CONCLUSION

The literature reviewed demonstrates a complex relationship between vitamin D supplementation and blood pressure regulation, with various studies highlighting both potential benefits and inconsistencies in outcomes across different populations and methodologies. The introduction outlines the growing recognition of vitamin D's role beyond bone health, particularly its association with cardiovascular diseases, including hypertension. The prevalence of vitamin D deficiency and its potential implications for chronic diseases underscore the urgency for further research to clarify whether vitamin D deficiency is a causative factor in hypertension or merely a consequence of underlying health issues (Kienreich et al., 2013).

Several articles contribute to the discourse, indicating that while observational studies have established a correlation between vitamin D levels and blood pressure, interventional studies yield mixed results. For instance, high-dose vitamin D supplementation showed no significant effect on blood pressure in resistant hypertension patients (Pawlicka et al., 2018), while systematic reviews indicate potential blood pressure reduction in vitamin D-deficient populations (Farapti et al., 2020). The nuanced relationship is further complicated by the mechanisms proposed for vitamin D's influence on blood pressure, such as its effects on the renin-angiotensin-aldosterone system (RAAS) and endothelial function [ref: 288f4604-dcd6-4d1e-9a52-dea38e60821a; ref: 7b5cd7dd-0262-472f-bf1b-7a0aba5e1b28].

Moreover, the NHANES study data suggest that optimal vitamin D levels may mitigate age-related increases in systolic blood pressure, although vitamin D

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

levels do not predict future hypertension (Che et al., 2023). This highlights the importance of individual patient factors, such as baseline vitamin D status and demographic variables, in understanding the effects of supplementation on blood pressure regulation.

In conclusion, the reviewed literature indicates a need for more rigorous, well-designed randomized controlled trials to establish causality and clarify the mechanisms through which vitamin D may influence blood pressure. While there is evidence suggesting that vitamin D supplementation may lower blood pressure, particularly in deficient populations, inconsistencies in study outcomes necessitate further investigation to develop definitive guidelines for vitamin D supplementation in hypertension management.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Kienreich, K., Tomaschitz, A., Verheyen, N., Pieber, T., Gaksch, M., R. Grübler, M., & Pilz, S., 2013. Vitamin D and Cardiovascular Disease. ncbi.nlm.nih.gov
2. Yun Jeong, H., Mi Park, K., Jung Lee, M., Ho Yang, D., Hoon Kim, S., & Lee, S. Y., 2017. Vitamin D and Hypertension. ncbi.nlm.nih.gov
3. Pawlicka, M., Fałkowska, U., Bałabuszek, K., Radzka, A., Mroczek, A., & Bednarski, J., 2018. Vitamin D and hypertension. [PDF]

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

4. Farapti, F., Fadilla, C., Yogiswara, N., & Adriani, M., 2020. Effects of vitamin D supplementation on 25(OH)D concentrations and blood pressure in the elderly: a systematic review and meta-analysis. ncbi.nlm.nih.gov
5. Che, J., Tong, J., Kuang, X., Zheng, C., Zhou, R., Song, J., Zhan, X., & Liu, Z., 2023. Relationship between serum 25-hydroxyvitamin D concentrations and blood pressure among US adults without a previous diagnosis of hypertension: evidence from NHANES 2005–2018. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved