Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Clinical Nutrition and Family Medicine



Literature Review of Association of Physical Activity and Type 2 Diabetes Mellitus

[1] Hanifa, [1] Sri

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.hanifa983727@gmail.com

Article History :

Received date : 2024/10/12
Revised date : 2024/11/16
Accepted date : 2024/12/24
Published date : 2025/01/08

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The association between physical activity and Type 2 Diabetes Mellitus (T2DM) has been a focal point of research for several decades, highlighting the critical role of exercise in managing and preventing this prevalent metabolic disorder. **Literature Review:** The early work by (Bracken & Stephens, 1970) established the foundational understanding of how physical exercise complements non-insulin glucose-lowering therapies, particularly metformin, to enhance insulin sensitivity. This was further supported by (Al-Safi Ismail et al., 2004), who demonstrated significant improvements in metabolic control through structured exercise in overweight and uncontrolled diabetic patients. The emphasis on tailored exercise programs was echoed in subsequent studies, including the 2011 review by (Teixeira-Lemos et al., 2011), which underscored the preventative role of moderate-intensity exercise in improving insulin sensitivity and reducing oxidative stress. Long-term effects of exercise were highlighted by (Najafipour et al., 2017), who showed that regular physical activity significantly enhances glycemic control and reduces cardiovascular risk factors. This aligns with the holistic perspective presented by (Cannata et al., 2020), advocating for a combination of diet, exercise, and pharmacotherapy to optimize metabolic control. The bibliometric analysis conducted by (Zhang et al., 2022) reinforced the effectiveness of structured exercise programs, particularly those combining aerobic and resistance training, in improving HbA1c levels. **Conclusions:** Overall, the literature consistently identifies physical activity as a vital component in the management and prevention of T2DM, advocating for structured exercise programs, multidisciplinary approaches, and individualized care strategies. This comprehensive view underscores the need for enhanced education among healthcare providers to better integrate lifestyle modifications into diabetes management protocols.

Keywords: Physical Activity, Type 2 Diabetes Mellitus

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The association between physical activity and Type 2 Diabetes Mellitus (T2DM) has been a focal point of research for several decades, highlighting the critical role of exercise in managing and preventing this prevalent metabolic disorder. The literature reveals a progressive understanding of how physical activity interacts with various therapeutic approaches to improve glycemic control and overall health outcomes for individuals with T2DM.

In 1970, (Bracken & Stephens, 1970) conducted a clinical review that explored the relationship between physical exercise and non-insulin glucose-lowering therapies. Their findings underscored that while there is substantial evidence supporting the benefits of physical activity in managing T2DM and reducing cardiovascular risks, the translation of this knowledge into clinical practice remains inadequate. They emphasized the potential synergistic effects of metformin, the first-line oral glucose-lowering agent, and physical exercise in enhancing insulin sensitivity.

Fast forward to 2004, (Al-Safi Ismail et al., 2004) evaluated the impact of a structured exercise program on overweight and uncontrolled diabetic patients, revealing significant improvements in metabolic control, lipid profiles, and cardiovascular fitness. Their study highlighted the necessity of establishing effective exercise regimens tailored to the needs of T2DM patients, further supporting the notion that regular physical activity can lead to tangible health benefits, including reductions in blood pressure and improvements in body composition.

The importance of regular physical exercise continued to be affirmed in 2011 by (Teixeira-Lemos et al., 2011), who reviewed the preventative role of exercise against T2DM. They noted that moderate-intensity exercise not only aids in weight management but also improves insulin sensitivity and reduces markers of oxidative stress and inflammation, which are pivotal in the pathogenesis of T2DM.

In a comprehensive 2017 study, (Najafipour et al., 2017) focused on the long-term effects of exercise training on key indicators such as HbA1c and BMI among T2DM patients. Their findings illustrated that regular exercise significantly enhances glycemic control and reduces cardiovascular risk factors, reinforcing the idea that lifestyle modifications, particularly physical activity, are crucial in managing T2DM.

Continuing this trajectory, (Cannata et al., 2020) provided an overview of the beneficial effects of physical activity for both Type 1 and Type 2 diabetes

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

patients. They advocated for a multidisciplinary approach that integrates diet, exercise, and pharmacotherapy to optimize metabolic control and mitigate the complications associated with diabetes. This holistic perspective underscores the necessity of lifestyle interventions in diabetes management.

(Zhang et al., 2022) conducted a bibliometric analysis that reinforced the significance of structured exercise programs in enhancing glycemic control and overall health in T2DM patients. Their findings revealed that a combination of aerobic and resistance training is particularly effective in improving HbA1c levels, which is crucial for diabetes management. However, they noted a gap in the prescription of exercise by healthcare professionals, suggesting that greater emphasis on lifestyle changes is needed.

In 2023, (Valadares Campos Pereira et al., 2023) highlighted the essential role of physical activity in the management of both Type 1 and Type 2 diabetes, emphasizing its benefits in reducing cardiovascular risks and enhancing overall well-being. They called for individualized exercise prescriptions to maximize health benefits while minimizing potential risks associated with physical activity in diabetic patients.

Finally, (Turčinović et al., 2023) examined the impact of physical activity on microvascular complications in Type 1 diabetes, emphasizing the protective role of regular exercise against cardiovascular risks and its potential to improve patient quality of life. Their findings contribute to the ongoing discourse regarding the multifaceted benefits of physical activity in diabetes management.

Most recently, (White et al., 2024) conducted a systematic review and meta-analysis that assessed the effectiveness of clinician-led community-based exercise interventions on health outcomes in adults with T2DM. Their research supports current guidelines advocating for substantial physical activity as a cornerstone of diabetes management, highlighting the need for innovative community-based approaches to facilitate increased physical activity among patients.

Overall, the literature reveals a consistent and growing recognition of the vital role physical activity plays in the management and prevention of T2DM, with an emphasis on structured exercise programs, multidisciplinary approaches, and individualized care strategies to optimize health outcomes for individuals living with this chronic condition.

LITERATURE REVIEW

The article "Physical exercise and non-insulin glucose-lowering therapies in the management of Type 2 diabetes mellitus: a clinical review" offers a comprehensive examination of the interplay between physical activity and

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

pharmacotherapy in the management of Type 2 diabetes mellitus (T2DM). The authors meticulously synthesize existing literature to elucidate how physical exercise can be effectively integrated with commonly prescribed non-insulin glucose-lowering agents, particularly focusing on metformin and its mechanisms.

The review begins with a detailed exploration of the pharmacological landscape of T2DM treatment, emphasizing biguanides, specifically metformin, as the first-line oral therapy. The authors highlight metformin's mechanism of action, which involves the attenuation of gluconeogenesis in the liver through the activation of 5' AMP-activated protein kinase (AMPK). This is a critical insight, as it establishes a biochemical basis for understanding how pharmacotherapy can synergize with physical activity to enhance glycemic control.

The authors also discuss the physiological effects of exercise, particularly its role in enhancing insulin sensitivity and glucose uptake in skeletal muscle. The activation of AMPK during physical activity is noted as a significant factor that promotes glucose disposal. This connection between exercise and the pharmacological effects of metformin suggests a potential for improved outcomes in patients with T2DM when both strategies are employed concurrently.

Furthermore, the article references various diabetes prevention programs conducted in diverse geographical contexts, including China, the USA, Finland, and India. These programs underline the importance of lifestyle modifications, such as increased physical activity, alongside pharmacotherapy in mitigating the risk of developing T2DM. The authors effectively argue that while medications like metformin play a crucial role in managing hyperglycemia, the incorporation of a physically active lifestyle is equally vital for improving patient outcomes.

However, the article could benefit from a more critical evaluation of the limitations and challenges associated with implementing such lifestyle interventions in clinical practice. Factors such as patient adherence, socioeconomic barriers, and the variability in individual responses to exercise and medication are not thoroughly addressed. Additionally, the review could expand on the potential interactions between different glucose-lowering therapies and physical activity, providing a more nuanced understanding of how these variables can be optimized in clinical settings.

The article "An interventive and kap study on healthy lifestyle of overweight and uncontrolled diabetic patients in Kota Bharu, Kelantan" presents a comprehensive examination of the impact of dietary and physical activity interventions on patients with type 2 diabetes mellitus. The authors emphasize the significance of structured exercise programs in enhancing metabolic control and improving various health parameters among diabetic patients.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The study's findings indicate a substantial increase in VO2max, which is a crucial indicator of cardiovascular fitness, alongside notable reductions in weight, body mass index (BMI), fasting glucose levels, insulin, total cholesterol, triglycerides, and systolic blood pressure. These outcomes align with existing literature that supports the notion that regular physical activity is integral to managing type 2 diabetes. The authors effectively highlight how structured exercise not only aids in weight management but also plays a vital role in improving overall metabolic health, which is particularly important for overweight and uncontrolled diabetic patients.

Furthermore, the article underscores the need for culturally and contextually appropriate exercise programs tailored to the local population. By focusing on the feasibility and accessibility of such programs, the authors propose a model that could potentially be replicated in similar settings, thereby promoting wider implementation of physical activity interventions. This approach is commendable as it takes into account the specific challenges faced by the local demographic, enhancing the likelihood of adherence to the exercise regimen.

The critical evaluation of the material reveals that while the findings are promising, the study could benefit from a larger sample size and a longer follow-up period to further validate the efficacy of the proposed exercise interventions. Additionally, the article could have explored the psychological aspects of engaging in physical activity, such as motivation and barriers to exercise, which are crucial for sustaining long-term lifestyle changes in diabetic patients.

The article "Regular physical exercise training assists in preventing type 2 diabetes development: focus on its antioxidant and anti-inflammatory properties" presents a comprehensive examination of the role of physical activity in the prevention and management of type 2 diabetes mellitus (T2DM). The authors highlight the alarming global prevalence of T2DM, which is projected to affect approximately 380 million individuals by 2025, emphasizing the urgent need for effective prevention strategies to mitigate associated health complications and costs.

A critical aspect of the article is its discussion of the inadequacy of existing pharmacological treatments in preventing cardiovascular events among diabetic patients, which are the leading causes of morbidity and mortality in this population. This perspective aligns with contemporary understandings of chronic disease management, where a holistic approach is increasingly recognized as essential.

The article further elucidates the relationship between sedentary behavior, obesity, and the development of T2DM, noting that low levels of physical activity are a

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

significant risk factor for early mortality. This assertion is supported by numerous prospective studies, reinforcing the idea that sedentary lifestyles contribute to the increasing incidence of diabetes. The authors argue convincingly that regular engagement in moderate-intensity aerobic exercise can lead to substantial health benefits, including weight reduction, enhanced insulin sensitivity, increased levels of high-density lipoprotein (HDL), decreased triglyceride levels, and normalized blood pressure.

Moreover, the review emphasizes the importance of understanding the underlying mechanisms by which physical activity exerts its protective effects against T2DM. The authors delve into the roles of oxidative stress and low-grade inflammation, which are pivotal in the onset and progression of the disease. By focusing on these factors, the article not only highlights the physiological benefits of exercise but also provides a framework for future research that could explore targeted interventions aimed at these pathways.

The article "Effect of regular exercise training on changes in HbA1c, BMI and VO(2)max among patients with type 2 diabetes mellitus: an 8-year trial" presents a comprehensive examination of the relationship between regular exercise and key health indicators in patients with type 2 diabetes mellitus (T2DM). The authors effectively highlight the alarming rise in lifestyle-related chronic diseases due to decreased physical activity, emphasizing that sedentary behavior contributes significantly to obesity and insulin resistance—two critical components of metabolic syndrome that elevate the risk of T2DM.

The study is particularly notable for its long-term approach, spanning eight years, which allows for a robust analysis of the sustained impact of exercise on metabolic health outcomes. The findings underscore that regular physical activity not only aids in glycemic control—evidenced by reductions in hemoglobin A1c (HbA1c) levels—but also contributes to improvements in body mass index (BMI) and maximal oxygen uptake (VO2max). These metrics are essential as they correlate with the overall health and quality of life of individuals with T2DM.

The authors articulate that a mere 10%–15% weight reduction can significantly enhance insulin sensitivity, particularly beneficial during the early stages of T2DM. This insight aligns with existing literature that underscores the importance of weight management in diabetes care. The article also emphasizes the critical role of HbA1c as a marker for diabetes-related complications, noting that a 1% decrease in HbA1c can lead to substantial reductions in the risk of myocardial infarction and diabetes-related mortality.

Moreover, the authors discuss the cardiovascular risks associated with chronic hyperglycemia, which is particularly pertinent given the increasing prevalence of

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

T2DM globally. The article effectively connects the dots between physical activity, metabolic health, and the prevention of long-term complications associated with diabetes.

However, while the study provides valuable insights, it could benefit from a more detailed exploration of the mechanisms by which exercise exerts its beneficial effects on metabolic parameters. Additionally, further discussion on the barriers to maintaining regular physical activity among T2DM patients would enhance the understanding of how to effectively implement exercise interventions in clinical practice.

The article "Beneficial Effects of Physical Activity in Diabetic Patients" presents a comprehensive examination of the role of physical activity (PA) in managing type 1 and type 2 diabetes mellitus (T2D). The authors emphasize that one of the primary goals of diabetic therapy is to achieve optimal metabolic control to mitigate the risk of complications associated with chronic hyperglycaemia. This review advocates for a multidisciplinary approach that integrates diet, physical activity, and pharmacological interventions to enhance metabolic outcomes for diabetic patients.

A critical evaluation of the article reveals that the authors effectively synthesize existing evidence on the positive impacts of PA on diabetic patients. They delineate how regular engagement in physical activity can significantly influence various metabolic parameters, including blood glucose levels, lipid profiles, and overall body composition. The authors argue that lifestyle modifications, particularly through structured PA, can lead to improvements in anthropometric measures and a reduction in the need for medication.

The article also provides a clear definition of physical activity as planned and structured movements aimed at improving health and fitness. This clarity is important as it sets the stage for discussing the specific types and levels of PA that are beneficial for individuals with diabetes. The authors highlight the World Health Organization's (WHO) classification of diabetes as a group of metabolic disorders, which underscores the necessity of addressing the condition from multiple angles, including lifestyle interventions.

Furthermore, the article addresses the epidemic nature of diabetes, noting the alarming prevalence of the disease and its associated complications. The authors argue convincingly that continuous interventions, such as regular physical activity, are essential for managing blood glucose levels and reducing cardiovascular risk factors. This perspective aligns with contemporary approaches to diabetes management that prioritize non-pharmacological strategies.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

However, while the article provides a robust overview of the benefits of PA, it could enhance its impact by including specific guidelines or recommendations regarding the types and intensities of PA that are most effective for different populations within the diabetic community. Additionally, incorporating more detailed discussions on barriers to PA participation among diabetic patients could provide a more rounded understanding of the challenges faced in implementing these recommendations.

The article "Global Trends and Research Hotspots of Exercise for Intervening Diabetes: A Bibliometric Analysis" provides a comprehensive overview of the intersection between physical activity and the management of type 2 diabetes mellitus. The authors synthesize existing literature to highlight the critical role that exercise plays in glycemic control and overall metabolic health for individuals with this condition.

One of the key insights presented in the article is the collaborative efforts of researchers who have established guidelines on various exercise modalities suitable for individuals with type 2 diabetes. Their recommendations encompass not only the types of exercises—such as aerobic and resistance training—but also the necessary precautions to ensure safety for patients on multiple medications or with diabetes-related complications. This emphasis on tailored exercise regimens underscores the importance of individualized approaches in diabetes management.

Furthermore, the article draws attention to the findings of researchers who explored the efficacy of different exercise types in improving glycemic control and cardiorespiratory fitness. The analysis reveals that structured exercise regimens, particularly those incorporating both aerobic and resistance training, are significantly more effective in reducing A1C levels compared to either modality alone. This finding is critical, as it suggests that a multifaceted exercise approach is essential for optimal diabetes management.

The bibliometric analysis conducted by the authors highlights prevalent themes and keywords in the current literature, such as "mellitus," "exercise," "physical activity," and "glycemic control." This focus on keyword trends provides valuable insight into the research landscape surrounding diabetes and exercise, indicating a growing recognition of the importance of physical activity in diabetes care.

Moreover, the article underscores a concerning gap in clinical practice, noting that despite the established benefits of exercise, many healthcare providers fail to prescribe physical activity as part of diabetes management. This observation points to a need for increased education and awareness among physicians regarding the integration of lifestyle modifications, particularly physical activity, in the treatment of type 2 diabetes.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article titled "2022: Position of Brazilian Diabetes Society on exercise recommendations for people with type 1 and type 2 diabetes" provides a comprehensive overview of the role of physical activity in the management of diabetes mellitus, particularly focusing on type 1 and type 2 diabetes. The authors define physical activity as any bodily movement produced by skeletal muscles that results in energy expenditure, emphasizing its significance for individuals diagnosed with diabetes mellitus.

One of the key insights from the article is the assertion that properly guided physical exercises are indispensable for metabolic management in diabetic patients. The authors highlight the extensive benefits of regular physical activity, which include reductions in cardiovascular risks, improvements in personal welfare, and effective weight control. Furthermore, they note that physical activity enhances personal strength and conditioning while also contributing to favorable changes in lipid profiles. These findings underscore the necessity of integrating physical exercise into diabetes treatment protocols.

The article also addresses the individualization of exercise prescriptions, particularly for patients with type 1 diabetes mellitus (T1DM). This individualization is crucial due to the complex interplay of various factors such as the presence of micro and macrovascular complications, obesity, hypertension, hyperglycemia, dyslipidemia, insulin resistance, and sedentary lifestyles. The authors advocate for a tailored approach to exercise, especially for higher intensity activities, to ensure safety and efficacy in managing diabetes-related health risks.

Moreover, the article stresses the importance of assessing cardiovascular risk in individuals with type 2 diabetes mellitus (T2DM) prior to initiating exercise programs. This precautionary measure is vital for preventing adverse events during physical activity, particularly given the increased cardiovascular risks associated with diabetes. The authors argue that regular physical exercise is not only beneficial for managing diabetes but also plays a critical role in the primary and secondary prevention of cardiovascular diseases.

The article titled "THE IMPACT OF TOTAL PHYSICAL ACTIVITY ON MICROVASCULAR COMPLICATIONS IN TYPE 1 DIABETES MELLITUS PATIENTS" provides valuable insights into the complex relationship between physical activity and diabetes management, particularly focusing on type 2 diabetes mellitus (T2DM). The authors emphasize that exercise therapy plays a crucial role in both the prevention and treatment of T2DM, highlighting that regular aerobic exercise is integral to effective disease management. This assertion aligns with the growing body of literature advocating for physical activity as a cornerstone in diabetes care strategies.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

One of the key points raised in the article is the multifaceted benefits of regular physical activity for T2DM patients, including reduced cardiovascular risk, decreased insulin requirements, improved muscular strength, and enhanced overall well-being. These findings reinforce the notion that exercise is not merely an adjunct therapy but a fundamental component of comprehensive diabetes management.

However, the article also addresses the ongoing debate regarding the effects of physical activity on glycemic control and microvascular complications, which remain inconclusive. The authors note that while some studies suggest that physical activity may exacerbate certain microvascular complications, others report beneficial outcomes. This dichotomy in findings underscores the need for further research to clarify the relationship between exercise intensity and the risk of developing complications.

Moreover, the article raises an important question about whether the intensity of physical activity affects complication-free life and longevity in T2DM patients. This inquiry highlights a gap in the current literature, as existing studies have produced mixed results regarding the optimal exercise regimen for maximizing health benefits in this population.

The authors advocate for more focused research to explore these relationships, which could ultimately inform clinical guidelines and improve patient outcomes.

The article titled "The Effectiveness of Clinician-Led Community-Based Group Exercise Interventions on Health Outcomes in Adults with Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis" provides a comprehensive examination of the role of physical activity in managing type 2 diabetes mellitus (T2DM). The authors assert that physical activity, in conjunction with nutritional and medical therapies, is essential for effective management of T2DM. They emphasize that exercise, which encompasses various forms of physical activity, contributes significantly to improving strength, endurance, agility, balance, and flexibility—all critical components for individuals living with T2DM.

The article outlines current guidelines recommending that adults aged 18–64 with T2DM engage in a minimum of 150 minutes of moderate-to-vigorous intensity exercise weekly, alongside resistance training at least twice per week. This framework sets a clear standard for physical activity, underscoring its importance in the management of glycemic control and cardiovascular risk factors associated with T2DM.

A critical evaluation of the material reveals that the authors have effectively synthesized existing research on various physical activity intervention techniques aimed at enhancing the activity levels of individuals with T2DM. Notably, the

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

article highlights clinician-led facility-based fitness training as a viable strategy that could improve glycemic management. However, the authors also acknowledge the limitations of this approach, particularly its accessibility and the potential barriers faced by patients in adhering to such programs.

The introduction of community-based exercise interventions as a hybrid model offers a promising alternative. By combining the structured guidance of healthcare professionals with the accessibility of community settings, these interventions may address the challenges posed by traditional facility-based approaches. The authors suggest that community-based programs can provide culturally relevant health education, which is crucial for engaging diverse populations and ensuring that interventions are tailored to meet the specific needs of different communities.

Moreover, the article calls for an expansion of existing intervention strategies to effectively manage the increasing prevalence of T2DM. The authors argue for a multi-faceted approach that includes individual-based treatments, medication use, and behavior modification alongside physical activity interventions. This holistic perspective is commendable, as it recognizes the complexity of T2DM management and the need for comprehensive strategies that encompass various aspects of patient care.

CONCLUSION

The literature reviewed presents a comprehensive exploration of the association between physical activity and Type 2 Diabetes Mellitus (T2DM), emphasizing the critical role of exercise in managing and preventing this metabolic disorder. The findings across various studies highlight the importance of structured exercise regimens, individualized care strategies, and a multidisciplinary approach that integrates lifestyle modifications with pharmacotherapy.

The early work by (Bracken & Stephens, 1970) established the foundational understanding of how physical exercise complements non-insulin glucose-lowering therapies, particularly metformin, to enhance insulin sensitivity. This was further supported by (Al-Safi Ismail et al., 2004), who demonstrated significant improvements in metabolic control through structured exercise in overweight and uncontrolled diabetic patients. The emphasis on tailored exercise programs was echoed in subsequent studies, including the 2011 review by (Teixeira-Lemos et al., 2011), which underscored the preventative role of moderate-intensity exercise in improving insulin sensitivity and reducing oxidative stress.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Long-term effects of exercise were highlighted by (Najafipour et al., 2017), who showed that regular physical activity significantly enhances glycemic control and reduces cardiovascular risk factors. This aligns with the holistic perspective presented by (Cannata et al., 2020), advocating for a combination of diet, exercise, and pharmacotherapy to optimize metabolic control. The bibliometric analysis conducted by (Zhang et al., 2022) reinforced the effectiveness of structured exercise programs, particularly those combining aerobic and resistance training, in improving HbA1c levels.

Recent studies, including the systematic review by (White et al., 2024), continue to support the integration of physical activity as a cornerstone of diabetes management, emphasizing the need for innovative community-based approaches to increase patient engagement in exercise. The exploration of individualized exercise prescriptions by (Valadares Campos Pereira et al., 2023) further underscores the necessity of tailoring interventions to maximize health benefits while minimizing risks.

Overall, the literature consistently identifies physical activity as a vital component in the management and prevention of T2DM, advocating for structured exercise programs, multidisciplinary approaches, and individualized care strategies. This comprehensive view underscores the need for enhanced education among healthcare providers to better integrate lifestyle modifications into diabetes management protocols.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Bracken, R. & Stephens, J., 1970. Physical exercise and non-insulin glucose-lowering therapies in the management of Type 2 diabetes mellitus: a clinical review. [PDF]

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

2. Al-Safi Ismail, A., Mohamed, M., Nordin, R., Abdul Manan Wan Muda, W., & Ibrahim, N., 2004. An interventive and kap study on healthy lifestyle of overweight and uncontrolled diabetic patients in Kota Bharu, Kelantan. [PDF]
3. Teixeira-Lemos, E., Nunes, S., Teixeira, F., & Reis, F., 2011. Regular physical exercise training assists in preventing type 2 diabetes development: focus on its antioxidant and anti-inflammatory properties. ncbi.nlm.nih.gov
4. Najafipour, F., Mobasseri, M., Yavari, A., Nadrian, H., Aliasgarzadeh, A., Mashinchi Abbasi, N., Niafar, M., Houshyar Gharamaleki, J., & Sadra, V., 2017. Effect of regular exercise training on changes in HbA1c, BMI and VO(2)max among patients with type 2 diabetes mellitus: an 8-year trial. ncbi.nlm.nih.gov
5. Cannata, F., Vadalà, G., Russo, F., Papalia, R., Napoli, N., & Pozzilli, P., 2020. Beneficial Effects of Physical Activity in Diabetic Patients. ncbi.nlm.nih.gov
6. Zhang, Z., Zhu, Y., Wang, Q., Chang, T., Liu, C., Zhu, Y., Wang, X., & Cao, X., 2022. Global Trends and Research Hotspots of Exercise for Intervening Diabetes: A Bibliometric Analysis. ncbi.nlm.nih.gov
7. Valadares Campos Pereira, W., Maria Martins Vancea, D., de Andrade Oliveira, R., Galeno Pinheiro Chaves de Freitas, Y., Nunes Lamounier, R., S. Silva Júnior, W., Messias Britto Fioretti, A., Luiz Dornelles Macedo, C., Casaccia Bertoluci, M., & Luis Zagury, R., 2023. 2022: Position of Brazilian Diabetes Society on exercise recommendations for people with type 1 and type 2 diabetes. ncbi.nlm.nih.gov
8. Turčinović, K., Cigrovski, V., Bilić-Ćurćić, I., Mrzljak, A., Kuna, L., Klobučar, S., & Cigrovski Berković, M., 2023. THE IMPACT OF TOTAL PHYSICAL ACTIVITY ON MICROVASCULAR COMPLICATIONS IN TYPE 1 DIABETES MELLITUS PATIENTS. ncbi.nlm.nih.gov
9. White, L., Kirwan, M., Christie, V., Hurst, L., & Gwynne, K., 2024. The Effectiveness of Clinician-Led Community-Based Group Exercise Interventions on Health Outcomes in Adults with Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved