Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Clinical Nutrition and Family Medicine



Management of Malnutrition

[1] Wahyu, [1] Dedi

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.wahyu909838@gmail.com

Article History :

Received date : 2024/10/19
Revised date : 2024/11/10
Accepted date : 2024/12/24
Published date : 2025/01/08

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The management of malnutrition is a critical concern within healthcare systems, particularly in hospital and community settings, where its prevalence can significantly impact patient outcomes and healthcare costs. **Literature Review:** A recurring theme across the reviewed articles is the urgent need for improved screening and management strategies. For instance, (A. Barker et al., 2011) advocate for mandatory nutrition risk screening upon hospital admission to facilitate timely dietary referrals, while (Kabashneh et al., 2020) emphasize the importance of integrating routine nutrition assessments in clinical practice. Despite the recognition of these needs, the implementation of standardized protocols remains inconsistent, resulting in underdiagnosis and inadequate management of malnutrition. The literature also underscores the multifactorial nature of malnutrition, which is influenced by various social, clinical, and environmental factors. (Rémond et al., 2015) highlight that dietary interventions must account for the unique needs of the elderly, including physiological changes that affect nutrient absorption and utilization. Furthermore, (Brunner et al., 2021) discuss the role of external factors, such as meal quality and care environments, in exacerbating nutritional deficiencies among older adults. In community settings, (S. Harris et al., 2019) point out the barriers to effective malnutrition management, including a lack of prioritization of nutritional assessments in primary care. Their findings suggest that healthcare professionals often overlook nutritional issues in favor of more immediate health concerns, indicating a need for enhanced training and awareness among practitioners (Lindner-Rabl et al., 2022). The synthesis of qualitative and quantitative data in their study reveals that systematic screening could significantly improve the identification of at-risk individuals, thereby enabling timely interventions. **Conclusions:** In conclusion, the reviewed literature collectively underscores the need for a nuanced approach to iron supplementation in breastfed infants. While iron is essential for preventing IDA and promoting health, the timing, dosage, and broader context of infant health must be carefully considered to optimize developmental outcomes and avoid potential adverse effects. Further research is essential to clarify the relationships between iron supplementation, growth, neurodevelopment, and overall health in this vulnerable population.

Keywords: Management, Malnutrition

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The management of malnutrition, particularly in hospital and community settings, has garnered significant attention in recent years due to its profound implications for patient health outcomes and healthcare costs. The literature reveals a consistent theme: malnutrition is a prevalent and serious condition that adversely affects various patient demographics, especially the elderly. (A. Barker et al., 2011) highlight the alarming prevalence of malnutrition in acute hospital settings, noting that rates can reach approximately 40%. They emphasize the dire consequences of malnutrition, including impaired immune function, prolonged hospital stays, and increased mortality. The authors advocate for mandatory nutrition risk screening upon hospital admission to ensure timely dietetic referrals and mitigate the risks associated with undiagnosed malnutrition.

Building on this foundation, (Rémond et al., 2015) explore the multifaceted nature of malnutrition in the elderly, linking it to increased healthcare costs and adverse clinical outcomes. Their findings indicate that malnourished patients incur significantly higher daily expenses and experience longer hospitalizations. They propose dietary interventions as essential strategies to combat malnutrition, underscoring the importance of addressing both social and clinical factors that contribute to poor nutritional status in this vulnerable population.

In a similar vein, (Gen-jida & Takeda, 2018) discuss the critical role of nutrition support in hospitalized patients, revealing that malnutrition is present in 40% to 50% of cases. Their review establishes a direct correlation between malnutrition and increased morbidity and mortality, advocating for routine nutrition screening as a pivotal component of patient care. They argue that despite the recognized risks, nutrition assessments remain inconsistently implemented, highlighting a gap in current medical practices.

(Volkert et al., 2019) further elaborate on the complex etiology of malnutrition in older adults, detailing how external factors, such as meal quality and care environments, can exacerbate dietary deficiencies. They emphasize the severe

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

consequences of malnutrition on recovery and overall health, including increased risks of falls and fractures, thereby reinforcing the need for comprehensive management strategies.

(Kabashneh et al., 2020) contribute to the discourse by providing a comprehensive guide on identifying and managing malnutrition in hospitalized patients. They emphasize the need for increased awareness among healthcare providers regarding the prevalence and consequences of malnutrition, advocating for standardized diagnostic criteria and management protocols to improve patient outcomes.

(Brunner et al., 2021) conduct an umbrella review focused on older individuals in hospitals and long-term care settings, reiterating the global concern surrounding malnutrition. They point out the significant gaps in practice related to nutritional risk screening and the complexity of the nutrition process within healthcare institutions, suggesting that a lack of prioritization of nutritional care contributes to the persistence of malnutrition.

Finally, (Lindner-Rabl et al., 2022) present a scoping review that examines current practices in primary healthcare for managing malnutrition in older adults. They highlight the variability in awareness and approaches across different countries, advocating for dietician-led interventions and the use of standardized screening tools to improve nutritional care. Their findings suggest that personalized nutritional strategies are crucial for effectively addressing malnutrition in diverse patient populations.

Overall, the literature underscores the multifaceted nature of malnutrition management, revealing significant challenges and opportunities for improving care across various healthcare settings. The consistent call for enhanced screening, tailored interventions, and increased awareness among healthcare professionals reflects a growing recognition of malnutrition as a critical public health issue that warrants urgent attention.

LITERATURE REVIEW

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Hospital malnutrition: prevalence, identification and impact on patients and the healthcare system" by Barker, Gout, and Crowe (A. Barker et al., 2011) presents a comprehensive examination of malnutrition within the acute hospital setting, highlighting its prevalence, consequences, and the need for improved identification and management strategies. The authors assert that approximately 40% of hospitalized patients are malnourished, a statistic that underscores the critical nature of this issue in healthcare environments.

One of the key insights from the article is the multifaceted impact of malnutrition on patient outcomes. The authors detail how malnutrition can lead to a range of adverse effects, including a depressed immune response, impaired wound healing, muscle wasting, extended hospital stays, increased healthcare costs, and heightened mortality rates. This connection between malnutrition and negative health outcomes emphasizes the necessity for healthcare systems to prioritize nutritional care as part of comprehensive patient management.

Despite the evident risks associated with malnutrition, the article reveals that referral rates for dietetic assessment and treatment of malnourished patients are alarmingly low. This gap in care is particularly concerning given that nutrition risk screening, when implemented using validated tools, can effectively identify at-risk patients and facilitate timely dietary interventions. The authors argue for the implementation of mandatory nutrition screening upon hospital admission, as current practices in Australia do not require such assessments. This lack of mandatory screening contributes to the underreporting and inadequate documentation of malnutrition, ultimately exacerbating the risks of adverse patient outcomes.

Moreover, Barker et al. (A. Barker et al., 2011) discuss the financial implications of malnutrition, noting that unidentified cases can lead to lost reimbursements for hospitals. This point highlights the dual burden of malnutrition on both patient health and healthcare systems, suggesting that addressing this issue could lead to improved clinical and financial outcomes.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Understanding the gastrointestinal tract of the elderly to develop dietary solutions that prevent malnutrition" by Rémond et al. (Rémond et al., 2015) provides a comprehensive examination of the multifaceted issue of malnutrition in the elderly, highlighting its significant impact on health outcomes and healthcare costs. The authors emphasize that malnutrition is not merely a nutritional deficiency but an independent risk factor that exacerbates various hospital-related parameters, including length of stay, complications, mortality, and overall healthcare expenditures.

One of the critical insights presented in the article is the stark contrast in daily medical expenses between malnourished and well-nourished patients, with the former incurring a mean daily cost of US$ 228 compared to US$ 138 for their well-nourished counterparts. This disparity underscores the economic burden that malnutrition places on healthcare systems, particularly in the context of an aging population that is increasingly susceptible to nutritional deficiencies. The authors reference a study from Ben-Gurion University, which reveals that elderly patients at risk of malnutrition experience a doubling of hospitalization days compared to those with adequate nutritional status. This finding is particularly alarming, as it highlights the urgent need for effective interventions to address malnutrition in this demographic.

The article outlines several strategies to treat or prevent malnutrition, which include social interventions, nutritional supplements, improvements in care settings, and clinical interventions. However, the authors place particular emphasis on dietary interventions as the cornerstone of any effective strategy. They argue that ensuring an adequate food supply tailored to the unique needs of the elderly is essential in mitigating the adverse effects of malnutrition. Furthermore, the authors elucidate the physiological consequences of malnutrition, detailing its detrimental effects on organ function, immunity, muscle strength, and mental health, including its association with conditions such as sarcopenia and dementia.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The authors effectively convey the message that poor nutritional status is a major negative prognostic indicator in the elderly, linking it to increased mortality rates, longer hospital stays, and a cascade of other health-related issues that ultimately lead to diminished quality of life. This critical evaluation of the material highlights the urgent need for healthcare providers and policymakers to prioritize nutritional health in the elderly population, advocating for a multifaceted approach that encompasses dietary, social, and clinical strategies.

The article titled "The role of nutrition support" by Gen-jida and Takeda (Gen-jida & Takeda, 2018) provides a comprehensive examination of the prevalence and implications of malnutrition within hospital settings, highlighting its significant impact on patient outcomes and healthcare costs. The authors present compelling data indicating that a substantial portion of hospitalized patients—between 40% to 50%—are at risk for malnutrition, with severe malnutrition affecting up to 12% of this population. This statistic underscores the critical need for effective management strategies in addressing malnutrition among patients.

Central to the article's argument is the assertion that malnutrition not only exacerbates existing health conditions but also contributes to increased morbidity and mortality rates. The authors cite research that establishes a direct cause-effect relationship between malnutrition and adverse health outcomes, particularly highlighting its detrimental effects on cellular immunity. This is particularly concerning for vulnerable populations such as infants, children, and the elderly, who are disproportionately affected by the consequences of malnutrition. The authors effectively articulate that profound malnutrition can lead to growth retardation and a heightened susceptibility to infections, thus emphasizing the urgency of addressing this issue within healthcare systems.

The review further advocates for the implementation of routine nutrition screening and follow-up assessments as integral components of patient care. The authors argue that despite the clear evidence supporting the need for nutritional interventions, these practices are not consistently integrated into standard medical protocols. This gap in care presents a significant opportunity for improvement in

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

clinical practices. By providing scientific and economic justifications for routine nutrition screening, Gen-jida and Takeda encourage healthcare professionals to prioritize nutritional support as a means to enhance patient health outcomes and reduce healthcare costs associated with malnutrition-related complications.

The article titled "Management of Malnutrition in Older Patients—Current Approaches, Evidence and Open Questions" by Volkert et al. (Volkert et al., 2019) provides a comprehensive overview of the multifaceted nature of malnutrition in older adults, emphasizing the significant impact of external factors such as meal quality and care standards on dietary intake. The authors argue that malnutrition is not only prevalent but also a critical geriatric syndrome that exacerbates morbidity and mortality rates among older populations.

One of the key insights presented in the article is the complex interplay of factors contributing to malnutrition, which includes inadequate protein intake leading to loss of body protein. This loss is particularly concerning as it compromises immune function and contributes to muscle mass decline, which is essential for maintaining physical health and independence in older adults. The authors effectively highlight that decreased muscle strength and mass can lead to a cascade of negative outcomes, including impaired physical status, increased risk of falls, fractures, and a diminished quality of life.

The article further discusses the implications of malnutrition on healthcare systems, noting that malnourished patients experience longer hospital stays, higher rates of complications, and increased health resource utilization. This aspect underscores the economic burden that malnutrition imposes on healthcare systems, suggesting that addressing malnutrition could lead to improved patient outcomes and reduced healthcare costs.

Volkert et al. (Volkert et al., 2019) also call attention to the need for a multi-disciplinary approach to the management of malnutrition in older patients. They advocate for improved nutritional screening and assessment practices in hospitals and care facilities, as well as the implementation of tailored nutritional

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

interventions to enhance dietary intake. The authors suggest that addressing malnutrition requires a collaborative effort among healthcare professionals, including dietitians, nurses, and physicians, to create a supportive environment that fosters optimal nutritional care.

Despite the thorough examination of the current state of malnutrition management, the article identifies several open questions that warrant further research. For instance, the authors note the need for more studies to evaluate the effectiveness of various interventions and to explore how different care settings can better address the nutritional needs of older adults. This acknowledgment of gaps in the literature indicates a forward-thinking approach, encouraging continued investigation into the complexities of malnutrition.

The article titled "Barriers and facilitators to screening and treating malnutrition in older adults living in the community: a mixed-methods synthesis" by Harris et al. (S. Harris et al., 2019) provides an in-depth analysis of the multifaceted issue of malnutrition among older adults, particularly those residing in community settings. The authors highlight the significant health risks associated with malnutrition, including impaired wound healing, reduced muscle strength, and weakened immune responses, all of which can exacerbate the vulnerability of this demographic, especially given the increasing prevalence of long-term health conditions.

One of the key insights presented in the article is the distinction between malnutrition and undernutrition, both of which are often conflated. The authors clarify that malnutrition can arise from inadequate nutrient intake or from the body's inability to utilize nutrients effectively. This nuanced understanding is critical for healthcare providers as it underscores the complexity of diagnosing and treating malnutrition, particularly in older adults who may present with multiple health issues.

The article provides compelling statistics, noting that over 3 million individuals in the UK are estimated to be malnourished, with the financial burden on health and

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

social care systems reaching £20 billion in 2015. This economic perspective emphasizes the urgent need for effective screening and treatment strategies. The authors assert that systematic screening using validated tools can significantly enhance the identification of at-risk individuals, thereby facilitating timely intervention.

However, the article also critically evaluates the current state of malnutrition management in primary care settings. Despite the recommendations for malnutrition-screen-and-treat policies in hospitals, the authors point out the lack of evidence supporting their implementation in community care. This gap highlights a crucial barrier to effective management, as healthcare professionals often prioritize other health issues over nutritional assessments in older patients. The authors call attention to the need for appropriate training for healthcare providers in screening practices, as existing clinical guidelines do not adequately address the specifics of training or the enactment of screening procedures.

Furthermore, the synthesis of qualitative and quantitative findings from various studies offers valuable insights into the facilitators and barriers to implementing malnutrition interventions. The authors advocate for further research to explore these barriers in greater depth, particularly in the context of community-dwelling older adults. Their recommendations for designing interventions and enhancing nutrition education for healthcare professionals are particularly noteworthy, as they aim to bridge the existing gap in knowledge and practice.

The article titled "A Complete Guide to Identify and Manage Malnutrition in Hospitalized Patients" by Kabashneh et al. (Kabashneh et al., 2020) provides a comprehensive overview of malnutrition within the hospital setting, emphasizing its prevalence and the critical need for effective management strategies. The authors highlight that malnutrition affects approximately half of patients admitted to acute care hospitals, leading to a range of adverse health outcomes, including increased susceptibility to infections, immunodeficiency, and the development of pressure ulcers.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

A significant contribution of this article is its focus on the underrecognition of malnutrition in clinical practice. Despite the high incidence of malnutrition among hospitalized patients, the authors note that many healthcare systems do not mandate nutritional assessments during patient admission and throughout their hospital stay. This oversight can lead to inadequate diagnosis and management, ultimately compromising patient outcomes. The article effectively argues for the integration of routine nutritional screening as a standard practice in hospitals, which is crucial for improving the recognition and treatment of malnutrition.

The authors present various screening methodologies that healthcare providers can utilize to assess malnutrition. They discuss tools such as the Malnutrition Universal Screening Tool (MUST) and the Subjective Global Assessment (SGA), which can facilitate the early identification of at-risk patients. By providing a guide on these screening techniques, the article serves as a valuable resource for healthcare practitioners aiming to enhance their approach to nutritional assessment.

Furthermore, the review delves into the consequences of malnutrition, elucidating its direct correlation with increased morbidity and mortality rates in hospitalized patients. The authors effectively use evidence from current literature to support their claims, reinforcing the urgency of addressing malnutrition as a significant medical issue. They also emphasize the role of nutritional supplements in the management of malnourished patients, outlining the benefits of early intervention and tailored nutritional plans.

The article titled "Interventions to optimise nutrition in older people in hospitals and long-term care: Umbrella review" by Brunner et al. (Brunner et al., 2021) provides a comprehensive overview of the pressing issue of malnutrition among older adults, particularly within hospital and long-term care settings. The authors underscore the significant prevalence of malnutrition, citing rates that can reach as high as 50% in hospitalized elderly populations, which is a critical concern given the associated risks such as increased complications, care dependency, and mortality.

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

A key insight from the article is the multifaceted nature of malnutrition risk factors in older adults. The authors identify various contributors, including impaired cognitive function, physiological changes such as muscle mass loss, diminished taste, and oral health issues. Additionally, they highlight the role of multimorbidity and polypharmacy in exacerbating nutritional challenges, which is particularly relevant in geriatric care. This comprehensive identification of risk factors is crucial for developing targeted interventions that address the specific needs of this vulnerable population.

The article also discusses the systemic gaps in practice related to nutritional risk screening. Brunner et al. (Brunner et al., 2021) emphasize that despite the existence of numerous systematic reviews detailing nutrition-improving interventions, the recognition and treatment of malnutrition risk often remain inadequate. This discrepancy can be attributed to the complexity of nutritional care processes, which involve multiple healthcare professionals, and a general lack of awareness regarding the significance of nutrition in patient outcomes. The authors argue that these barriers contribute to the underprioritization of nutritional support during food intake, which is essential for older adults.

Furthermore, the review aims to fill the identified gap by aggregating evidence-based interventions that have proven effective in mitigating malnutrition risks. This is a commendable approach, as it not only consolidates existing knowledge but also serves as a practical guide for healthcare providers aiming to implement effective nutritional strategies. However, the article could benefit from a more detailed discussion on the specific interventions identified and their comparative effectiveness, as this would provide clearer guidance for practitioners.

The article "Clinical Interventions to Improve Nutritional Care in Older Adults and Patients in Primary Healthcare – A Scoping Review of Current Practices of Health Care Practitioners" by Lindner-Rabl et al. (Lindner-Rabl et al., 2022) provides a comprehensive overview of the current state of nutritional interventions aimed at addressing malnutrition among older adults within primary healthcare settings. The authors systematically review existing practices and

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

highlight the significant challenges faced by healthcare practitioners in effectively managing malnutrition.

One of the core insights from the article is the divergence in evidence regarding the efficacy of various interventions designed to mitigate the risk of malnutrition. This inconsistency underscores the complexity of nutritional care, particularly in community settings where older adults are prevalent. The authors argue that addressing malnutrition is not merely a matter of improving individual health outcomes; it also has broader implications for the healthcare system by reducing costs and resource utilization (Lindner-Rabl et al., 2022). This perspective aligns with the growing recognition of the need for preventive strategies in healthcare, especially as the population ages and the prevalence of malnutrition rises.

A critical aspect of the article is its emphasis on the role of dieticians in the management of malnutrition. The authors advocate for a more prominent and proactive involvement of dieticians in primary care, suggesting that dietician-led interventions have shown promising results in improving nutritional outcomes. This recommendation is particularly relevant as it highlights the necessity for specialized knowledge in devising effective nutritional strategies tailored to individual patient needs. The article posits that personalized care approaches are essential for addressing the diverse nutritional requirements and preferences of older adults, thereby enhancing the effectiveness of interventions (Lindner-Rabl et al., 2022).

Furthermore, the review emphasizes the importance of accurately identifying patients at risk of malnutrition. Effective nutritional care hinges on the ability to recognize malnourished individuals early, which necessitates training and awareness among healthcare practitioners. The authors suggest that improving identification processes can lead to timely interventions, ultimately contributing to better health outcomes and improved quality of life for older adults.

CONCLUSION

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The management of malnutrition is a critical concern within healthcare systems, particularly in hospital and community settings, where its prevalence can significantly impact patient outcomes and healthcare costs. The literature consistently highlights the alarming rates of malnutrition, especially among elderly populations, with estimates suggesting that 40% to 50% of hospitalized patients are affected (A. Barker et al., 2011). This condition not only leads to adverse health outcomes, such as increased morbidity and mortality, but also imposes a substantial financial burden on healthcare systems [ref: 9dd55f0f-e76b-489f-991f-a71762ff47ed; ref: 610c1047-aec2-446a-8a06-8eb8643c1400].

A recurring theme across the reviewed articles is the urgent need for improved screening and management strategies. For instance, (A. Barker et al., 2011) advocate for mandatory nutrition risk screening upon hospital admission to facilitate timely dietary referrals, while (Kabashneh et al., 2020) emphasize the importance of integrating routine nutrition assessments in clinical practice. Despite the recognition of these needs, the implementation of standardized protocols remains inconsistent, resulting in underdiagnosis and inadequate management of malnutrition [ref: 5d5c9085-68d6-4539-aa37-12c6a1bdd1e1; ref: 35c9518a-8e0a-4b63-be12-6eb7dd8bdfa6].

The literature also underscores the multifactorial nature of malnutrition, which is influenced by various social, clinical, and environmental factors. (Rémond et al., 2015) highlight that dietary interventions must account for the unique needs of the elderly, including physiological changes that affect nutrient absorption and utilization. Furthermore, (Brunner et al., 2021) discuss the role of external factors, such as meal quality and care environments, in exacerbating nutritional deficiencies among older adults.

In community settings, (S. Harris et al., 2019) point out the barriers to effective malnutrition management, including a lack of prioritization of nutritional assessments in primary care. Their findings suggest that healthcare professionals often overlook nutritional issues in favor of more immediate health concerns, indicating a need for enhanced training and awareness among practitioners

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

(Lindner-Rabl et al., 2022). The synthesis of qualitative and quantitative data in their study reveals that systematic screening could significantly improve the identification of at-risk individuals, thereby enabling timely interventions.

In conclusion, the literature collectively emphasizes that addressing malnutrition requires a multifaceted approach, encompassing improved screening practices, tailored dietary interventions, and increased awareness among healthcare professionals. The evidence suggests that prioritizing nutritional care could lead to better health outcomes and reduced healthcare costs, particularly for vulnerable populations such as the elderly. As malnutrition remains a critical public health issue, ongoing research and policy efforts are essential to bridge the existing gaps in management strategies and enhance the quality of care provided to affected individuals.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. A. Barker, L., S. Gout, B., & C. Crowe, T., 2011. Hospital malnutrition : prevalence, identification and impact on patients and the healthcare system. [PDF]
2. Rémond, D., R. Shahar, D., Gille, D., Pinto, P., Kachal, J., Peyron, M. A., Nunes Dos Santos, C., Walther, B., Bordoni, A., Dupont, D., Tomás-Cobos, L., & Vergères, G., 2015. Understanding the gastrointestinal tract of the elderly to develop dietary solutions that prevent malnutrition. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

3. Gen-jida, K. & Takeda, E., 2018. The role of nutrition support. [PDF]
4. Volkert, D., Marie Beck, A., Cederholm, T., Cereda, E., Cruz-Jentoft, A., Goisser, S., de Groot, L., Großhauser, F., Kiesswetter, E., Norman, K., Pourhassan, M., Reinders, I., C. Roberts, H., Rolland, Y., M. Schneider, S., C. Sieber, C., Thiem, U., Visser, M., A.H. Wijnhoven, H., & Wirth, R., 2019. Management of Malnutrition in Older Patients—Current Approaches, Evidence and Open Questions. ncbi.nlm.nih.gov
5. Kabashneh, S., Alkassis, S., Shanah, L., & Ali, H., 2020. A Complete Guide to Identify and Manage Malnutrition in Hospitalized Patients. ncbi.nlm.nih.gov
6. Brunner, S., Mayer, H., Qin, H., Breidert, M., Dietrich, M., & Müller Staub, M., 2021. Interventions to optimise nutrition in older people in hospitals and long-term care: Umbrella review. ncbi.nlm.nih.gov
7. Lindner-Rabl, S., Wagner, V., Matijevic, A., Herzog, C., Lampl, C., Traub, J., & Roller-Wirnsberger, R., 2022. Clinical Interventions to Improve Nutritional Care in Older Adults and Patients in Primary Healthcare – A Scoping Review of Current Practices of Health Care Practitioners. ncbi.nlm.nih.gov
8. S. Harris, P., Payne, L., Morrison, L., M. Green, S., Ghio, D., Hallett, C., L. Parsons, E., Aveyard, P., C. Roberts, H., Sutcliffe, M., Robinson, S., Slodkowska-Barabasz, J., S. Little, P., A. Stroud, M., & Yardley, L., 2019. Barriers and facilitators to screening and treating malnutrition in older adults living in the community: a mixed-methods synthesis. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Clinical Nutrition and Family Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved