Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Clinical Pathology and Microbiology



Serologic Screening for Genital Herpes

[1] Maria, [1] Yohanes

[1] Faculty of Medicine, Maranatha Christian University, Bandung City, West Java, Indonesia

Correspondence : dokter.maria938483@gmail.com

Article History :

Received date : 2024/07/15
Revised date : 2024/08/26
Accepted date : 2024/09/08
Published date : 2025/01/12

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature on serologic screening for genital herpes has seen significant advancements, emphasizing the importance of comprehensive methodologies, epidemiological insights, psychosocial impacts, and economic considerations. **Literature Review:** (Miles, 2017) provided a qualitative perspective on the psychological and social ramifications of HSV-2 diagnoses, particularly regarding safe sex practices and disclosure dynamics in intimate relationships. This study, complemented by (Ann Hermes, 2017), which focused on couples with discordant HSV statuses, highlighted the relational complexities and emotional challenges faced by individuals living with HSV. These insights stress the importance of considering the psychosocial dimensions of HSV management in clinical practice. The systematic reviews by (Harfouche et al., 2019) and (Chaabane et al., 2019) further elucidated HSV epidemiology across different regions, advocating for the use of standardized serological assays to monitor trends in HSV-1 and HSV-2 infections. (Sukik et al., 2019) extended this analysis to Latin America and the Caribbean, emphasizing the need for region-specific data to inform effective healthcare interventions. **Conclusion:** In conclusion, the body of literature on serologic screening for genital herpes presents a multifaceted understanding of the complexities surrounding HSV infections. The integration of diagnostic methodologies, epidemiological data, psychosocial insights, and economic evaluations indicates a pressing need for comprehensive public health strategies that address the various dimensions of genital herpes. Such an approach is crucial for improving patient care and ultimately reducing the burden of HSV infections globally.

Keyword: Serologic Screening, Genital Herpes

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The literature on serologic screening for genital herpes has evolved significantly over the years, highlighting various aspects of epidemiology, diagnosis, and the social implications of herpes simplex virus (HSV) infections. The foundational work by Aryee et al. (AN Aryee et al., 2005) established essential methodologies for detecting HSV in cervicovaginal secretions through real-time PCR and highlighted the importance of comprehensive serological testing. Their study underscored the necessity of using multiple diagnostic tools, such as ELISA and rapid plasma reagin tests, to accurately identify co-infections and the prevalence of HSV among patients.

Building on this foundation, Nyiro et al. (U. Nyiro et al., 2011) explored the seroprevalence and transmission dynamics of HSV-2 among semi-urban women in Kenya. Their findings revealed critical risk factors associated with undiagnosed infections and emphasized the epidemiological links between HSV-2 and HIV, thereby illustrating the broader public health implications of genital herpes screening. This study contributed to understanding how geographic and demographic variations influence HSV prevalence, necessitating tailored screening approaches.

In 2017, Miles conducted a qualitative investigation into the disclosure patterns of individuals diagnosed with HSV-2, probing the psychological and social ramifications of such a diagnosis. This work illuminated the personal and relational challenges faced by those living with HSV, particularly concerning safe sex practices and the dynamics of disclosure in intimate relationships. The insights from this study emphasized the need for healthcare providers to consider the emotional and social contexts of their patients when discussing HSV diagnosis and management.

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Continuing this exploration of relationships impacted by HSV, Hermes (Ann Hermes, 2017) utilized a grounded theory approach to examine how couples navigate discordant HSV statuses. This research highlighted the complexities of managing relationships when one partner is infected, revealing significant gaps in existing literature regarding the psychosocial effects of STDs on couples. Hermes's findings advocate for improved clinical practices that address the relational dimensions of STD management, thus enhancing the support provided to affected individuals and couples.

The systematic reviews conducted by Harfouche et al. (Harfouche et al., 2019) and Chaabane et al. (Chaabane et al., 2019) further expanded the understanding of HSV epidemiology across different regions, including Africa and the Middle East. These reviews synthesized data on the changing trends of HSV-1 and HSV-2 infections, demonstrating a notable shift in the etiology of genital herpes. They emphasized the importance of utilizing standardized serological assays to monitor these trends, which is crucial for public health strategies aimed at controlling HSV transmission.

Sukik et al. (Sukik et al., 2019) extended this analysis to Latin America and the Caribbean, providing a comprehensive overview of HSV-1 seroprevalence and its implications for genital herpes management. This systematic review highlighted the need for region-specific data to inform effective healthcare interventions and screening practices.

In 2021, AlMukdad et al. (AlMukdad et al., 2021) conducted a systematic review of HSV-2 epidemiology in Asia, contributing valuable insights into seroincidence and seroprevalence across diverse populations. Their findings reiterated the necessity of robust serological screening to understand the burden of HSV-2 and its implications for public health policies.

Finally, Lee et al. (Wen Huey Lee et al., 2022) provided a systematic review of healthcare resource utilization and costs associated with HSV diagnosis and management. Their analysis of 38 studies revealed substantial heterogeneity in reporting and highlighted the economic burden of HSV-related healthcare,

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

particularly in high-income countries. This review underscored the importance of understanding the resource implications of HSV management to inform healthcare policy and optimize resource allocation.

Together, these articles illustrate a comprehensive landscape of serologic screening for genital herpes, encompassing diagnostic methodologies, epidemiological trends, psychosocial impacts, and economic considerations. The evolution of research in this field underscores the importance of a multifaceted approach to understanding and managing genital herpes, ultimately aiming to improve public health outcomes and patient care.

LITERATURE REVIEW

The article titled "Detection, quantification and genotyping of Herpes Simplex Virus in cervicovaginal secretions by real-time PCR: a cross sectional survey" by (AN Aryee et al., 2005) presents a comprehensive examination of serologic screening methods for genital herpes, specifically focusing on the detection of Herpes Simplex Virus (HSV) in cervicovaginal secretions. The authors employ a real-time PCR technique, which is notable for its sensitivity and specificity in identifying HSV, particularly in the context of asymptomatic carriers who may not present with overt clinical manifestations.

One of the key insights from the article is the rigorous methodology used for serologic screening, including the detection of serum anti-HSV-2 IgG using the Kalon IgG kit. This approach underscores the importance of accurate serological testing in clinical settings, as it allows for the identification of individuals who may be at risk for transmission despite the absence of symptoms. The authors further enhance the reliability of their findings by subjecting reactive samples to additional testing, such as Monospecific ELISA for HIV-2 and Wellcozyme HIV Recombinant for HIV-1, thereby ensuring that co-infections do not confound the results.

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Moreover, the article highlights the multifaceted approach to diagnosing sexually transmitted infections (STIs) by incorporating various tests for other pathogens, including Hepatitis B and Treponema pallidum. The use of the MACRO-VUE Rapid Plasma Reagin test for syphilis, alongside the Abbott Determine™ HBsAg qualitative immunoassay for Hepatitis B, illustrates a thorough screening protocol that addresses the complexity of STI diagnostics. The authors also utilize statistical analysis through software such as EPIInfo, SPSS, and Minitab, which adds a layer of credibility to their data interpretation and conclusions.

However, while the article provides valuable insights into the methodologies for serologic screening, it could benefit from a more detailed discussion on the implications of these findings for public health strategies and clinical practice. The authors briefly mention the importance of identifying asymptomatic carriers, yet a deeper exploration of how these screening methods can influence disease management and prevention strategies in high-risk populations would enhance the overall impact of the research.

The article titled "Seroprevalence, predictors and estimated incidence of maternal and neonatal Herpes Simplex Virus Type 2 infection in semi-urban women in Kilifi, Kenya" by (U. Nyiro et al., 2011) presents a comprehensive analysis of the seroprevalence and risk factors associated with Herpes Simplex Virus Type 2 (HSV-2) infection in a specific demographic. The study highlights critical insights into the epidemiology of HSV-2, particularly in pregnant women, and its implications for maternal and neonatal health.

One of the key findings of the article is the high seroprevalence of HSV-2 among the study population, which underscores the significant burden of this infection in semi-urban settings. The authors provide a detailed examination of the predictors of HSV-2 infection, revealing that certain demographic and behavioral factors, such as age, marital status, and sexual history, are strongly associated with the likelihood of being seropositive. This information is vital for public health strategies aimed at reducing transmission rates, particularly in populations at higher risk.

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Furthermore, the article discusses the performance of various serological assays, including the Focus ELISA tests and HerpeSelect assays, in detecting HSV-2 antibodies. The evaluation of these diagnostic tools is particularly relevant, as accurate serologic screening is essential for identifying undiagnosed infections and implementing appropriate interventions. The authors detail the sensitivity and specificity of these tests, providing a valuable resource for clinicians and researchers working in the field of sexually transmitted infections.

The relationship between HSV-2 infection and HIV susceptibility is another critical aspect addressed in the study. The authors note that co-infection with HSV-2 may increase the risk of acquiring HIV, highlighting the need for integrated screening and prevention strategies. This association is particularly pertinent for public health initiatives in regions where both infections are prevalent, as it emphasizes the importance of addressing multiple sexually transmitted infections concurrently.

The article titled "This is herpes: A qualitative investigation of the disclosure patterns of HSV2 individuals" by (Miles, 2017) provides a nuanced exploration of the personal experiences of individuals diagnosed with HSV-2, particularly focusing on how their diagnosis affects their sexual practices and the implications of disclosing their status to partners.

Miles employs a qualitative methodology, allowing for in-depth insights into the lived experiences of participants. The findings indicate that the diagnosis of HSV-2 significantly alters individuals' approaches to sexual activity. Many participants reported a heightened awareness of safe sex practices post-diagnosis, which often included increased use of condoms and a more cautious approach to sexual encounters. This shift in behavior underscores the importance of education and awareness regarding sexually transmitted infections (STIs) and their implications for sexual health.

Furthermore, the article delves into the complexities surrounding the disclosure of one's HSV-2 status. Participants expressed varying degrees of comfort and anxiety about sharing their diagnosis with potential partners. The act of disclosure was often

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

fraught with concerns about stigma and rejection, which can significantly impact one's sexual relationships. This aspect of the study highlights the psychological burden that accompanies the diagnosis of HSV-2 and the need for supportive frameworks that encourage open communication about sexual health.

Miles' work contributes to the broader discourse on sexual health by emphasizing the importance of both safe sex practices and the social dynamics of disclosure in the context of STIs. The qualitative nature of the research allows for a rich understanding of the emotional and relational dimensions of living with HSV-2, which is often overlooked in more quantitative studies.

The article titled "How Discordant HSV Status Impacts Dyadic Relationships: A Grounded Theory Study" by (Ann Hermes, 2017) presents a nuanced exploration of the relational dynamics faced by couples where one partner is HSV-positive and the other is HSV-negative. The research is significant as it addresses a notable gap in the existing literature concerning the psychological and relational implications of herpes simplex virus (HSV) infections, particularly in dyadic contexts.

Hermes employs a constructivist grounded theory approach, which allows for an in-depth understanding of how couples navigate the complexities introduced by discordant HSV statuses. One of the key insights from the study is the difficulty that mental health professionals face in conceptualizing the impact of STDs on relationships due to the lack of focused research in this area. This gap is critical as it underscores the need for clinicians to develop a better understanding of the emotional and relational challenges posed by HSV, which can significantly affect communication, intimacy, and overall relationship satisfaction.

The article highlights the prevalence of HSV, noting that approximately 5% of individuals aged 14-49 are affected by genital HSV-2, with around 776,000 new infections occurring annually. This statistic not only contextualizes the importance of the study but also emphasizes the potential for widespread relational issues stemming from HSV infections. Furthermore, Hermes discusses the severe implications of HSV during childbirth, including the risk of neonatal HSV, which

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

can lead to serious health complications for infants. This aspect of the research stresses the urgency of addressing HSV in clinical settings, particularly for couples planning to start families.

Moreover, the article draws attention to the immunological implications of HSV, particularly the increased CD4 cell counts in individuals with HSV, which heightens their risk of contracting HIV. This intersection of sexual health and mental health is crucial, as it indicates that the ramifications of HSV extend beyond relational dynamics and into broader public health concerns.

The article titled "Epidemiology of herpes simplex virus type 2 in Asia: A systematic review, meta-analysis, and meta-regression" by (AlMukdad et al., 2021) provides a comprehensive overview of the epidemiological landscape of herpes simplex virus type 2 (HSV-2) in Asia. The systematic review and meta-analysis methodology employed in this study is a significant strength, allowing for a robust synthesis of existing data on HSV-2 seroincidence and seroprevalence across the region.

The authors meticulously screened a wide array of literature, focusing on studies that presented primary data on at least ten subjects related to HSV-2 outcomes. This rigorous selection process ensured that the included studies were relevant and of sufficient quality, thereby enhancing the reliability of the findings. The exclusion of case reports, commentaries, reviews, and qualitative studies further underscores the authors' commitment to a quantitative approach, which is crucial for drawing statistically valid conclusions regarding serologic screening for genital herpes.

One of the key insights from the article is the identification of varying rates of HSV-2 seroprevalence across different Asian countries. This variability highlights the need for region-specific strategies in serologic screening and public health interventions. The meta-regression analysis conducted by the authors also sheds light on potential demographic and behavioral factors influencing HSV-2 transmission, which could inform targeted prevention efforts.

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

However, the article could benefit from a deeper exploration of the implications of these epidemiological findings for clinical practice, particularly concerning serologic screening protocols for genital herpes. While the authors provide a solid overview of HSV-2 epidemiology, a discussion on how these insights translate into actionable guidelines for healthcare providers would enhance the practical relevance of the research. Additionally, the impact of sociocultural factors on the acceptance and implementation of serologic screening in different Asian contexts could be explored further to provide a more holistic understanding of the challenges and opportunities in addressing genital herpes.

The article titled "Healthcare resource utilisation pattern and costs associated with herpes simplex virus diagnosis and management: a systematic review" by (Wen Huey Lee et al., 2022) provides a comprehensive analysis of the healthcare resource utilization and associated costs tied to the diagnosis and management of herpes simplex virus (HSV). The systematic review synthesizes findings from 38 studies, highlighting a significant gap in the literature regarding the economic implications of HSV management, particularly in low- and middle-income countries (LMICs).

The authors conducted an extensive search that initially identified 11,443 articles, ultimately narrowing the focus to 38 relevant studies after rigorous screening. This methodological rigor enhances the credibility of the findings, ensuring that the included studies are pertinent to the topic of HSV management. Notably, the review distinguishes between studies that solely report resource utilization, those that focus on costs, and those that address both. This categorization allows for a nuanced understanding of the various aspects of HSV management, which is crucial for healthcare providers and policymakers.

Among the key insights, the review underscores that while a substantial number of studies report on costs related to HSV care, the majority are derived from high-income countries (HICs), with only one study originating from an LMIC. This disparity suggests a need for more research in diverse economic settings to better understand the global burden of HSV and inform resource allocation. Furthermore,

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

the review highlights that laboratory testing costs are a critical component in the diagnosis of HSV, yet only three studies specifically addressed these costs, indicating another area ripe for further investigation.

CONCLUSION

The literature on serologic screening for genital herpes has seen significant advancements, emphasizing the importance of comprehensive methodologies, epidemiological insights, psychosocial impacts, and economic considerations. (AN Aryee et al., 2005) laid the groundwork by establishing effective diagnostic methodologies, utilizing real-time PCR alongside serological tests like ELISA and rapid plasma reagin to detect HSV in cervicovaginal secretions. This foundational work highlighted the necessity of accurate serological testing in identifying asymptomatic carriers, which is crucial for effective public health strategies ((AN Aryee et al., 2005)).

(U. Nyiro et al., 2011) expanded the understanding of HSV-2 epidemiology by investigating seroprevalence and transmission dynamics among semi-urban women in Kenya. Their findings revealed significant risk factors for undiagnosed infections and underscored the correlation between HSV-2 and HIV, suggesting that integrated screening approaches are essential for populations at higher risk. This aligns with the work of (AlMukdad et al., 2021), who conducted a systematic review of HSV-2 epidemiology in Asia, emphasizing the need for robust serological screening to inform public health policies.

(Miles, 2017) provided a qualitative perspective on the psychological and social ramifications of HSV-2 diagnoses, particularly regarding safe sex practices and disclosure dynamics in intimate relationships. This study, complemented by (Ann Hermes, 2017), which focused on couples with discordant HSV statuses, highlighted the relational complexities and emotional challenges faced by individuals living with HSV. These insights stress the importance of considering the psychosocial dimensions of HSV management in clinical practice.

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The systematic reviews by (Harfouche et al., 2019) and (Chaabane et al., 2019) further elucidated HSV epidemiology across different regions, advocating for the use of standardized serological assays to monitor trends in HSV-1 and HSV-2 infections. (Sukik et al., 2019) extended this analysis to Latin America and the Caribbean, emphasizing the need for region-specific data to inform effective healthcare interventions.

Finally, (Wen Huey Lee et al., 2022) provided an essential overview of healthcare resource utilization and costs associated with HSV diagnosis and management. Their findings highlighted the economic burden of HSV-related healthcare, particularly in high-income countries, underscoring the importance of understanding resource implications to optimize healthcare policies.

In conclusion, the body of literature on serologic screening for genital herpes presents a multifaceted understanding of the complexities surrounding HSV infections. The integration of diagnostic methodologies, epidemiological data, psychosocial insights, and economic evaluations indicates a pressing need for comprehensive public health strategies that address the various dimensions of genital herpes. Such an approach is crucial for improving patient care and ultimately reducing the burden of HSV infections globally.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. AN Aryee, E., L Bailey, R., Natividad-Sancho, A., Kaye, S., & J Holland, M., 2005. Detection, quantification and genotyping of Herpes Simplex

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Virus in cervicovaginal secretions by real-time PCR: a cross sectional survey.. [PDF]

2. U. Nyiro, J., J. Sanders, E., Ngetsa, C., Wale, S., Awuondo, K., Anne Bukusi, E., A. Price, M., N. Amornkul, P., & James Nokes, D., 2011. Seroprevalence, predictors and estimated incidence of maternal and neonatal Herpes Simplex Virus Type 2 infection in semi-urban women in Kilifi, Kenya. [PDF]
3. Ann Hermes, H., 2017. How Discordant HSV Status Impacts Dyadic Relationships: A Grounded Theory Study. [PDF]
4. Harfouche, M., Chemaitelly, H., & J Abu-Raddad, L., 2019. Herpes simplex virus type 1 epidemiology in Africa: Systematic review, meta-analyses, and meta-regressions.. [PDF]
5. Chaabane, S., Harfouche, M., Chemaitelly, H., Schwarzer, G., & J Abu-Raddad, L., 2019. Herpes simplex virus type 1 epidemiology in the Middle East and North Africa: systematic review, meta-analyses, and meta-regressions.. [PDF]
6. Sukik, L., Alyafei, M., Harfouche, M., & J. Abu-Raddad, L., 2019. Herpes simplex virus type 1 epidemiology in Latin America and the Caribbean: Systematic review and meta-analytics. ncbi.nlm.nih.gov
7. AlMukdad, S., Harfouche, M., Wettstein, A., & J. Abu-Raddad, L., 2021. Epidemiology of herpes simplex virus type 2 in Asia: A systematic review, meta-analysis, and meta-regression. ncbi.nlm.nih.gov
8. Wen Huey Lee, S., L Gottlieb, S., & Chaiyakunapruk, N., 2022. Healthcare resource utilisation pattern and costs associated with herpes simplex virus diagnosis and management: a systematic review. ncbi.nlm.nih.gov
9. Miles, K., 2017. This is herpes: A qualitative investigation of the disclosure patterns of HSV2 individuals. [PDF]

Downloaded from The Medical Journal of Clinical Pathology and Microbiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved