Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Dentistry of Oral Maxillofacial Surgery



Cervical Necrotizing Fasciitis Of Odontogenic Origin

[1] Fahmi, [1] Lisa

[1] Faculty of Dentistry, Padjadjaran University, Bandung City, West Java, Indonesia

Correspondence : drg.fahmi@gmail.com

Article History :

Received date : 2024/07/26
Revised date : 2024/08/17
Accepted date : 2024/09/08
Published date : 2025/01/24

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: Cervical necrotizing fasciitis (CNF) of odontogenic origin is a severe condition characterized by rapid soft tissue necrosis, primarily resulting from dental infections. The literature indicates that early diagnosis and intervention are paramount to improving patient outcomes, as this condition poses significant challenges due to its aggressive nature and potential for high mortality rates. **Literature Review:** The case presented by (Cecchini et al., 2021) further reinforces the notion that even healthy individuals can develop CNF from odontogenic sources, such as wisdom tooth abscesses. This underscores the need for swift medical and surgical interventions to ensure successful recovery. The multifactorial nature of odontogenic infections, as discussed by (Pinkston et al., 2022), highlights the polymicrobial characteristics of these infections and the critical importance of prompt surgical intervention to prevent complications like airway obstruction and pericarditis. The aggressive nature of odontogenic infections leading to CNF is also examined by (L Huff et al., 2022), who emphasizes the need for rapid diagnosis and aggressive surgical management, especially when infections are located near vital anatomical structures. Additionally, (Shankar Chaurasiya et al., 2022) identifies Pseudomonas aeruginosa as a rare but serious pathogen in CNF cases, particularly among diabetic patients, further stressing the need for early intervention. In pediatric cases, as discussed by (Simona Coșarcă et al., 2023), the rarity of CNF poses unique challenges, and the case of a 12-year-old boy with a dental origin highlights the potential for rapid progression. This necessitates a comprehensive understanding of the disease and its management in younger populations. **Conclusion:** In conclusion, the reviewed literature collectively emphasizes the critical importance of early diagnosis, aggressive intervention, and interdisciplinary management in treating cervical necrotizing fasciitis of odontogenic origin. The varied case studies illustrate that both healthy individuals and those with underlying health conditions are at risk, necessitating heightened awareness among healthcare providers regarding the potential for rapid disease progression and severe complications.

Keyword: Cervical Necrotizing Fasciitis, Odontogenic Origin

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

Cervical necrotizing fasciitis (CNF) of odontogenic origin is a grave condition characterized by the rapid necrosis of soft tissue, often stemming from dental infections. The literature surrounding this topic underscores the critical nature of early diagnosis and intervention, as well as the multifaceted challenges presented by this disease.

In 2014, (Song et al., 2014) provided a compelling case study illustrating the complexities of managing CNF in a patient with renal failure and MRSA infection, who was treated with an interdisciplinary approach that included adjusted doses of vancomycin. Their findings highlight the necessity of tailored antibiotic therapies in high-risk patients to prevent severe complications.

Two years later, (Yenigun et al., 2016) expanded on the understanding of necrotizing fasciitis by discussing its association with compromised immune systems, particularly in cases linked to sialadenitis. Their research emphasizes the importance of recognizing the fulminant nature of the disease, where early diagnosis can significantly alter outcomes, especially given the mortality rates associated with cervical necrotizing fasciitis, which range from 7% to 20%.

(Cecchini et al., 2021) presented a noteworthy case of a healthy young female who developed CNF from a wisdom tooth abscess, underscoring that even individuals without predisposing factors can be affected. Their study reinforces the idea that swift medical and surgical interventions are crucial for successful recovery in such cases.

The multifactorial nature of odontogenic infections was further explored by (Pinkston et al., 2022), who documented a case of necrotizing mediastinitis following dental extraction. They emphasized the polymicrobial nature of these infections and the need for prompt surgical intervention to prevent high mortality rates associated with complications such as airway obstruction and pericarditis.

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

In the same year, (L Huff et al., 2022) examined a severe case of odontogenic infection leading to necrotizing fasciitis of the anterior chest wall and neck. Their findings reinforced the notion that rapid diagnosis and aggressive surgical management are imperative, particularly given the aggressive nature of these infections and their proximity to vital anatomical structures.

(Shankar Chaurasiya et al., 2022) added to the discourse by identifying Pseudomonas aeruginosa as a rare but potentially fatal cause of cervical necrotizing fasciitis, particularly in diabetic patients. Their case study highlighted the critical importance of early intervention in improving patient outcomes, drawing attention to the polymicrobial characteristics of these infections.

(Simona Coșarcă et al., 2023) introduced the unique challenges presented by pediatric cases of cervical necrotizing fasciitis, presenting a case of a 12-year-old boy with a dental origin. Their study illustrated the rarity of such occurrences in children and the complexities involved in management due to the potential for rapid progression.

Most recently, (A Ba-shammakh et al., 2023) examined the role of vacuum-assisted closure therapy in managing neck necrotizing fasciitis. Their case study underscored the aggressive nature of this condition and the necessity of comprehensive management strategies, including surgical debridement and ongoing monitoring for complications post-discharge.

Together, these articles present a comprehensive overview of the complexities associated with cervical necrotizing fasciitis of odontogenic origin, highlighting the critical importance of prompt diagnosis, aggressive intervention, and interdisciplinary management to improve patient outcomes.

LITERATURE REVIEW

The article titled "Cervical Necrotizing Fasciitis Caused by Dental Infection" by (Song et al., 2014) provides a comprehensive overview of a case involving cervical necrotizing fasciitis originating from a dental infection. The authors present a

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

detailed account of the clinical management and interdisciplinary approach taken to address the complexities of the case, particularly in the context of the patient's renal failure and methicillin-resistant Staphylococcus aureus (MRSA) infection.

A critical evaluation of the material reveals several key insights into the management of cervical necrotizing fasciitis. The authors highlight the significant challenge posed by the patient's renal failure, which precluded the use of standard antibiotic therapies. This limitation necessitated a tailored approach to antibiotic administration, employing adjusted doses of vancomycin. The successful implementation of this strategy underscores the importance of individualized treatment plans in cases where standard protocols may not be applicable due to underlying health conditions. The consultation with specialists in infectious diseases and nephrology exemplifies the necessity of an interdisciplinary approach in managing complex medical cases, particularly those involving severe infections.

Furthermore, the article notes the absence of diabetes mellitus (DM) in the patient, a factor often associated with delayed healing and increased risk of complications. The authors assert that the lack of DM contributed positively to the patient's recovery trajectory, allowing for a more favorable outcome following the interdisciplinary intervention. This observation emphasizes the need for thorough patient assessments to identify potential risk factors that could complicate treatment and recovery.

The article "Nonodontogenic Cervical Necrotizing Fasciitis Caused by Sialadenitis" by (Yenigun et al., 2016) provides a comprehensive examination of necrotizing fasciitis, particularly emphasizing its rapid progression and the critical nature of early diagnosis and intervention. The authors detail the clinical features that characterize this condition, such as the fulminant progression of symptoms and the presence of grey-black necrotic areas, which serve as key indicators for healthcare professionals.

In their analysis, the authors highlight that necrotizing fasciitis can arise from localized infections, often following minor trauma, which underscores the importance of vigilance in clinical settings where such injuries may occur. The

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

article notes that while necrotizing fasciitis is a serious condition, its occurrence in the head and neck region is relatively rare, making awareness and prompt diagnosis particularly crucial in these cases. The reported mortality rates ranging from 7% to 20% further illustrate the potentially fatal nature of the disease if not addressed swiftly.

The authors effectively communicate the urgency required in managing necrotizing fasciitis, as delayed treatment can lead to severe complications and increased mortality. Their findings suggest that clinicians should maintain a high index of suspicion for necrotizing fasciitis in patients presenting with rapid deterioration of soft tissue, especially in the context of local infections.

The article titled "Odontogenic Infection Complicated by Cervicofacial Necrotizing Fasciitis in a Healthy Young Female" by (Cecchini et al., 2021) provides a compelling examination of a rare but critical condition: necrotizing fasciitis (NF) stemming from odontogenic infections. The authors effectively highlight the severity of NF, particularly in the head and neck region, where it is not only rare but also associated with a high mortality rate. This underscores the importance of prompt recognition and intervention in such cases.

The case presented involves a young female patient who, notably, had no significant risk factors typically associated with NF. This aspect is particularly noteworthy as it challenges the common perception that NF predominantly affects individuals with pre-existing comorbidities. The authors emphasize that dental infections, specifically those originating from wisdom tooth abscesses, are among the most frequent causes of NF, suggesting that dental practitioners should maintain a high index of suspicion for such complications in otherwise healthy individuals.

One of the critical insights from the article is the emphasis on early diagnosis and intervention. The authors detail how the timely administration of appropriate medical therapy combined with surgical intervention led to a successful outcome for the patient. This aspect of the study reinforces the need for healthcare providers to be vigilant in monitoring patients with odontogenic infections, particularly when symptoms escalate or when there is evidence of rapid progression.

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The authors also address the broader implications of their findings, suggesting that while NF of the head and neck is uncommon, its potential for rapid deterioration necessitates immediate action. The mortality rates associated with NF in patients with comorbidities serve as a stark reminder of the risks involved, but the case presented illustrates that even healthy individuals can face severe outcomes if NF is not swiftly addressed.

The article "Necrotizing Mediastinitis Following Dental Extraction: A Case Report" by (Pinkston et al., 2022) provides a critical examination of the severe complications that can arise from odontogenic infections, particularly focusing on the phenomenon of descending necrotizing mediastinitis. This condition, as highlighted in the article, is a grave consequence of polymicrobial infections originating from dental sources, which can traverse the cervical fascial planes, leading to life-threatening outcomes.

The authors effectively outline the pathophysiology of how odontogenic infections can escalate into necrotizing fasciitis and further into the mediastinum. They emphasize the importance of recognizing the polymicrobial nature of these infections, which complicates treatment and management strategies. The article underscores that the initial source of the infection often stems from dental procedures, such as extractions, which may inadvertently allow for bacterial dissemination into deeper tissue layers.

Moreover, the authors discuss the clinical implications of delayed diagnosis and treatment, which can result in significant morbidity and mortality. The reported mortality rates of 25–40%, and as high as 60% in some cases, reflect the urgency of prompt intervention. This high mortality rate persists despite advancements in diagnostic imaging, such as computed tomography (CT), and improvements in surgical techniques, antibiotic therapies, and intensive care management. The article effectively conveys that while these advancements have improved patient outcomes, they are not foolproof, and the risk of fatality remains a pressing concern.

The article also highlights the critical role of surgical subspecialties in the management of such infections. Prompt identification and drainage of the source of

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

the odontogenic infection are essential to mitigate the risks associated with descending necrotizing mediastinitis. The authors advocate for a multidisciplinary approach, emphasizing that collaboration among dental, surgical, and medical teams is vital to improving patient prognosis.

The article titled "A Severe Case of Odontogenic Infection and Necrotizing Fasciitis of the Anterior Chest Wall and Neck" by (L Huff et al., 2022) presents a critical examination of the complexities surrounding cervical necrotizing fasciitis (CCNF) stemming from odontogenic infections. The authors emphasize the need for early identification and immediate surgical intervention, which are pivotal in enhancing patient outcomes in cases of necrotizing fasciitis.

The article underscores the aggressive nature of odontogenic infections, which can rapidly progress to CCNF, particularly due to the anatomical proximity of these infections to vital structures in the head and neck. This proximity necessitates a heightened awareness among clinicians, as the consequences of delayed diagnosis can be dire. The authors highlight the atypical presentation of their case—a young man without significant past medical history—contrasting with the common demographic of elderly males who often present with preexisting immunocompromised conditions. This observation serves to broaden the understanding of risk factors associated with CCNF, suggesting that age and health status should not be the sole criteria for assessment.

Furthermore, the study reinforces the principles of infection control and wound care, which are crucial in the management of necrotizing fasciitis. The authors provide a clear rationale for the implementation of aggressive surgical debridement and the importance of multidisciplinary approaches in treating such infections. By detailing the clinical management of the presented case, the authors contribute valuable insights into the treatment protocols that can significantly improve survival rates.

The article titled "Pseudomonas aeruginosa as a culprit of cervical necrotizing fasciitis: A case report" by (Shankar Chaurasiya et al., 2022) provides a critical examination of a rare and serious condition known as cervical necrotizing fasciitis,

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

specifically highlighting its odontogenic origins. This case report underscores the significance of recognizing the polymicrobial nature of necrotizing fasciitis, a condition often arising from dental infections that can lead to severe complications in patients, particularly those with underlying health issues such as diabetes.

The authors detail a case where Pseudomonas aeruginosa was identified as the primary pathogen responsible for the cervical necrotizing fasciitis in a diabetic patient. This identification is crucial as it sheds light on the atypical presentations of this infection, which is predominantly associated with multiple microbial agents. The rarity of Pseudomonas aeruginosa as a sole pathogen in such cases emphasizes the need for heightened clinical awareness and consideration of this organism in similar presentations, especially in immunocompromised individuals.

The article also discusses the clinical implications of odontogenic infections as potential foci for cervical necrotizing fasciitis. Odontogenic infections can often progress rapidly, leading to significant morbidity. The authors provide a comprehensive overview of the pathophysiology involved, illustrating how an initial dental infection can escalate into a life-threatening condition if not promptly addressed. This emphasizes the importance of early diagnosis and intervention, as well as the need for a multidisciplinary approach in managing such cases.

Furthermore, the report highlights the diagnostic challenges and treatment protocols necessary for effectively managing cervical necrotizing fasciitis. The authors advocate for aggressive surgical intervention combined with appropriate antibiotic therapy, which is vital in controlling the spread of the infection and preventing further complications.

The article "Pediatric Cervicofacial Necrotizing Fasciitis—A Challenge for a Medical Team" by (Simona Coșarcă et al., 2023) provides a comprehensive overview of cervical necrotizing fasciitis (NF) of odontogenic origin, particularly within the pediatric population. The authors highlight the rarity of this condition, especially in children, and emphasize its potential to arise from dental complications, specifically pericoronitis associated with the second inferior molar.

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The case study presented in the article illustrates a 12-year-old boy who developed this severe infection following pericoronitis, underlining the critical link between dental health and systemic infections. The authors effectively detail the pathophysiology of NF, noting its rapid progression and the potential for serious complications, including mediastinal involvement. This aspect is particularly alarming, as it underscores the urgent need for prompt diagnosis and intervention in cases of odontogenic infections that may escalate to necrotizing fasciitis.

(Simona Coșarcă et al., 2023) also provide insight into the microbiological classification of NF, distinguishing between type 1 (polymicrobial) and type 2 (monomicrobial) infections. They note that polymicrobial infections are more prevalent, which is an important consideration for clinicians when determining appropriate antibiotic therapy and surgical intervention strategies. The multidisciplinary approach advocated by the authors is essential, as the management of NF often requires collaboration among various specialties, including surgery, infectious disease, and critical care.

One notable strength of the article is its emphasis on the challenges faced during treatment, particularly in the context of prolonged intubation and the complexities of managing a young patient with a severe infection. This highlights the need for careful monitoring and supportive care, which are crucial for improving patient outcomes in such critical scenarios.

However, while the article provides valuable insights into this rare condition, it could benefit from a more extensive discussion on the long-term implications of cervical necrotizing fasciitis in pediatric patients. Additionally, the authors might consider elaborating on preventative measures that could reduce the incidence of such infections arising from odontogenic sources.

The article titled "Vacuum-Assisted Closure Therapy: A Lifesaver in a Case of Neck Necrotizing Fasciitis" by (A Ba-shammakh et al., 2023) provides a comprehensive examination of cervical necrotizing fasciitis (CNF), emphasizing its association with odontogenic infections. The authors effectively highlight the aggressive nature of CNF, a severe form of necrotizing soft tissue infections

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

(NSTIs), characterized by rapid tissue disintegration affecting various anatomical structures, including skeletal muscle and subcutaneous fat.

One of the key insights presented in the article is the identification of odontogenic infections, particularly those stemming from the second and third mandibular molars, as significant contributors to the development of CNF. The authors detail how infections in these dental regions can infiltrate the submandibular space and extend through fascial planes, leading to complications such as mediastinitis. This progression underscores the critical need for timely diagnosis and intervention, especially in patients with predisposing chronic conditions like diabetes mellitus and obesity, which are frequently associated with poorer outcomes.

The article further discusses the necessity of a multifaceted therapeutic approach to manage CNF effectively. The authors advocate for an integrated strategy that includes hemodynamic support, broad-spectrum antibiotics, and surgical debridement. This holistic approach is crucial given the rapid deterioration that can occur with CNF, necessitating swift and decisive action to prevent further systemic involvement.

Additionally, the authors introduce vacuum-assisted closure therapy as a potential lifesaving intervention in the management of CNF. This innovative technique may enhance wound healing and reduce the risk of further infection, presenting an important advancement in the treatment of this condition. The incorporation of such modern therapeutic modalities reflects a growing recognition of the need for adaptive and effective treatment strategies in the face of complex infections.

CONCLUSION

Cervical necrotizing fasciitis (CNF) of odontogenic origin is a severe condition characterized by rapid soft tissue necrosis, primarily resulting from dental infections. The literature indicates that early diagnosis and intervention are paramount to improving patient outcomes, as this condition poses significant challenges due to its aggressive nature and potential for high mortality rates.

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The case study by (Song et al., 2014) illustrates the complexities involved in managing CNF, particularly in patients with underlying health concerns such as renal failure and MRSA infection. The study emphasizes the necessity of tailored antibiotic therapies in high-risk patients, highlighting the importance of interdisciplinary approaches to treatment. Similarly, (Yenigun et al., 2016) expands on the significance of early diagnosis, particularly in cases associated with compromised immune systems, where the mortality rates can range from 7% to 20%.

The case presented by (Cecchini et al., 2021) further reinforces the notion that even healthy individuals can develop CNF from odontogenic sources, such as wisdom tooth abscesses. This underscores the need for swift medical and surgical interventions to ensure successful recovery. The multifactorial nature of odontogenic infections, as discussed by (Pinkston et al., 2022), highlights the polymicrobial characteristics of these infections and the critical importance of prompt surgical intervention to prevent complications like airway obstruction and pericarditis.

The aggressive nature of odontogenic infections leading to CNF is also examined by (L Huff et al., 2022), who emphasizes the need for rapid diagnosis and aggressive surgical management, especially when infections are located near vital anatomical structures. Additionally, (Shankar Chaurasiya et al., 2022) identifies Pseudomonas aeruginosa as a rare but serious pathogen in CNF cases, particularly among diabetic patients, further stressing the need for early intervention.

In pediatric cases, as discussed by (Simona Coșarcă et al., 2023), the rarity of CNF poses unique challenges, and the case of a 12-year-old boy with a dental origin highlights the potential for rapid progression. This necessitates a comprehensive understanding of the disease and its management in younger populations.

Lastly, (A Ba-shammakh et al., 2023) introduces vacuum-assisted closure therapy as an innovative management strategy for CNF, underscoring the need for comprehensive and adaptive treatment approaches in the face of complex infections.

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

In conclusion, the reviewed literature collectively emphasizes the critical importance of early diagnosis, aggressive intervention, and interdisciplinary management in treating cervical necrotizing fasciitis of odontogenic origin. The varied case studies illustrate that both healthy individuals and those with underlying health conditions are at risk, necessitating heightened awareness among healthcare providers regarding the potential for rapid disease progression and severe complications.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Song, C. W., Yoon, H. J., Jung, D. W., & Lee, S. H., 2014. Cervical Necrotizing Fasciitis Caused by Dental Infection. ncbi.nlm.nih.gov
2. Yenigun, A., Veyseller, B., Vural, O., & Ozturan, O., 2016. Nonodontogenic Cervical Necrotizing Fasciitis Caused by Sialadenitis. ncbi.nlm.nih.gov
3. Cecchini, A., J Cox, C., A Cecchini, A., Solanki, K., & McSharry, R., 2021. Odontogenic Infection Complicated by Cervicofacial Necrotizing Fasciitis in a Healthy Young Female. ncbi.nlm.nih.gov
4. Pinkston, J., A. Khoury, C., & D. Raper, J., 2022. Necrotizing Mediastinitis Following Dental Extraction: A Case Report. ncbi.nlm.nih.gov
5. L Huff, M., S Wilson, K., E Kane, K., L Wheel, K., & J Stirparo, J., 2022. A Severe Case of Odontogenic Infection and Necrotizing Fasciitis of the Anterior Chest Wall and Neck. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

6. Shankar Chaurasiya, P., Gurung, S., Karki, S., Timilsina, B., Shah, R., & Neupane, S., 2022. Pseudomonas aeruginosa as a culprit of cervical necrotizing fasciitis: A case report. ncbi.nlm.nih.gov
7. Simona Coșarcă, A., Száva, D., Bögözi, B., Iacob, A., Frățilă, A., & Sergiu, G., 2023. Pediatric Cervicofacial Necrotizing Fasciitis—A Challenge for a Medical Team. ncbi.nlm.nih.gov
8. A Ba-shammakh, S., M Alrayes, B., M Abu-obead, H., A Ba-Shammakh, M., M Al-Mohd, F., H Ramzoun, E., & Abu-Nowar, L., 2023. Vacuum-Assisted Closure Therapy: A Lifesaver in a Case of Neck Necrotizing Fasciitis. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Dentistry and Oral Maxillofacial Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved