Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Dermatology and Venereology



# Association of Vitiligo and Alopecia Areata With Atopic Dermatitis

[1] Pertiwi, [1] Devi

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.pertiwi29387@gmail.com

**Article History :**

Received date : 2024/10/11
Revised date : 2024/15/14
Accepted date : 2024/12/09
Published date : 2025/01/07

***Copyright:*** *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

## ABSTRACT

**Background:** The literature concerning the association of vitiligo and alopecia areata with atopic dermatitis reveals a multifaceted interplay of genetic, immunological, and environmental factors. The studies reviewed indicate a significant overlap in the pathogenesis of these autoimmune conditions, suggesting that they may share common underlying mechanisms. **Literature Review:** The characterization of alopecia areata and vitiligo as "sister diseases" by (Popa et al., 2023) further elucidates their relationship, highlighting associations with atopic dermatitis and asthma. The study advocates for thyroid hormone assessments in alopecia areata patients, reflecting the intricate connections among various autoimmune conditions. A large-scale retrospective analysis by (Ezzedine et al., 2023) reveals a higher incidence of autoimmune comorbidities, including alopecia areata, among patients with vitiligo, emphasizing the multifaceted nature of vitiligo as an autoimmune disorder. This study also points to significant correlations between vitiligo and psychological conditions, further complicating its clinical presentation. The exploration of the aetiopathogenesis of vitiligo by (Boniface, 2023) highlights the role of immune responses and genetic determinants in disease susceptibility, offering insights into shared pathogenic mechanisms with alopecia areata. Lastly, the discussion of dietary influences on skin health by (Sharma et al., 2024) suggests that nutritional interventions may play a role in managing both alopecia areata and vitiligo, linking these disorders through lifestyle factors. **Conclusion:** In conclusion, the literature presents a compelling narrative that underscores the intricate relationships between vitiligo, alopecia areata, and atopic dermatitis. The shared genetic and immunological landscape highlighted in these studies warrants further exploration, particularly regarding the underlying mechanisms that contribute to the co-occurrence of these autoimmune conditions.

**Keyword:** Vitiligo, Alopecia Areata, Atopic Dermatitis



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

## INTRODUCTION

The association between vitiligo and alopecia areata, particularly in the context of autoimmune mechanisms, has been a focal point of recent dermatological research. The literature reveals a complex interplay of genetic, immunological, and environmental factors that contribute to the pathogenesis of these conditions.

(Meredith Petukhova, 2013) provides foundational insights into alopecia areata, highlighting its increased prevalence among individuals with psychiatric disorders and its association with autoimmune conditions. The study emphasizes the emotional disturbances linked to alopecia areata, suggesting that stress and psychological factors may exacerbate the condition. Furthermore, it identifies a history of atopy or autoimmunity as a risk factor for developing alopecia areata, setting the stage for understanding its relationship with vitiligo.

In the same year, (Riad et al., 2013) explore the anatomical colocalization of alopecia areata and vitiligo, documenting cases where one condition precedes the other. Their findings from a genome-wide association study reveal shared susceptibility loci, indicating a genetic overlap between the two diseases. This work underscores the notion that the mechanisms underlying follicular melanocyte destruction in both conditions may be interrelated, thereby reinforcing the link between alopecia areata and vitiligo.

(Baldini et al., 2017) expand on the autoimmune aspect by investigating the role of Th17-related cytokines in vitiligo. They discuss how dysregulation of melanocyte function can lead to depigmentation, further supporting the idea of shared autoimmune pathways between vitiligo and alopecia areata. Their study emphasizes the prevalence of comorbid autoimmune diseases in patients with vitiligo, including alopecia areata, thus highlighting the interconnectedness of these disorders.

(Anbar et al., 2022) contribute to this discourse by examining the videodermoscopic changes in vitiligo lesions, identifying similarities in pathogenesis with alopecia



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

areata. Their research suggests that both conditions may share common genetic risk factors and exhibit colocalization, particularly in scalp regions. This reinforces the idea that both diseases may arise from similar immunological disturbances.

(Popa et al., 2023) further elucidate the relationship between alopecia areata and vitiligo, referring to them as “sister diseases.” They identify atopic dermatitis and asthma as additional comorbidities associated with alopecia areata, suggesting a broader autoimmune background. Their findings advocate for the assessment of thyroid hormones in patients with alopecia areata, reflecting the intricate connections among various autoimmune conditions.

(Ezzedine et al., 2023) provide a large-scale retrospective analysis that underscores the higher incidence of autoimmune comorbidities, including alopecia areata, among patients with vitiligo. Their study reveals significant correlations between vitiligo and various psychological conditions, thus reinforcing the notion that vitiligo is a multifaceted autoimmune disorder with various associated comorbidities.

(Boniface, 2023) delves deeper into the aetiopathogenesis of vitiligo, exploring the cellular interactions that lead to melanocyte loss. This review highlights the role of immune responses in the development of vitiligo lesions and points to genetic determinants that contribute to disease susceptibility. The insights presented here are pivotal in understanding the shared pathogenic mechanisms between vitiligo and alopecia areata.

Lastly, (Sharma et al., 2024) discuss dietary influences on skin health, particularly in relation to alopecia areata. They emphasize the role of vitamin D in keratinocyte proliferation and its potential therapeutic implications for autoimmune conditions. Their findings suggest that dietary interventions may play a role in managing both alopecia areata and vitiligo, further linking these disorders through lifestyle factors.

Overall, the literature presents a compelling narrative that underscores the intricate relationships between vitiligo, alopecia areata, and other autoimmune conditions,



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

revealing a shared genetic and immunological landscape that warrants further exploration.

## LITERATURE REVIEW

The article "The Genetic Architecture of Alopecia Areata" by (Meredith Petukhova, 2013) presents a comprehensive examination of the multifaceted relationship between alopecia areata (AA) and various psychiatric disorders, particularly focusing on the emotional and psychological implications for affected individuals. The article systematically explores the increased prevalence of psychiatric conditions, such as depressive disorders, among patients with alopecia areata, emphasizing the need for a holistic approach to treatment that encompasses both dermatological and psychological care.

One of the key insights from the article is the association between alopecia areata and a history of atopy or autoimmunity, which may heighten the risk of developing this condition. (Meredith Petukhova, 2013) highlights that individuals with a background of atopic dermatitis or other autoimmune diseases may be predisposed to alopecia areata, suggesting a potential shared genetic or environmental pathway that warrants further investigation. This connection between autoimmune conditions and alopecia areata could also extend to vitiligo, as both disorders are characterized by immune-mediated processes that lead to the destruction of pigment-producing cells or hair follicles.

The article further delves into the psychiatric symptomatology observed in children with alopecia areata, noting that life events and emotional disturbances often exacerbate the condition. (Meredith Petukhova, 2013)'s evaluation of patients underscores the importance of recognizing the psychological burden of alopecia areata, which can lead to significant emotional distress and social challenges for those affected. This aspect of the condition is crucial, as it not only impacts the individual’s quality of life but also complicates the management of the disorder itself.



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Diphenylcyclopropenone-Induced Vitiligo in a Patient with Alopecia Universalis" by (Riad et al., 2013) presents a detailed examination of the interplay between vitiligo and alopecia areata, particularly in the context of diphenylcyclopropenone (DCP) treatment. The authors highlight the anatomical colocalization of these two skin disorders, noting a significant female-to-male ratio of 3:1 among the cases studied, with a notable prevalence in Asian populations, particularly from India.

One of the critical insights from the article is the temporal relationship between the onset of alopecia areata and vitiligo. The authors provide compelling evidence that in some patients, alopecia preceded the development of vitiligo, while in others, the reverse was true. This suggests a complex interaction between the two conditions, potentially influenced by underlying autoimmune mechanisms. The pathological examination confirming diagnoses in 7 of the 8 reported cases underscores the necessity for careful clinical evaluation in patients presenting with these overlapping symptoms.

The article further discusses the efficacy of DCP in inducing vitiligo in patients with severe forms of alopecia areata. It is noteworthy that 75% of cases exhibited vitiligo as focal lesions, either in remote areas or adjacent to the DCP application site. This observation raises questions about the localized immune response triggered by DCP and its implications for understanding the pathogenesis of both disorders.

Moreover, the authors reference a genome-wide association study that identifies several susceptibility loci associated with both vitiligo and alopecia areata, including three loci within the human leukocyte antigen region. This genetic overlap suggests a shared autoimmune pathway that may contribute to the development of both conditions. The mention of autoimmune polyendocrinopathy and Vogt-Koyanagi-Harada syndrome adds another layer of complexity, indicating that patients with these syndromes may also present with vitiligo and alopecia areata, further supporting the notion of a systemic autoimmune process.



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Vitiligo and autoimmune thyroid disorders" by (Baldini et al., 2017) provides a comprehensive examination of the intricate relationship between vitiligo and various autoimmune conditions, particularly focusing on the role of Th17-related cytokines and the implications of regulatory T cells in the pathogenesis of vitiligo vulgaris. The authors present a compelling argument regarding the dysregulation of melanocyte function, which is significantly influenced by Th17 cell infiltration, highlighting its relevance in the broader context of autoimmune disorders.

One of the critical insights from the article is the discussion on the expression and signaling of the tyrosine kinase FGFR2b/KGFR, which is shown to regulate phagocytosis and melanosome uptake in human keratinocytes. This finding is pivotal as it suggests a mechanistic pathway through which autoimmune processes may disrupt normal melanocyte function, leading to the characteristic depigmentation seen in vitiligo. The authors effectively connect this dysregulation to the increased prevalence of comorbid autoimmune diseases in patients with vitiligo, as evidenced by their cross-sectional studies and retrospective analyses.

Furthermore, the article delves into the notion of polyglandular autoimmune diseases, emphasizing that patients with generalized vitiligo often present with a higher incidence of other autoimmune conditions, including autoimmune thyroid disorders. This correlation is particularly relevant for clinicians, as it underscores the need for comprehensive evaluations of patients with vitiligo for potential coexisting autoimmune diseases.

The discussion on familial generalized vitiligo and its association with early disease onset and increased risk of other autoimmune diseases is another significant aspect of the paper. This highlights the genetic predisposition that may underlie these conditions, suggesting that individuals with a family history of autoimmune disorders should be monitored more closely for the development of vitiligo and related comorbidities.



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Videodermoscopic Changes of the Hair in Vitiligo Lesions in Relation to Disease Duration" by (Anbar et al., 2022) provides a comprehensive examination of the interrelationship between vitiligo and alopecia areata (AA), particularly through the lens of autoimmune processes and genetic predispositions. The authors articulate that vitiligo is characterized as an acquired chronic depigmenting autoimmune disease, which results in the loss of melanocytes and manifests as leukoderma and leukotrichia. This characterization underscores the autoimmune nature of the condition, which is pivotal in understanding its association with other autoimmune disorders, including AA.

The article highlights the increased prevalence of autoimmune diseases among individuals with vitiligo and their relatives, suggesting a potential genetic predisposition that may contribute to the co-occurrence of these conditions. This is particularly relevant when considering the shared genetic risk factors and pathogenesis that both vitiligo and AA exhibit. The authors note that colocalization of these disorders, especially on the scalp, has been documented, which is significant as it may indicate a common pathway or mechanism that predisposes individuals to both conditions.

Furthermore, the findings regarding decreased hair density in depigmented areas are particularly noteworthy. The authors present evidence from skin biopsies that reveal features indicative of both vitiligo and AA, reinforcing the notion that these conditions may not only coexist but may also influence each other’s clinical presentation. This suggests that clinicians should maintain a high index of suspicion for AA in patients presenting with vitiligo, particularly when there are observable changes in hair density or texture.

The article titled "Study of the Thyroid Profile of Patients with Alopecia" by (Popa et al., 2023) delves into the intricate relationships between alopecia areata (AA), vitiligo, and atopic dermatitis (AD), highlighting their classification as 'sister diseases.' The authors present compelling evidence that AA shares a common autoimmune background with other conditions, including atopic dermatitis, asthma,



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

and allergic rhinitis. This association is supported by a comprehensive study involving 61,899 individuals diagnosed with AA, which reinforces the prevalence of these comorbidities.

The article underscores the significance of assessing thyroid hormones and antithyroid antibodies in patients exhibiting specific manifestations of AA. This is particularly relevant given the autoimmune nature of these conditions. The authors suggest that the common autoimmune pathophysiology may underpin the observed associations, which could have implications for clinical practice. For instance, understanding the thyroid profile in patients with AA could lead to better management strategies and outcomes for those also presenting with vitiligo or atopic dermatitis.

Furthermore, the discussion on hyperandrogenemia-related alopecia in females introduces another layer to the understanding of hair loss disorders. The authors note that androgen excess can contribute to alopecia, suggesting that hormonal evaluations may be necessary in the diagnostic process. This aspect is particularly critical when considering treatment options, as interventions may need to address underlying hormonal imbalances.

Additionally, the role of vitamin D in skin metabolism and its potential link to alopecia is a noteworthy point raised in the article. The authors emphasize that hypovitaminosis D might manifest as alopecia, suggesting a need for further exploration into the relationship between vitamin D levels and autoimmune skin conditions. This highlights the importance of a holistic approach in managing patients with AA, vitiligo, and atopic dermatitis, where nutritional factors could play a pivotal role.

The article titled "Comorbidity Burden Among Patients with Vitiligo in the United States: A Large-Scale Retrospective Claims Database Analysis" by (Ezzedine et al., 2023) presents a comprehensive examination of the comorbidities associated with vitiligo, particularly focusing on autoimmune conditions such as atopic dermatitis



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

and alopecia areata. The study utilizes a large retrospective claims database to illustrate a higher incidence of both psychiatric and autoimmune diseases among patients diagnosed with vitiligo compared to a control cohort.

One of the critical insights from the analysis is the significantly elevated risk of developing autoimmune comorbidities, including atopic dermatitis and alopecia areata, in individuals with vitiligo. This finding is particularly relevant as it underscores the interconnectedness of autoimmune disorders, suggesting a shared pathogenesis. The authors highlight that patients with vitiligo who also have at least one comorbid autoimmune disease tend to exhibit more extensive vitiligo, measured by body surface area, than those without such comorbidities. This observation may indicate that the presence of additional autoimmune conditions exacerbates the severity of vitiligo, which warrants further investigation into the mechanisms behind this relationship.

Moreover, the article discusses the genetic predispositions that may contribute to the susceptibility to vitiligo and its associated comorbidities. The identification of shared genetic abnormalities among various autoimmune diseases supports the hypothesis of a common underlying pathophysiological process. This aspect of the study is crucial as it opens avenues for future research aimed at understanding the genetic and environmental factors that may contribute to the onset and progression of these autoimmune conditions.

The authors also provide statistical evidence of the increased incidence rates of conditions such as inflammatory bowel disease and autoimmune thyroid disease among patients with vitiligo, which further emphasizes the broader implications of vitiligo as a complex autoimmune disorder. The reported twofold and eightfold higher incidences, respectively, demonstrate the significant health burden faced by this population and the need for healthcare providers to be vigilant in monitoring for additional autoimmune diseases in patients with vitiligo.



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Aetiopathogenesis of Vitiligo" by (Boniface, 2023) provides a comprehensive exploration of the underlying mechanisms contributing to vitiligo, a chronic autoimmune condition characterized by skin depigmentation due to melanocyte loss. The narrative review emphasizes the pivotal role of resident T memory cells in both the development and recurrence of vitiligo lesions, thereby highlighting a critical component of the skin's immune response.

(Boniface, 2023) elucidates the complex interplay of various factors that contribute to the pathophysiology of vitiligo. The author identifies genetic predisposition, environmental triggers, intrinsic abnormalities of melanocytes, and oxidative stress as key elements that collectively lead to an exaggerated immune response. Notably, the article underscores the significance of genome-wide association studies (GWAS) in advancing the understanding of vitiligo's genetic architecture, revealing that genetic determinants account for approximately 80% of the overall risk for developing the condition. This assertion is supported by the identification of over fifty susceptibility genes linked to immune response pathways, which is crucial for understanding the disease's aetiology.

The article's critical evaluation of the immune mechanisms involved in vitiligo offers valuable insights into the relationship between this condition and other autoimmune diseases, such as alopecia areata and atopic dermatitis. The shared immunological pathways, particularly the involvement of T cells, suggest a potential commonality in the mechanisms driving these conditions. This intersection raises important questions regarding the co-occurrence of vitiligo with other autoimmune disorders, as both alopecia areata and atopic dermatitis involve similar immune dysregulations.

Moreover, (Boniface, 2023)'s review calls attention to the need for further research to elucidate the precise intercellular interactions that activate the local immune response in vitiligo. Understanding these interactions could pave the way for novel therapeutic strategies aimed at modulating the immune response, which may also



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

be applicable to treating associated conditions like alopecia areata and atopic dermatitis.

The article titled "Dietary Influences on Skin Health in Common Dermatological Disorders" by (Sharma et al., 2024) provides a comprehensive examination of the relationship between dietary factors and skin health, particularly focusing on autoimmune conditions like Vitiligo and Alopecia Areata. The authors emphasize the role of these conditions in affecting skin and hair health, highlighting Alopecia Areata (AA) as a prevalent autoimmune disorder that impacts approximately 2% of the population, presenting with diverse clinical manifestations.

One of the key insights from the article is the significant role of Vitamin D in the proliferation of keratinocytes, which are essential for skin health. The authors draw a connection between Vitamin D deficiency and the exacerbation of autoimmune disorders, suggesting that maintaining adequate levels of this vitamin could be crucial in managing conditions like AA. They also discuss the therapeutic potential of topical calcipotriol, a Vitamin D analog, noting its effectiveness as an adjunct treatment for AA. This finding underscores the importance of exploring vitamin-based therapies in the management of autoimmune skin conditions.

Furthermore, the article delves into dietary interventions, particularly the impact of anti-inflammatory diets. The authors advocate for gluten-free diets and those rich in antioxidants, positing that these dietary choices may help mitigate inflammation associated with AA. This perspective aligns with growing interest in the role of nutrition in managing chronic inflammatory conditions, although the authors acknowledge that more research is needed to establish definitive therapeutic protocols.

The mention of micronutrients such as zinc and iron is particularly noteworthy. The authors indicate that while these micronutrients are vital for the hair follicle cycle, their effectiveness in oral supplementation for AA remains inconclusive. This highlights a critical gap in current understanding and suggests the need for further



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

studies to elucidate the relationship between these micronutrients and hair health in autoimmune disorders.

## CONCLUSION

The literature concerning the association of vitiligo and alopecia areata with atopic dermatitis reveals a multifaceted interplay of genetic, immunological, and environmental factors. The studies reviewed indicate a significant overlap in the pathogenesis of these autoimmune conditions, suggesting that they may share common underlying mechanisms.

The foundational work by (Meredith Petukhova, 2013) establishes the prevalence of alopecia areata among individuals with psychiatric disorders, emphasizing the emotional impact of the condition. This study highlights the link between a history of atopy or autoimmunity and the development of alopecia areata, which is crucial for understanding its relationship with vitiligo.

In a complementary study, (Riad et al., 2013) documents cases of anatomical colocalization between alopecia areata and vitiligo, revealing shared susceptibility loci through genome-wide association studies. This genetic overlap supports the hypothesis of interrelated mechanisms of follicular melanocyte destruction in both conditions.

Further exploration by (Baldini et al., 2017) focuses on the role of Th17-related cytokines in vitiligo, linking dysregulation of melanocyte function to autoimmune pathways shared with alopecia areata. The study underscores the prevalence of comorbid autoimmune diseases in vitiligo patients, reinforcing the interconnected nature of these disorders.

The investigation of videodermoscopic changes in vitiligo by (Anbar et al., 2022) reveals similarities in pathogenesis with alopecia areata, suggesting common genetic risk factors and colocalization, particularly in scalp regions. This reinforces the notion that both diseases may arise from similar immunological disturbances.



Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The characterization of alopecia areata and vitiligo as “sister diseases” by (Popa et al., 2023) further elucidates their relationship, highlighting associations with atopic dermatitis and asthma. The study advocates for thyroid hormone assessments in alopecia areata patients, reflecting the intricate connections among various autoimmune conditions.

A large-scale retrospective analysis by (Ezzedine et al., 2023) reveals a higher incidence of autoimmune comorbidities, including alopecia areata, among patients with vitiligo, emphasizing the multifaceted nature of vitiligo as an autoimmune disorder. This study also points to significant correlations between vitiligo and psychological conditions, further complicating its clinical presentation.

The exploration of the aetiopathogenesis of vitiligo by (Boniface, 2023) highlights the role of immune responses and genetic determinants in disease susceptibility, offering insights into shared pathogenic mechanisms with alopecia areata.

Lastly, the discussion of dietary influences on skin health by (Sharma et al., 2024) suggests that nutritional interventions may play a role in managing both alopecia areata and vitiligo, linking these disorders through lifestyle factors.

In conclusion, the literature presents a compelling narrative that underscores the intricate relationships between vitiligo, alopecia areata, and atopic dermatitis. The shared genetic and immunological landscape highlighted in these studies warrants further exploration, particularly regarding the underlying mechanisms that contribute to the co-occurrence of these autoimmune conditions.

## DISCLOSURE STATEMENT


- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -




Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

## REFERENCES


1. Meredith Petukhova, L., 2013. The Genetic Architecture of Alopecia Areata. [PDF]
2. Riad, H., Al Mannai, H., Mansour, K., Al Qaatri, K., Al Dosari, S., Al Obaidaly, A., & Sultan, E., 2013. Diphenylcyclopropenone-Induced Vitiligo in a Patient with Alopecia Universalis. ncbi.nlm.nih.gov
3. Baldini, E., Odorisio, T., Sorrenti, S., Catania, A., Tartaglia, F., Carbotta, G., Pironi, D., Rendina, R., D'armiento, E., Persechino, S., & Ulisse, S., 2017. Vitiligo and autoimmune thyroid disorders. [PDF]
4. Anbar, T., Abdelraouf, H., Abd Elfattah Afify, A., Hussien Ragaie, M., Abulfotooh Eid, A., & Moneib, H., 2022. Videodermoscopic Changes of the Hair in Vitiligo Lesions in Relation to Disease Duration. ncbi.nlm.nih.gov
5. Popa, A., Carsote, M., Cretoiu, D., Cristian Dumitrascu, M., Nistor, C. E., & Sandru, F., 2023. Study of the Thyroid Profile of Patients with Alopecia. ncbi.nlm.nih.gov
6. Ezzedine, K., M. Soliman, A., Li, C., S. Camp, H., & G. Pandya, A., 2023. Comorbidity Burden Among Patients with Vitiligo in the United States: A Large-Scale Retrospective Claims Database Analysis. ncbi.nlm.nih.gov
7. Boniface, K., 2023. Aetiopathogenesis of Vitiligo. ncbi.nlm.nih.gov
8. Sharma, N., M Chaudhary, S., Khungar, N., K Aulakh, S., Idris, H., Singh, A., & Sharma, K., 2024. Dietary Influences on Skin Health in Common Dermatological Disorders. ncbi.nlm.nih.gov




Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved