Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Dermatology and Venereology



Chronic Spontaneous Urticaria

[1] Andi, [1] Aceng

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.andi9283727@gmail.com

Article History :

Received date : 2024/10/27
Revised date : 2024/15/12
Accepted date : 2024/12/05
Published date : 2025/01/07

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: Chronic Spontaneous Urticaria (CSU) is a prevalent and complex condition characterized by the recurrent appearance of itchy wheals and angioedema lasting for more than six weeks without an identifiable trigger. The literature on CSU has evolved significantly over the years, providing deeper insights into its pathophysiology, management strategies, and the impact on patients' quality of life. **Literature Review:** Despite the availability of various treatment options, including antihistamines and the monoclonal antibody omalizumab, many patients continue to experience inadequate symptom control (Lin et al., 2022). The literature suggests that up to 50% of patients may not achieve adequate symptom relief even with increased dosages of antihistamines (Lin et al., 2022). This gap in effective management underscores the need for ongoing research to refine treatment protocols and explore novel therapeutic targets (Kocatürk et al., 2017). The exploration of biomarkers in CSU is a promising area of research, as identifying reliable markers could enhance the understanding of disease severity and therapeutic responses (Puxeddu et al., 2019). The complexity of CSU, with its multifaceted nature involving immune dysregulation, varying treatment responses, and significant impacts on quality of life, calls for a comprehensive approach to management and ongoing research to improve patient outcomes. The psychological burden associated with CSU is also noteworthy, as high levels of anxiety and psychiatric comorbidities are common among affected individuals (Sussman et al., 2015). This highlights the necessity for a multidisciplinary approach to treatment that addresses both the physical symptoms and the psychological impact on patients. **Conclusion:** In conclusion, the literature on Chronic Spontaneous Urticaria reveals a complex interplay of autoimmune mechanisms, psychological factors, and treatment challenges. While advancements in management strategies, such as the use of omalizumab, have shown promise, the persistent inadequacy of symptom control for many patients emphasizes the need for further research and innovative therapeutic approaches. The ongoing exploration of biomarkers and the understanding of the underlying pathophysiology will be crucial for developing more effective treatments and improving the overall quality of life for individuals suffering from this debilitating condition.

Keyword: Chronic Spontaneous Urticaria

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

Chronic Spontaneous Urticaria (CSU) is a prevalent and complex condition characterized by the recurrent appearance of itchy wheals and angioedema lasting for more than six weeks without an identifiable trigger. The literature on CSU has evolved significantly over the years, providing deeper insights into its pathophysiology, management strategies, and the impact on patients' quality of life.

In 2015, (Sussman et al., 2015) highlighted that CSU predominantly affects women aged 20 to 40, with a notable autoimmune component present in about 45% of cases. The study emphasized the psychological burden associated with CSU, linking high anxiety levels and psychiatric comorbidities to diminished quality of life. (Ferrer, 2015) further elaborated on the management of CSU, underscoring the challenges posed by the condition's persistence despite treatment, and identifying an autoimmune origin in a subset of patients. They also noted the emergence of omalizumab as a promising therapeutic option for patients unresponsive to antihistamines.

(Ferrer, 2015) contributed to the understanding of the immunological events in CSU, revealing that a significant percentage of patients exhibit unique mast cell and basophil characteristics, which may play a role in the disease's pathogenesis. The complexity of these immune responses highlights the need for further research to establish a comprehensive model explaining the interactions between various immune cells in CSU.

By 2017, (Kocatürk et al., 2017) acknowledged the substantial impact of CSU on patients' daily lives, emphasizing the necessity for improved treatment options. They pointed out that while omalizumab has enhanced management strategies, many patients continue to struggle with existing therapies, indicating a gap in effective treatments for this population.

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The exploration of autoimmune theories by (J. Bracken et al., 2019) reinforced the notion that CSU may be an autoimmune condition for many patients. They discussed the potential mechanisms involving mast cell and basophil activation, suggesting that dysregulation in these cells is central to CSU's pathophysiology. This aligns with findings from (Puxeddu et al., 2019) who identified mediators as potential biomarkers for CSU, further indicating the involvement of autoimmunity in the disease.

(Lin et al., 2022) conducted a network meta-analysis examining the effectiveness and safety of various treatments for CSU, noting that a significant proportion of patients remain inadequately controlled even with advanced therapies. The findings highlighted the importance of ongoing research to refine treatment protocols and improve patient outcomes.

(Khan et al., 2022) provided a narrative review of treatment options for resistant CSU, emphasizing the necessity of a thorough clinical evaluation to identify possible autoimmune or infectious etiologies. They noted that high doses of antihistamines are often required, and in cases resistant to standard treatment, clinicians may need to explore established immunomodulatory agents.

Overall, the literature underscores the complexity of CSU, revealing its multifaceted nature involving autoimmune mechanisms, varying responses to treatment, and significant impacts on patients' quality of life. The ongoing research and evolving management strategies reflect a growing understanding of this debilitating condition, paving the way for future advancements in treatment and care.

LITERATURE REVIEW

The article "Insights and advances in chronic urticaria: a Canadian perspective" by (Sussman et al., 2015) provides a comprehensive overview of chronic spontaneous urticaria (CSU), highlighting its epidemiology, pathophysiology, and implications for patient management. The authors emphasize that CSU predominantly affects

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

women, with a notable peak incidence occurring between the ages of 20 and 40 years. This demographic insight is critical for understanding the population most at risk and for tailoring preventive strategies.

One of the significant findings presented is the prevalence of angioedema, which affects a substantial proportion of CSU patients, ranging from 33% to 66%. This association underscores the need for clinicians to be vigilant in diagnosing and managing both conditions concurrently, as angioedema can lead to more severe outcomes and increased patient distress.

The article also addresses the autoimmune aspect of CSU, noting that approximately 45% of patients exhibit an autoimmune response, characterized by antibodies targeting the IgE receptor alpha subunit or IgE itself on mast cells. This finding is pivotal as it points to a potential therapeutic avenue for patients with an autoimmune component, suggesting that treatments aimed at modulating the immune response may be beneficial.

Furthermore, the authors highlight the psychological impact of CSU, noting high levels of anxiety and psychiatric co-morbidities among affected individuals. This aspect is often overlooked in dermatological conditions, yet it is crucial for understanding the overall burden of the disease on quality of life. The recognition that CSU can lead to significant psychological distress necessitates a multidisciplinary approach to treatment that includes mental health support.

The article also discusses diagnostic strategies, emphasizing the importance of a thorough history and physical examination. The authors specifically mention the need to inquire about non-steroidal anti-inflammatory drug (NSAID) use, as a notable percentage of patients report exacerbations linked to these medications. This insight is essential for clinicians, as it can influence management decisions and help mitigate flare-ups.

Lastly, the article notes the chronic nature of CSU, with many patients experiencing symptoms for over five years, particularly those with more severe manifestations.

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

This long-term perspective is critical for healthcare providers to set realistic expectations for patients and to develop long-term management plans.

The article "Management of urticaria: not too complicated, not too simple" by provides a comprehensive overview of Chronic Spontaneous Urticaria (CSU), a condition that significantly impacts the quality of life for many patients. Defined as urticaria that persists for more than six weeks, CSU is characterized by the spontaneous appearance of wheals and, sometimes, angioedema. The authors highlight the complexity of managing this condition, noting that despite the availability of various treatment options, patients often experience persistent symptoms.

A critical aspect of the article is its exploration of the underlying mechanisms of CSU, which remain poorly understood. The authors point out that while an autoimmune origin has been identified in a subset of patients, distinguishing these individuals from those with non-autoimmune CSU is challenging. This lack of clarity complicates treatment strategies, as the clinical presentation does not differ significantly between the two groups. The authors emphasize the need for further research to elucidate the pathophysiology of CSU, which is essential for developing targeted therapies.

The article also discusses treatment options, particularly the role of omalizumab, a monoclonal antibody that has shown efficacy in controlling CSU symptoms in patients who do not respond to high doses of antihistamines or combination therapies. Although omalizumab has emerged as a promising treatment, the authors note that its mechanism of action is not fully understood, raising questions about its long-term use and potential side effects. This highlights an important gap in the current understanding of CSU management.

The article "Immunological events in chronic spontaneous urticaria" by provides a comprehensive overview of the immunological mechanisms implicated in chronic spontaneous urticaria (CSU), a condition characterized by the recurrent appearance

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

of pruritic wheals and angioedema lasting longer than six weeks. The author highlights the debilitating nature of CSU and its systemic features, emphasizing the need for a deeper understanding of its underlying physiopathology, which remains largely unknown.

Ferrer presents compelling evidence supporting an autoimmune basis for CSU, particularly through the observation that serum from CSU patients can activate mast cells and basophils from healthy donors. This finding suggests that the immune response in CSU may be driven by autoantibodies or other serum factors that provoke an inappropriate activation of these cells, which are crucial mediators in allergic responses. The role of mast cells and basophils is further elaborated, with the article noting that basophils from CSU patients exhibit a unique hyporesponsiveness to IgE receptor stimuli while becoming hyperresponsive when exposed to patient sera. This paradoxical behavior indicates a complex dysregulation of the immune response in CSU, which could contribute to the chronic inflammation characteristic of the condition.

Additionally, the presence of eosinophils in skin biopsies from CSU patients is acknowledged, although their specific contributions to the pathophysiology of CSU remain unclear. The article suggests that these cells may play a role in the inflammatory milieu, yet further research is needed to elucidate their function within the context of CSU.

Ferrer also discusses the involvement of endothelial cells, noting an increase in vasoactive peptides in the skin and plasma of CSU patients. This observation points to a systemic inflammatory response that may exacerbate the symptoms of CSU, as these peptides can influence vascular permeability and contribute to the formation of wheals and angioedema.

The article "Looking forward to new targeted treatments for chronic spontaneous urticaria" by (Kocatürk et al., 2017) provides a comprehensive overview of chronic spontaneous urticaria (CSU), characterized by the spontaneous appearance of itchy

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

wheals, angioedema, or both for a duration of at least six weeks. The authors emphasize that while CSU is generally a self-limiting disorder, with a typical duration of 2–5 years, a significant proportion of patients (20%) endure symptoms for over five years. This chronicity can lead to a substantial reduction in quality of life, affecting various aspects such as sleep, daily activities, social interactions, work productivity, and emotional well-being.

The economic implications of CSU are also highlighted, with the authors noting both direct costs, such as medication and healthcare visits, and indirect costs associated with reduced work efficiency and absenteeism. This dual burden underscores the need for effective management strategies to alleviate the symptoms and improve the overall well-being of affected individuals.

A pivotal advancement in the treatment landscape for CSU is the introduction of omalizumab, which has been shown to significantly enhance therapeutic outcomes when used as an add-on therapy to H1 antihistamines. However, the article critically notes that a subset of patients does not respond adequately to omalizumab or cannot tolerate it, indicating a gap in the current treatment paradigm.

In addressing future directions for therapy, the authors discuss potential novel therapeutic targets and treatment options that could emerge as viable alternatives for those who do not benefit from existing therapies. This forward-looking perspective is crucial, as it not only highlights the ongoing challenges in managing CSU but also points to the necessity for continued research and innovation in this field.

The article "Autoimmune Theories of Chronic Spontaneous Urticaria" by (J. Bracken et al., 2019) provides a comprehensive overview of chronic spontaneous urticaria (CSU), emphasizing its classification, potential autoimmune underpinnings, and the complexities of its pathophysiology. CSU is characterized by recurrent hives lasting longer than six weeks without identifiable triggers, and

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

the authors highlight recent guidelines that expand this definition to include isolated idiopathic angioedema, provided that other causes are excluded.

The article critically examines the notion that a significant portion of CSU cases may be autoimmune in nature. This assertion is supported by multiple studies, yet the authors underscore the necessity of identifying possible triggers and ruling out differential diagnoses before arriving at a definitive diagnosis. This caution is vital, as misdiagnosis could lead to inappropriate management strategies.

A key focus of the article is the pathophysiological mechanisms underlying CSU. The authors detail the central role of mast cells and basophils as primary effectors in chronic urticaria. The discussion of histamine and other mast cell-derived products as pivotal in the development of CSU is particularly noteworthy. The authors propose two significant mechanisms contributing to the pathogenesis of chronic urticaria. The first mechanism involves the dysregulation of intracellular signaling pathways within mast cells and basophils, which can lead to defects in their trafficking or functional capabilities. The second mechanism posits the formation of autoantibodies targeting FcεRIα or IgE on mast cells and basophils, suggesting that the activation of the high-affinity IgE receptor, FcεR1, is crucial in the manifestation of urticaria.

The article "Autoimmune Theories of Chronic Spontaneous Urticaria" by (J. Bracken et al., 2019) provides a comprehensive overview of chronic spontaneous urticaria (CSU), a condition characterized by recurrent hives lasting longer than six weeks without an identifiable trigger. The authors emphasize the significant prevalence of urticaria, affecting 15 to 25% of individuals at some point in their lives, with a noted increase in occurrence among adults and particularly women in their third to fifth decades of life.

A critical aspect of the article is its classification of urticaria into acute and chronic forms, with CSU being defined by its persistent nature. The authors highlight the estimated prevalence of CSU, which ranges from 0.5 to 5% in the general

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

population, although they acknowledge the challenges in accurately determining these figures. This acknowledgment of the limitations in prevalence data is an important aspect of the article, as it underscores the need for further research to understand the true incidence of CSU.

The article also addresses the autoimmune theories surrounding CSU, positing that a substantial proportion of cases may be linked to autoimmune mechanisms. This perspective is crucial as it shifts the focus towards understanding the underlying pathophysiology of CSU, which could lead to more targeted therapeutic approaches. The authors suggest that while autoimmune factors may play a role, it is vital to identify potential triggers and rule out other differential diagnoses before concluding a diagnosis of CSU. This cautious approach is commendable, as it encourages thorough clinical evaluation and consideration of a broad differential diagnosis in affected patients.

Furthermore, the inclusion of isolated idiopathic angioedema within the definition of CSU as per recent guidelines reflects an evolving understanding of the condition. This change may enhance the clinical management of patients who present with symptoms of angioedema alongside urticaria, thereby improving patient outcomes.

The article "Biomarkers In Chronic Spontaneous Urticaria: Current Targets And Clinical Implications" by (Puxeddu et al., 2019) provides a comprehensive examination of chronic urticaria (CU), particularly focusing on chronic spontaneous urticaria (CSU). The authors highlight the negative impact of CU on patients' quality of life, emphasizing the importance of understanding its underlying mechanisms to improve management strategies.

The article delineates CU as a condition characterized by recurrent itchy wheals and/or angioedema lasting more than six weeks, which is further classified into CSU and inducible urticaria. A significant contribution of this review is the discussion on the role of skin mast cells as the primary effector cells in CSU's pathogenesis. However, the authors note that the specific triggers for mast cell

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

activation remain largely unidentified, suggesting a gap in current understanding that warrants further research.

The investigation into autoimmunity as a contributing factor to CSU is particularly noteworthy. The authors present evidence of circulating IgG autoantibodies directed towards thyroid antigens and IgE or its high-affinity receptor (FcεRI). This association implies an autoimmune component in some patients with CSU. However, the authors also point out that the absence of detectable autoantibodies in a significant proportion of CSU cases indicates the potential involvement of alternative mechanisms in the disease's pathogenesis.

Moreover, the article emphasizes the need for identifying reliable biomarkers that could aid in assessing disease severity and predicting therapeutic responses. This focus on biomarkers is crucial, as it could lead to more personalized treatment approaches for patients suffering from CSU. The authors suggest that ongoing research in this area may illuminate the complex interplay of factors contributing to CSU and lead to improved clinical outcomes.

The article "Effectiveness and Safety of Immunosuppressants and Biological Therapy for Chronic Spontaneous Urticaria: A Network Meta-Analysis" by (Lin et al., 2022) provides a comprehensive evaluation of treatment options for Chronic Spontaneous Urticaria (CSU), a condition characterized by persistent hives, pruritus, and angioedema lasting longer than six weeks without identifiable external triggers. The authors highlight that CSU is prevalent, affecting approximately 1% of the population, with a notable peak incidence in individuals aged 20 to 40 years and a higher prevalence in females.

The article effectively underscores the significant impact of CSU on patients' quality of life, noting that individuals with this condition often experience diminished daily functioning and social interactions. Furthermore, the association between chronic urticaria and increased psychiatric disorders is a crucial point, as

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

it emphasizes the need for comprehensive management that addresses both physical and psychological aspects of the disease.

The authors present a systematic review and network meta-analysis of randomized controlled trials, which is a robust methodological approach to evaluate the effectiveness and safety of various treatment modalities for CSU. They delineate the current first-line treatment recommendations, which advocate for non-sedative antihistamine medications at standard doses. However, the article reveals a concerning statistic: up to 50% of patients may not achieve adequate symptom control even when receiving higher doses (3-4 times the recommended amount). This finding highlights a significant gap in effective management strategies for a substantial portion of the CSU population.

Moreover, the discussion of advanced treatment options, such as omalizumab and cyclosporin, is particularly relevant, as it provides insight into alternative therapies for patients who do not respond adequately to first-line treatments. The network meta-analysis presented in the article offers valuable data on symptom improvement and safety outcomes, which can guide clinicians in making informed treatment decisions.

The article titled "Resistant Chronic Spontaneous Urticaria – A Case Series Narrative Review of Treatment Options" by (Khan et al., 2022) provides a comprehensive overview of chronic spontaneous urticaria (CSU), emphasizing its diagnostic challenges and treatment options. The authors effectively delineate CSU as a condition characterized by recurrent, itchy wheals that persist for more than six weeks without identifiable triggers. This definition is crucial for clinicians, as it sets the stage for understanding the complexities involved in diagnosing and managing the condition.

One of the key insights presented in the article is the importance of a detailed clinical history and physical examination in diagnosing CSU, supported by specific investigations that may indicate underlying autoimmune or infectious etiologies.

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The authors highlight the utility of the Urticaria Activity Score (UAS), which quantifies disease severity through a scoring system that assesses both the presence of wheals and pruritus. This structured approach to monitoring disease activity is essential for tailoring treatment strategies effectively.

The article underscores the inadequacy of standard doses of antihistamines for many patients with CSU, advocating for doses up to four times the standard amount in cases of resistance. This recommendation is particularly relevant given the prevalence of treatment-resistant CSU, which poses significant challenges for both patients and healthcare providers. The authors also discuss alternative therapies, including immunomodulatory agents like dapsone and ciclosporin, for patients who do not respond to high-dose antihistamines and omalizumab. While dapsone shows promise in a subset of patients, the lack of identifiable clinical or laboratory markers for responsiveness presents a limitation that warrants further investigation.

Moreover, the article introduces the concept of neutrophilic urticaria as a distinct entity, characterized by more painful and burning lesions rather than the typical pruritic wheals associated with conventional urticaria. This differentiation is critical, as it suggests a different pathogenic mechanism potentially involving the inflammasome pathway. The authors' discussion of the pathophysiology of neutrophilic urticaria adds depth to the understanding of CSU and highlights the need for tailored therapeutic approaches based on underlying mechanisms.

CONCLUSION

Chronic Spontaneous Urticaria (CSU) is a condition that significantly impacts patients' quality of life due to its chronic nature, characterized by recurrent itchy wheals and angioedema lasting longer than six weeks without an identifiable trigger. The literature indicates that CSU predominantly affects women aged 20 to 40, with a substantial autoimmune component present in approximately 45% of cases (Sussman et al., 2015). This autoimmune aspect is echoed by several studies, which suggest that the activation of mast cells and basophils, potentially driven by

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

autoantibodies, plays a central role in the pathophysiology of CSU [ref: b5e3348d-0d50-491f-a516-3fcafdda1ab; ref: d36161c1-eb9b-459d-ba41-406ed44861ec].

The psychological burden associated with CSU is also noteworthy, as high levels of anxiety and psychiatric comorbidities are common among affected individuals (Sussman et al., 2015). This highlights the necessity for a multidisciplinary approach to treatment that addresses both the physical symptoms and the psychological impact on patients.

Despite the availability of various treatment options, including antihistamines and the monoclonal antibody omalizumab, many patients continue to experience inadequate symptom control (Lin et al., 2022). The literature suggests that up to 50% of patients may not achieve adequate symptom relief even with increased dosages of antihistamines (Lin et al., 2022). This gap in effective management underscores the need for ongoing research to refine treatment protocols and explore novel therapeutic targets (Kocatürk et al., 2017).

The exploration of biomarkers in CSU is a promising area of research, as identifying reliable markers could enhance the understanding of disease severity and therapeutic responses (Puxeddu et al., 2019). The complexity of CSU, with its multifaceted nature involving immune dysregulation, varying treatment responses, and significant impacts on quality of life, calls for a comprehensive approach to management and ongoing research to improve patient outcomes.

In conclusion, the literature on Chronic Spontaneous Urticaria reveals a complex interplay of autoimmune mechanisms, psychological factors, and treatment challenges. While advancements in management strategies, such as the use of omalizumab, have shown promise, the persistent inadequacy of symptom control for many patients emphasizes the need for further research and innovative therapeutic approaches. The ongoing exploration of biomarkers and the understanding of the underlying pathophysiology will be crucial for developing

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

more effective treatments and improving the overall quality of life for individuals suffering from this debilitating condition.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Sussman, G., Hébert, J., Gulliver, W., Lynde, C., Waserman, S., Kanani, A., Ben-Shoshan, M., Horemans, S., Barron, C., Betschel, S., H Yang, W., Dutz, J., Shear, N., Lacuesta, G., Vadas, P., Kobayashi, K., Lima, H., & Estelle R Simons, F., 2015. Insights and advances in chronic urticaria: a Canadian perspective. ncbi.nlm.nih.gov
2. Ferrer, M., 2015. Immunological events in chronic spontaneous urticaria. ncbi.nlm.nih.gov
3. Kocatürk, E., Maurer, M., Metz, M., & Grattan, C., 2017. Looking forward to new targeted treatments for chronic spontaneous urticaria. ncbi.nlm.nih.gov
4. J. Bracken, S., Abraham, S., Abraham, S., S. MacLeod, A., & S. MacLeod, A., 2019. Autoimmune Theories of Chronic Spontaneous Urticaria. [PDF]
5. Puxeddu, I., Petrelli, F., Angelotti, F., Croia, C., & Migliorini, P., 2019. Biomarkers In Chronic Spontaneous Urticaria: Current Targets And Clinical Implications. ncbi.nlm.nih.gov
6. Lin, W. K., Lin, S. J., Lee, W. R., Lin, C. C., Lin, W. C., Chang, H. C., Cheng, C. T., & C. Hsu, J., 2022. Effectiveness and Safety of

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Immunosuppressants and Biological Therapy for Chronic Spontaneous Urticaria: A Network Meta-Analysis. ncbi.nlm.nih.gov

7. Khan, S., Chopra, C., Mitchell, A., Nakonechna, A., Yong, P., & Yousuf Karim, M., 2022. Resistant Chronic Spontaneous Urticaria – A Case Series Narrative Review of Treatment Options. ncbi.nlm.nih.gov
8. J. Bracken, S., Abraham, S., & S. MacLeod, A., 2019. Autoimmune Theories of Chronic Spontaneous Urticaria. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved