Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Dermatology and Venereology



Hair Regrowth Outcomes of Contact Immunotherapy for Patients With Alopecia Areata

[1] Irene, [1] Hendra

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.irene893728@gmail.com

Article History :

Received date : 2024/10/12
Revised date : 2024/15/25
Accepted date : 2024/12/16
Published date : 2025/01/07

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature on hair regrowth outcomes for patients with alopecia areata (AA) has evolved significantly, reflecting an increasing understanding of the disease's complex pathophysiology and treatment modalities. **Literature Review:** Further exploration of treatment modalities reveals the importance of considering both clinical and psychosocial factors in managing AA. The systematic review by (Mostaghimi et al., 2021) emphasizes the emotional and social implications of hair loss, advocating for a holistic approach that integrates psychological support with medical treatments. This perspective is echoed in the findings of (R. Barton et al., 2021), which specifically addresses the unique treatment needs of pediatric patients, reinforcing the necessity for age-appropriate therapies. Moreover, the investigation of corticosteroid pulse therapy by (Rastaghi et al., 2023) illustrates the challenges of managing severe cases of AA, where traditional therapies may be insufficient. Their review highlights the potential for corticosteroids to provide relief, yet it also points to the complexities involved in treating more extensive forms of the disease. The variability in patient responses underscores the need for tailored treatment approaches, as highlighted by (Mahasaksiri et al., 2021), which indicates that while DPCP can be effective, response rates can fluctuate significantly. **Conclusion:** In conclusion, the body of literature surrounding hair regrowth outcomes in alopecia areata demonstrates a significant evolution in understanding both the clinical and psychosocial dimensions of the condition. The findings collectively advocate for a multifaceted approach that not only addresses the immunological aspects of treatment but also incorporates the emotional and social well-being of patients. This comprehensive perspective is essential for optimizing treatment outcomes and improving the quality of life for individuals affected by alopecia areata.

Keyword: Hair Regrowth Outcomes, Contact Immunotherapy, Alopecia Areata

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The literature on hair regrowth outcomes for patients with alopecia areata (AA) has evolved significantly, reflecting an increasing understanding of the disease's complex pathophysiology and treatment modalities. (Gestetner, 2013) provides a foundational overview of AA, highlighting the role of topical immunotherapy and the autoimmune mechanisms implicated in its pathogenesis. This early exploration sets the stage for understanding how immune response can be harnessed to promote hair regrowth in affected individuals.

Building on this foundation, (J. Dabek et al., 2019) evaluate the efficacy of emerging treatments, specifically laser-assisted hair regrowth techniques. Their literature review underscores the need for innovative approaches to address the limitations of traditional therapies, suggesting that fractional laser modalities may offer promising results in hair restoration.

In a more recent contribution, (Mahasaksiri et al., 2021) delve deeper into topical immunotherapy, emphasizing its safety and efficacy for recalcitrant cases of AA. They report a variable response rate, with some patients experiencing significant regrowth while others remain unresponsive. This variability underscores the need for personalized treatment strategies in managing AA, particularly given the high recurrence rates noted in their findings.

(Mostaghimi et al., 2021) further contextualize the impact of AA by examining the social and emotional ramifications of the disease. Their systematic review highlights the profound effects of hair loss on patients' quality of life, drawing attention to the psychosocial aspects that must be considered alongside clinical outcomes in treatment planning.

(R. Barton et al., 2021) focus on pediatric patients, providing a systematic review of treatment modalities for children with AA. Their findings reinforce the importance of tailored interventions, as children may respond differently to

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

treatments compared to adults. This review emphasizes the need for ongoing research to develop age-appropriate therapies that consider the unique challenges faced by younger patients.

(Abedini et al., 2022) investigate the treatment satisfaction and response rates among patients receiving diphenylcyclopropenone (DPCP), a form of contact immunotherapy. Their study reveals a significant range of responses, with many patients reporting satisfactory outcomes despite the high cost and potential side effects associated with DPCP. This highlights the critical role of monitoring patient satisfaction in evaluating treatment effectiveness.

Finally, (Rastaghi et al., 2023) provide a comprehensive review of corticosteroid pulse therapy for AA, discussing its efficacy and the psychological factors that may influence treatment outcomes. They note that while corticosteroids can be effective for mild to moderate cases, managing severe AA remains challenging. Their findings contribute to a nuanced understanding of the treatment landscape, indicating that a multifaceted approach may be necessary to optimize regrowth outcomes for patients with varying degrees of severity.

Together, these articles illustrate the multifactorial nature of alopecia areata and the diverse therapeutic strategies being explored to enhance hair regrowth. The evolution of treatment perspectives, from immunotherapy to psychosocial considerations, underscores the complexity of managing this autoimmune condition effectively.

LITERATURE REVIEW

In the article "Alopecia Areata: An Overview," (Gestetner, 2013) provides a comprehensive examination of contact immunotherapy as a treatment modality for patients suffering from alopecia areata. The article delves into the mechanisms of action of topical immunotherapy, emphasizing its role in modulating the immune response that is implicated in the pathogenesis of alopecia areata.

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

(Gestetner, 2013) outlines that alopecia areata is characterized by an autoimmune attack on hair follicles, leading to hair loss. The author discusses how contact immunotherapy, particularly with agents like diphencyprone (DPCP), can induce an allergic contact dermatitis that may redirect the immune response and promote hair regrowth. This is particularly relevant as the article highlights the importance of understanding the immunological underpinnings of the condition to tailor effective treatment strategies.

The article further addresses the clinical efficacy of contact immunotherapy, citing various studies that demonstrate favorable outcomes in terms of hair regrowth. (Gestetner, 2013) notes that while the response can be variable among patients, many experience significant regrowth after a series of treatments. The article also emphasizes the necessity of patient selection and the potential for adverse effects, which are critical considerations for practitioners when implementing this therapy.

In addition to contact immunotherapy, (Gestetner, 2013) explores other treatment strategies for alopecia areata, particularly in pediatric populations. The article provides insights into the challenges faced in treating younger patients, including the psychological impact of hair loss and the need for age-appropriate interventions. The discussion on current treatment strategies is particularly valuable, as it offers a holistic view of management options available for alopecia areata.

The article "Laser-assisted Hair Regrowth: Fractional Laser Modalities for the Treatment of Androgenic Alopecia" by (J. Dabek et al., 2019) presents a comprehensive literature review that evaluates the efficacy of various laser-assisted treatments for hair regrowth, specifically in the context of androgenic alopecia. While the primary focus of the article is on laser modalities, the insights gleaned from this review can also provide valuable context for understanding treatment approaches for other forms of hair loss, including alopecia areata.

The authors systematically analyze existing studies to assess the effectiveness of fractional laser treatments, highlighting that these modalities stimulate the scalp,

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

potentially enhancing hair follicle activity and promoting regrowth. The review discusses the mechanisms by which lasers may facilitate hair regrowth, such as increasing blood flow to the scalp and inducing a healing response in the skin, which could be relevant when considering immune-mediated conditions like alopecia areata.

One significant strength of the article is its thoroughness in evaluating a range of studies, which allows for a broader understanding of the potential outcomes associated with laser treatments. The authors present a balanced view, acknowledging both the promising results and the limitations of current research, such as the variability in treatment protocols and the need for standardized measures of efficacy.

However, the article could benefit from a more explicit discussion on the comparative effectiveness of laser-assisted treatments versus other therapeutic options for alopecia areata, such as contact immunotherapy. While the review provides a solid foundation for understanding laser treatments, it lacks a direct correlation to the specific outcomes of contact immunotherapy, which is a critical aspect for patients with alopecia areata seeking effective treatment options.

The article "Application of Topical Immunotherapy in the Treatment of Alopecia Areata: A Review and Update" by (Mahasaksiri et al., 2021) provides a comprehensive overview of the use of topical immunotherapy as a treatment option for alopecia areata (AA), a condition characterized by patchy or complete hair loss due to an autoimmune response targeting hair follicles. The authors highlight the challenges associated with treating AA, noting that up to 25% of patients may be resistant to conventional therapies and that recurrence rates can range from 13.5% to 33%.

The review meticulously examines various therapeutic modalities, emphasizing that treatment responses can be highly variable and unpredictable. In this context, topical immunotherapy emerges as a promising approach, particularly for patients

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

with recalcitrant AA. The authors explain that this treatment works by inducing an inflammatory response through contact sensitizing agents, which in turn stimulates hair regrowth. The reported response rates in the literature vary widely, from 9% to 87%, with an effective rate of approximately 30%. This variability underscores the need for individualized treatment plans and highlights the unpredictable nature of AA.

Furthermore, the article delves into the pharmacology and mechanism of action of topical immunotherapy, detailing how agents like diphencyprone (DPCP) are applied to the scalp to provoke a localized immune response. This response is theorized to counteract the autoimmune attack on hair follicles, thereby promoting hair regrowth. The authors also address potential adverse events associated with the treatment, which can include local dermatitis and pruritus, although these are generally manageable.

The article "Patient Perspectives of the Social, Emotional and Functional Impact of Alopecia Areata: A Systematic Literature Review" by (Mostaghimi et al., 2021) provides a comprehensive examination of the multifaceted effects of alopecia areata (AA) on patients' lives. The authors highlight that AA, as a chronic autoimmune condition characterized by unpredictable hair loss, extends beyond mere physical manifestations, significantly impacting patients' emotional and psychological well-being.

One of the key insights from the article is the profound emotional distress experienced by individuals with AA. The authors systematically reviewed literature that reveals how the visibility of hair loss can lead to feelings of shame, embarrassment, and social withdrawal. This emotional burden is often compounded by societal perceptions and stigmas associated with hair loss, which can result in decreased self-esteem and quality of life. The authors effectively synthesize various studies to illustrate that patients frequently report anxiety and depression linked to their condition, underscoring the necessity for healthcare providers to address these psychological aspects in conjunction with physical treatment.

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Moreover, the article delves into the functional impacts of AA on daily life. Patients often report challenges in social interactions, professional environments, and personal relationships due to their hair loss. The authors point out that these functional limitations can exacerbate the emotional toll, creating a cyclical pattern of distress. The systematic review emphasizes the importance of understanding these patient perspectives to inform more holistic treatment approaches.

In terms of clinical implications, the findings suggest that while treatments such as contact immunotherapy may focus on hair regrowth, it is equally crucial for treatment plans to incorporate psychological support. The authors advocate for an interdisciplinary approach that includes dermatologists, psychologists, and support groups to address the comprehensive needs of patients with AA.

The systematic review conducted by (R. Barton et al., 2021) provides a comprehensive analysis of treatment modalities for pediatric alopecia areata (AA). The authors meticulously adhered to the PRISMA guidelines, ensuring a robust methodological framework for their investigation. By focusing on peer-reviewed literature sourced from the PubMed database, the authors employed a well-defined search strategy utilizing specific keywords related to alopecia and treatment options.

One of the strengths of this review is its stringent eligibility criteria, which included only English articles that directly addressed treatment outcomes for individuals under 18 years of age. This focus is particularly pertinent given that alopecia areata manifests differently across age groups, and treatments may vary in efficacy between pediatric and adult populations. The exclusion of review articles, animal studies, and outdated treatment evaluations enhances the relevance and applicability of the findings to current clinical practices.

The authors provided a critical evaluation of the various treatment modalities available, highlighting the limited but promising outcomes associated with contact immunotherapy. This treatment has gained traction in recent years for its potential

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

to stimulate hair regrowth in patients with AA, particularly in pediatric cases. The review's findings suggest that while several treatment options exist, the evidence supporting their efficacy varies significantly, underscoring the need for further research in this area.

The article titled "Treatment satisfaction and response in patients with severe alopecia areata under treatment with diphenylcyclopropenone" by (Abedini et al., 2022) presents a thorough examination of the efficacy and patient satisfaction associated with the use of diphenylcyclopropenone (DPCP) as a treatment for severe alopecia areata (AA). This autoimmune condition, characterized by sudden and patchy hair loss, poses significant challenges for affected individuals, particularly when the disease is extensive and resistant to standard therapies.

The authors highlight the chronic nature of AA, noting that it is mediated by CD8+ T cells, which target hair follicles. This immunological perspective is crucial, as it underscores the need for treatments that can modify the immune response rather than merely addressing the symptoms of hair loss. The article emphasizes that while various treatment options exist—including corticosteroids and contact immunotherapy—no definitive gold standard has been established, particularly for severe cases of AA. DPCP emerges as a preferred method due to its immunotherapeutic properties, with the study indicating a 40%-60% acceptable response rate among patients undergoing this treatment.

A significant aspect of the study is the focus on treatment satisfaction, which is often overlooked in clinical assessments. Utilizing the Treatment Satisfaction Questionnaire for Medication (TSQM) Version II, the authors assessed the subjective experiences of Iranian patients receiving DPCP. This approach not only provides insight into the clinical effectiveness of the treatment but also gauges the psychological and emotional impacts of the therapy on patients, which are critical for comprehensive care.

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The findings suggest that while a notable percentage of patients experience positive outcomes in terms of hair regrowth, the overall prognosis remains guarded, particularly for those with extensive hair loss. The authors argue that monitoring patient satisfaction is essential for evaluating the quality of care, as it can influence adherence to treatment and overall health outcomes.

However, the study could benefit from a more extensive discussion on the long-term sustainability of hair regrowth following DPCP treatment, as well as potential side effects associated with prolonged use. Additionally, the demographic homogeneity of the sample may limit the generalizability of the findings to wider populations with AA.

The article titled "The Efficacy and Adverse Effects of Corticosteroid Pulse Therapy in Alopecia Areata: A Review Article" by (Rastaghi et al., 2023) provides a comprehensive examination of the role of corticosteroid pulse therapy in the management of alopecia areata (AA). The authors highlight the multifactorial nature of AA, emphasizing that both genetic and environmental factors contribute significantly to its pathology. This understanding is critical, as it sets the stage for evaluating treatment efficacy and the challenges associated with managing this condition.

The review articulates the range of severity in AA, from localized patches to complete loss of hair, which can have profound psychosocial impacts on affected individuals, including increased rates of depression, anxiety, and social phobia. This acknowledgment of the psychological ramifications of AA underscores the importance of effective treatment strategies not only for physical restoration but also for improving the overall quality of life for patients.

(Rastaghi et al., 2023) delve into the various treatment modalities available for AA, noting that the efficacy of these treatments can be highly variable. They specifically focus on corticosteroid therapy, which has been a cornerstone in the treatment of AA. The authors explain that while topical and intralesional corticosteroids are

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

generally effective for mild to moderate cases, severe AA presents a more complex challenge. In such cases, systemic corticosteroids are often employed, reflecting a need for a more aggressive treatment approach.

The introduction of corticosteroid pulse therapy by Burton and Shuster in 1975 is a pivotal point in the article. This method aims to mitigate the side effects associated with long-term corticosteroid use while enhancing treatment efficacy. The review meticulously assesses the literature surrounding pulse steroid therapy, evaluating its effectiveness, relapse rates, side effects, and prognostic factors influencing treatment responses. The authors employed a systematic literature search, ensuring that their review is grounded in relevant clinical trials, thereby enhancing the credibility of their findings.

However, while the article successfully synthesizes existing literature on corticosteroid pulse therapy, it could benefit from a more detailed discussion on the mechanisms by which corticosteroids exert their effects in AA. Additionally, the review could explore alternative treatment options and their comparative effectiveness, providing a broader context for clinicians considering various therapeutic strategies for AA.

CONCLUSION

The literature on hair regrowth outcomes for patients with alopecia areata (AA) provides a comprehensive understanding of the multifaceted nature of this autoimmune condition and the diverse therapeutic strategies employed to promote hair regrowth. Early contributions, such as those by (Gestetner, 2013), establish the foundational role of contact immunotherapy, particularly diphenylcyclopropenone (DPCP), in modulating the immune response to facilitate hair regrowth. The variability in patient responses underscores the need for tailored treatment approaches, as highlighted by (Mahasaksiri et al., 2021), which indicates that while DPCP can be effective, response rates can fluctuate significantly.

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Further exploration of treatment modalities reveals the importance of considering both clinical and psychosocial factors in managing AA. The systematic review by (Mostaghimi et al., 2021) emphasizes the emotional and social implications of hair loss, advocating for a holistic approach that integrates psychological support with medical treatments. This perspective is echoed in the findings of (R. Barton et al., 2021), which specifically addresses the unique treatment needs of pediatric patients, reinforcing the necessity for age-appropriate therapies.

Moreover, the investigation of corticosteroid pulse therapy by (Rastaghi et al., 2023) illustrates the challenges of managing severe cases of AA, where traditional therapies may be insufficient. Their review highlights the potential for corticosteroids to provide relief, yet it also points to the complexities involved in treating more extensive forms of the disease.

In conclusion, the body of literature surrounding hair regrowth outcomes in alopecia areata demonstrates a significant evolution in understanding both the clinical and psychosocial dimensions of the condition. The findings collectively advocate for a multifaceted approach that not only addresses the immunological aspects of treatment but also incorporates the emotional and social well-being of patients. This comprehensive perspective is essential for optimizing treatment outcomes and improving the quality of life for individuals affected by alopecia areata.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

REFERENCES

1. Gestetner, C., 2013. Alopecia Areata: An Overview. [PDF]
2. J. Dabek, R., G. Austen, W., & Bojovic, B., 2019. Laser-assisted Hair Regrowth: Fractional Laser Modalities for the Treatment of Androgenic Alopecia. ncbi.nlm.nih.gov
3. Mahasaksiri, T., Kositkuljorn, C., Anuntrangsee, T., & Suchonwanit, P., 2021. Application of Topical Immunotherapy in the Treatment of Alopecia Areata: A Review and Update. ncbi.nlm.nih.gov
4. Mostaghimi, A., Napatalung, L., Sikirica, V., Winnette, R., Xenakis, J., H. Zwillich, S., & Gorsh, B., 2021. Patient Perspectives of the Social, Emotional and Functional Impact of Alopecia Areata: A Systematic Literature Review. ncbi.nlm.nih.gov
5. R. Barton, V., Toussi, A., Awasthi, S., & Kiuru, M., 2021. Treatment of pediatric alopecia areata: A systematic review. ncbi.nlm.nih.gov
6. Abedini, R., Abdshah, A., Ghandi, N., Janatalipour, A., Torabi, S., & Nasimi, M., 2022. Treatment satisfaction and response in patients with severe alopecia areata under treatment with diphenylcyclopropenone. ncbi.nlm.nih.gov
7. Rastaghi, F., Kaveh, R., Yazdanpanah, N., Sadat Sahaf, A., & Ahramyanpour, N., 2023. The Efficacy and Adverse Effects of Corticosteroid Pulse Therapy in Alopecia Areata: A Review Article. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Dermatology and Venereology
Copyright © International Medical Journal Corp. Ltd. All rights reserved