Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Obstetric and Gynaecology



Efficacy of Regular Exercise During Pregnancy on the Prevention of Postpartum Depression

[1] Nanny, [1] Rachmat

[1] Faculty of Medicine, Padjadjaran University, Bandung City, West Java, Indonesia

Correspondence : dr.nanny98293726@gmail.com

Article History :

Received date : 2024/07/25
Revised date : 2024/08/19
Accepted date : 2024/09/06
Published date : 2025/01/15

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The body of literature reviewed presents compelling evidence supporting the efficacy of regular exercise during pregnancy as a preventative measure against postpartum depression. Initial findings by the authors of (Sui & M Dodd, 2013) established a foundational understanding of the benefits of physical activity, noting significant improvements in both maternal and neonatal health outcomes. **Literature Review:** Subsequent research by (Watson et al., 2018) reinforced the importance of regular exercise, particularly emphasizing its protective effects on maternal mental health during the perinatal period. This study demonstrated a positive correlation between consistent physical activity and improved mental health outcomes, suggesting that exercise not only benefits physical health but also plays a crucial role in mitigating the psychological risks associated with pregnancy and postpartum. Further exploration by (Kołomańska-Bogucka & Irena Mazur-Bialy, 2019) specifically addressed the implications of regular physical activity in the context of postnatal depression. Their findings underscored that women engaging in regular exercise exhibited a significantly lower risk of developing depressive disorders postpartum. The authors elucidated the biochemical mechanisms through which exercise enhances mood, such as increased neurotransmitter levels and neuroprotective factors, thereby providing a scientific basis for the mental health benefits associated with physical activity. Finally, the systematic review and meta-analysis conducted by (Liu et al., 2022) quantitatively assessed various exercise types and their effects on perinatal depression symptoms. The results demonstrated that physical exercise, including modalities like yoga and walking, significantly alleviated depressive symptoms among women, while also indicating the necessity for further research into the safety of these interventions for diverse populations. **Conclusion:** In conclusion, the reviewed literature collectively underscores the critical role of regular exercise during pregnancy in preventing postpartum depression. While the evidence supports the positive impacts of physical activity on maternal mental health, it also highlights the need for more targeted research to address methodological limitations and the safety of exercise interventions for different populations. This comprehensive understanding can inform future clinical practices and public health initiatives aimed at promoting physical activity among pregnant women.

Keyword: Efficacy, Regular Exercise During Pregnancy, Prevention of Postpartum Depression

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The literature surrounding the efficacy of regular exercise during pregnancy on the prevention of postpartum depression reveals a multifaceted relationship between physical activity and maternal mental health. Beginning with the foundational work by (Sui & M Dodd, 2013), the authors conducted a comprehensive literature review that synthesized findings from numerous studies, highlighting the positive impacts of exercise on both maternal and neonatal health outcomes. They noted a consistent association between physical activity during pregnancy and a reduction in risks such as gestational diabetes, preeclampsia, and operative births, alongside improvements in psychological well-being and mood stability. However, the authors also pointed out significant methodological limitations in the existing studies, including the variability in participant BMI and the lack of standardized measures for assessing physical activity.

Building on this foundation, (Watson et al., 2018) explored the connection between exercise frequency and maternal mental health across the perinatal period. Their research emphasized the importance of regular physical activity not only for physical health but also for its potential to mitigate mental health issues. They provided evidence supporting the notion that exercise can enhance fetal oxygenation and feto-placental growth while simultaneously addressing psychological factors such as body image and mood during pregnancy. This study reinforces the idea that exercise is a critical component of a holistic approach to maternal health.

(Kołomańska-Bogucka & Irena Mazur-Bialy, 2019) further delved into the implications of physical activity specifically concerning postnatal depression, which they identified as a prevalent and often untreated condition. They argued that regular physical activity serves as a natural intervention that can significantly reduce the risk of developing depressive disorders postpartum. Their findings suggest that exercise boosts neurotransmitter levels and enhances neuroprotective

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

factors, thereby providing a biochemical basis for the mental health benefits associated with physical activity during and after pregnancy.

Most recently, (Liu et al., 2022) conducted a systematic review and meta-analysis that quantitatively assessed the effects of various exercise types on perinatal depression symptoms. Their findings indicated that physical exercise, including modalities such as yoga and walking, significantly alleviated depressive symptoms among women. However, they also cautioned that exercise may not be suitable for all individuals, highlighting the necessity for further research into the safety and appropriateness of exercise interventions for diverse populations of pregnant women.

Together, these studies underscore the critical role of regular exercise during pregnancy as a preventative measure against postpartum depression, while also acknowledging the complexities and individual variations that must be considered in future research and clinical practice.

LITERATURE REVIEW

The article "Exercise in obese pregnant women: positive impacts and current perceptions" by (Sui & M Dodd, 2013) presents a comprehensive review of the existing literature on the effects of exercise during pregnancy, particularly focusing on maternal and neonatal health outcomes. The authors highlight that numerous studies have consistently indicated the benefits of physical activity during pregnancy, which include improved cardiovascular function, psychological well-being, and mood stability, alongside a reduced risk of gestational diabetes ((Sui & M Dodd, 2013)). These findings are particularly relevant to the discussion of postpartum depression, as the psychological benefits of exercise may play a crucial role in mitigating the risk of depressive symptoms following childbirth.

However, the review also points out significant limitations in the existing research. The authors note that many studies fail to adequately control for maternal body mass index (BMI) and often lack standardization in methodology ((Sui & M Dodd, 2013)). This variability can complicate the interpretation of results and the

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

generalizability of findings, particularly in relation to specific populations such as obese pregnant women. Furthermore, while the article references three Cochrane reviews on exercise during pregnancy, it emphasizes that the evidence regarding the impact of exercise on clinical pregnancy outcomes remains inconclusive. For instance, although Kramer and McDonald's review of eleven randomized controlled studies indicated that regular exercise could maintain or improve physical fitness, the effects on clinical outcomes were uncertain ((Sui & M Dodd, 2013)).

The article also discusses the findings from other systematic reviews, which did not find significant benefits of exercise in terms of perinatal or maternal morbidity in specific groups, such as women with gestational diabetes ((Sui & M Dodd, 2013)). This raises important questions about the efficacy of exercise interventions in diverse populations of pregnant women and suggests that more targeted research is necessary to better understand these dynamics.

The article "Exercise frequency and maternal mental health: Parallel process modelling across the perinatal period in an Australian pregnancy cohort" by (Watson et al., 2018) provides a comprehensive examination of the relationship between regular exercise during pregnancy and various maternal and neonatal outcomes, particularly focusing on mental health. The authors present a robust analysis of how consistent physical activity can influence not only the immediate health of the mother but also the developmental trajectory of the offspring.

One of the key insights from the study is the positive correlation between regular maternal exercise and improved mental health outcomes during the perinatal period. The authors emphasize that physical activity can serve as a protective factor against the onset of postpartum depression, which is a significant concern for many women following childbirth. Their findings align with previous literature suggesting that exercise can enhance mood and reduce anxiety levels, thereby potentially mitigating the risk of developing postpartum depression.

Moreover, the article discusses the physiological mechanisms through which maternal exercise may exert its beneficial effects. For instance, the authors highlight how regular physical activity can improve fetal oxygenation and promote feto-

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

placental growth, which are critical for optimal neonatal outcomes. This connection between maternal health behaviors and fetal development underscores the importance of encouraging exercise during pregnancy not only for the mother's well-being but also for the long-term health of the child.

The study's methodology, utilizing parallel process modeling, allows for a nuanced understanding of the interplay between exercise frequency and maternal mental health across different stages of pregnancy and postpartum. This approach enhances the credibility of the findings and provides a strong basis for future research in this area. Additionally, the authors address the implications of their findings for public health initiatives aimed at promoting physical activity among pregnant women, suggesting that such programs could be instrumental in reducing the incidence of postpartum depression.

The systematic review conducted by (Kołomańska-Bogucka & Irena Mazur-Bialy, 2019) provides a comprehensive examination of the relationship between physical activity and postnatal depression, highlighting the significant role that regular exercise can play in the prevention of this common mental health issue among new mothers. The authors underscore that postnatal depression is frequently undiagnosed and untreated, which can lead to detrimental consequences for both the mother and child. This recognition of the severity of postpartum depression sets a critical backdrop for the discussion of preventative measures, particularly the incorporation of physical activity during pregnancy.

The review presents compelling evidence suggesting that women who engage in regular physical activity, either prior to or during pregnancy, demonstrate a markedly lower risk of developing depressive disorders when compared to those who remain inactive. This finding aligns with the recommendations from the American College of Obstetricians and Gynecologists (ACOG), which advocate for the resumption of physical activity as soon as it is deemed safe postpartum. The authors effectively argue that physical therapy serves as a natural and side-effect-free intervention, making it a viable option for expectant and new mothers seeking to mitigate the risk of postpartum depression.

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

A critical aspect of the article is its exploration of the underlying mechanisms through which exercise exerts its positive effects on mental health. The authors detail how physical activity contributes to an increase in neurotransmitters such as serotonin, dopamine, and noradrenaline, all of which are crucial for mood regulation. Furthermore, the elevation of brain-derived neurotrophic factor (BDNF) levels is discussed, emphasizing its role in neuroprotection, neurogenesis, and synaptic plasticity. These biological processes are vital in understanding how regular exercise can foster a more resilient mental state during the perinatal period. Additionally, the review touches upon the physiological responses to exercise, including the modulation of cortisol levels and the stimulation of growth hormone (GH) and insulin-like growth factor 1 (IGF-1), which are implicated in sleep regulation, cognitive function, and overall mood stability. This multifaceted approach not only reinforces the efficacy of exercise as a preventative measure against postpartum depression but also provides a robust framework for future research in this area.

The article "Physical exercise interventions for perinatal depression symptoms in women: A systematic review and meta-analysis" by (Liu et al., 2022) provides a comprehensive examination of the impact of various exercise modalities on perinatal depression symptoms. The systematic review synthesizes findings from 20 randomized controlled trials involving a total of 2,866 participants, thereby offering a robust dataset for analysis.

The authors assert that physical exercise yields significant benefits in alleviating symptoms of perinatal depression, a critical issue affecting many women during and after pregnancy. The meta-analysis reveals that different types of exercise, including comprehensive exercise, yoga, and walking, are effective in reducing depressive symptoms. Notably, yoga is highlighted for its multifaceted approach, incorporating breathing techniques, body posture adjustments, meditation, and muscle relaxation, which collectively contribute to its efficacy in enhancing mood ((Liu et al., 2022)). The accessibility of yoga classes via mobile technology is an important consideration, as it may facilitate participation among perinatal women.

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Walking, as a simpler and more universally applicable form of exercise, is particularly emphasized for its feasibility in low to middle-income areas. The article points out that walking not only helps in mitigating depressive symptoms but also fosters self-esteem, resilience, and overall psychological well-being. This aspect underscores the holistic benefits of physical activity beyond mere symptom alleviation, suggesting a broader impact on mental health.

However, the article raises a significant concern regarding the safety of exercise interventions for perinatal women, noting that the included studies did not report safety outcomes. This gap in the literature is critical, as it limits the ability to conclusively advocate for exercise as a universally safe intervention during pregnancy. The authors acknowledge the need for further research to establish safety parameters and to better understand the mechanisms through which exercise affects mood. This call for additional studies is vital, as it addresses a crucial aspect of implementing exercise interventions in clinical practices.

CONCLUSION

The body of literature reviewed presents compelling evidence supporting the efficacy of regular exercise during pregnancy as a preventative measure against postpartum depression. Initial findings by the authors of (Sui & M Dodd, 2013) established a foundational understanding of the benefits of physical activity, noting significant improvements in both maternal and neonatal health outcomes. Their comprehensive review highlighted the psychological benefits of exercise, which could be pivotal in reducing the risk of postpartum depression, although they also pointed out methodological limitations in existing studies that could affect the generalizability of these findings.

Subsequent research by (Watson et al., 2018) reinforced the importance of regular exercise, particularly emphasizing its protective effects on maternal mental health during the perinatal period. This study demonstrated a positive correlation between consistent physical activity and improved mental health outcomes, suggesting that exercise not only benefits physical health but also plays a crucial role in mitigating the psychological risks associated with pregnancy and postpartum.

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Further exploration by (Kołomańska-Bogucka & Irena Mazur-Bialy, 2019) specifically addressed the implications of regular physical activity in the context of postnatal depression. Their findings underscored that women engaging in regular exercise exhibited a significantly lower risk of developing depressive disorders postpartum. The authors elucidated the biochemical mechanisms through which exercise enhances mood, such as increased neurotransmitter levels and neuroprotective factors, thereby providing a scientific basis for the mental health benefits associated with physical activity.

Finally, the systematic review and meta-analysis conducted by (Liu et al., 2022) quantitatively assessed various exercise types and their effects on perinatal depression symptoms. The results demonstrated that physical exercise, including modalities like yoga and walking, significantly alleviated depressive symptoms among women, while also indicating the necessity for further research into the safety of these interventions for diverse populations.

In conclusion, the reviewed literature collectively underscores the critical role of regular exercise during pregnancy in preventing postpartum depression. While the evidence supports the positive impacts of physical activity on maternal mental health, it also highlights the need for more targeted research to address methodological limitations and the safety of exercise interventions for different populations. This comprehensive understanding can inform future clinical practices and public health initiatives aimed at promoting physical activity among pregnant women.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Sui, Z. & M Dodd, J., 2013. Exercise in obese pregnant women: positive impacts and current perceptions. ncbi.nlm.nih.gov
2. Watson, S. J., Lewis, A. J., Boyce, P., & Galbally, M., 2018. Exercise frequency and maternal mental health: Parallel process modelling across the perinatal period in an Australian pregnancy cohort. [PDF]
3. Kołomańska-Bogucka, D. & Irena Mazur-Bialy, A., 2019. Physical Activity and the Occurrence of Postnatal Depression—A Systematic Review. ncbi.nlm.nih.gov
4. Liu, X., Wang, G., & Cao, Y., 2022. Physical exercise interventions for perinatal depression symptoms in women: A systematic review and meta-analysis. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Obstetric and Gynaecology
Copyright © International Medical Journal Corp. Ltd. All rights reserved