Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Orthopaedic and Traumatology



Management Of Femur Fracture

[1] Joshua, [1] Andrian

[1] Faculty of Medicine, Maranatha Christian University, Bandung City, West Java, Indonesia

Correspondence : dr.joshua9983747@gmail.com

Article History :

Received date : 2024/07/19
Revised date : 2024/08/17
Accepted date : 2024/09/28
Published date : 2025/01/10

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The management of femur fractures has evolved significantly, with a focus on optimizing patient outcomes through various treatment modalities. The literature indicates a shift towards surgical interventions, particularly in cases of complex fractures, where traditional nonoperative methods have shown limitations. **Literature Review:** (Boopathi, 2011) emphasized the challenges associated with nonoperative methods for managing compound supracondylar fractures, noting complications such as knee stiffness and delayed mobilization (Boopathi, 2011). This study underscored the necessity for surgical stabilization, advocating for anatomic reduction and early mobilization to enhance long-term recovery outcomes. (V. Neumann-Langen et al., 2023) further explored the surgical management of distal lower leg fractures, revealing that while intramedullary nailing is a preferred technique due to its effectiveness, it is not without significant postoperative complications (V. Neumann-Langen et al., 2023). This highlights the importance of careful surgical planning and technique selection to mitigate risks. In 2018, (Jackson et al., 2018) reviewed subtrochanteric proximal femur fractures, advocating for intramedullary fixation as the gold standard due to its advantages in reducing operative time and improving functional outcomes (Jackson et al., 2018). Their insights into classification systems further emphasize the need for tailored treatment based on specific fracture characteristics. **Conclusion:** In conclusion, the literature indicates a clear trend towards surgical management for femur fractures, particularly in complex cases and among the elderly. The findings across studies emphasize the significance of individualized treatment plans, timely intervention, and the careful selection of surgical techniques to optimize patient outcomes and minimize complications.

Keyword: Management, Femur Fracture

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The management of femur fractures encompasses a range of treatment modalities, each with distinct implications for patient outcomes. The literature reflects a progressive evolution in understanding these injuries and their optimal management strategies.

In 2011, K Boopathi's study highlighted the various treatment options available for compound supracondylar fractures of the femur, noting that while nonoperative methods often yielded better immediate outcomes, they were fraught with complications such as knee stiffness and delayed mobilization (Boopathi, 2011). This work underscored the necessity of surgical stabilization for distal femoral fractures, emphasizing the importance of anatomic reduction and early mobilization to mitigate long-term disability.

Advancing the discourse, Neumann et al. (V. Neumann et al., 2016) examined the challenges associated with surgical management of distal lower leg fractures, particularly the complications arising from intramedullary nailing and plate osteosynthesis. Their findings revealed that while intramedullary nailing is favored for its effectiveness, it is also associated with significant postoperative complications, necessitating careful consideration of surgical techniques and patient management (V. Neumann et al., 2016).

In 2018, Jackson, Tanios, and Ebraheim reviewed the management of subtrochanteric proximal femur fractures, advocating for intramedullary fixation as the gold standard due to its advantages in reducing operative time and improving functional outcomes (Jackson et al., 2018). Their analysis of classification systems, including the Russell-Taylor and AO systems, further emphasized the importance of tailored treatment approaches based on fracture characteristics.

Fischer et al. (Fischer et al., 2021) built upon these insights by focusing on the elderly demographic, where intramedullary nailing was again highlighted as the

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

preferred method for both intertrochanteric and subtrochanteric fractures. Their work pointed out that biomechanical advantages of intramedullary devices contribute to reduced soft tissue damage and enhanced recovery, reinforcing the trend towards surgical intervention in this population (Fischer et al., 2021).

A systematic review by Lodde et al. (F. Lodde et al., 2022) further elucidated the outcomes associated with double plating for femoral fractures, indicating that this technique may offer superior mechanical stability compared to other methods. This systematic approach to evaluating various surgical techniques provided a broader understanding of their effectiveness and complications (F. Lodde et al., 2022).

Sharma et al. (Sharma et al., 2022) addressed the critical issue of timing in surgical intervention for neck of femur fractures, linking early surgical management to better functional outcomes and reduced morbidity. Their findings underscored the public health implications of femoral neck fractures, particularly in terms of mortality and the need for effective treatment protocols (Sharma et al., 2022).

In a retrospective cohort study, Neumann-Langen et al. (V. Neumann-Langen et al., 2023) reinforced the significance of surgical techniques in achieving favorable outcomes in distal femoral fractures, noting that dual plating constructs were particularly beneficial for patients at risk of non-union. Their findings also suggest that innovative approaches, such as megaprothesis in elderly patients, could be advantageous in specific cases (V. Neumann-Langen et al., 2023).

Lastly, Fu and Zheng (Fu & Zheng, 2023) provided a narrative review on surgical treatments for femoral shaft fractures, emphasizing the importance of timely intervention and careful intraoperative techniques to prevent complications like malrotation (Fu & Zheng, 2023). Their insights into the need for comprehensive preoperative assessments and postoperative care highlight the multifaceted nature of femur fracture management.

This literature review will delve deeper into these studies, critically evaluating the evolving strategies and outcomes associated with the management of femur

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

fractures, ultimately aiming to provide a comprehensive understanding of current practices and future directions in this critical area of orthopedic care.

LITERATURE REVIEW

The article titled "Prospective Outcome Analysis of Various Methods of Management of Compound Supracondylar Fractures of Femur" by (Boopathi, 2011) provides a comprehensive examination of the treatment modalities available for distal femur fractures, particularly focusing on compound supracondylar fractures. The author highlights the evolution of treatment strategies, emphasizing that while skeletal traction has traditionally been employed for managing these injuries, it is associated with several complications, including angular deformity, joint incongruity, and delayed mobilization.

(Boopathi, 2011) argues convincingly that surgical intervention, specifically the use of an angled blade plate, has revolutionized the management of these fractures. The article underscores the importance of anatomic reduction of the articular surface and restoration of limb alignment, which are critical for ensuring favorable outcomes. Early mobilization is also emphasized as a key factor in recovery, demonstrating a shift in clinical practice towards more aggressive surgical approaches to mitigate the risks associated with conservative treatment.

Despite these advancements, the article does not shy away from acknowledging the ongoing challenges in managing distal femoral fractures. (Boopathi, 2011) notes that long-term disabilities can arise, particularly in cases involving extensive articular cartilage damage, significant bone comminution, and severe soft tissue injuries. The discussion on the limitations of conservative methods, which may lead to knee stiffness, malunion, and non-union, provides a balanced view of the complexities involved in treatment decision-making.

Moreover, the article highlights the variability in patient presentations, suggesting that no single implant or treatment approach is universally applicable. This point is particularly pertinent, as it reflects the need for individualized treatment plans that consider the specific characteristics of each fracture and the patient’s overall health

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

status. The association of osteoporosis and severe comminution with poor outcomes further complicates the management landscape, underscoring the necessity for thorough preoperative assessment and planning.

The article "Management of proximal femur fractures in the elderly: current concepts and treatment options" by (Fischer et al., 2021) provides a comprehensive overview of the management strategies for proximal femur fractures, particularly in the elderly population. The authors emphasize the paramount goal of treatment, which is to facilitate a return to the patient's previous level of activity and enable full weight-bearing post-injury.

A critical aspect of the article is its discussion of the preferred treatment modalities for intertrochanteric and subtrochanteric fractures. The authors advocate for intramedullary nailing as the treatment of choice due to its advantages, including reduced soft tissue damage and the ability to allow early weight-bearing (Fischer et al., 2021). This is particularly relevant in the elderly, where prolonged immobility can lead to significant complications. The article highlights the importance of evaluating the stability of the fracture pattern, particularly the integrity of the lateral cortical wall, which informs the choice of implant. The authors argue that extramedullary devices may be appropriate when this wall is intact, underscoring the necessity of a thorough evaluation of the fracture pattern prior to surgical intervention (Fischer et al., 2021).

Furthermore, the article notes the biomechanical advantages of intramedullary devices over extramedullary options, indicating that the Proximal Femoral Nail Antirotation (PFNA) is particularly effective, resulting in less blood loss and improved functional outcomes. The recommendation of long intramedullary nails as the gold standard for subtrochanteric fractures is well-supported, aligning with current best practices in orthopedic surgery (Fischer et al., 2021). Additionally, the authors introduce the concept of cement augmentation in osteosynthesis as a method to mitigate the risk of screw and blade cut-out in osteoporotic bone, thereby addressing a common challenge in this demographic.

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article titled "Outcome Evaluation of Distal Femoral Fractures Following Surgical Management: A Retrospective Cohort Study" by (V. Neumann-Langen et al., 2023) provides a comprehensive examination of the management outcomes for distal femur fractures, focusing on the efficacy of various surgical interventions. The authors conducted a retrospective cohort study to assess the influence of pre-existing risk factors on bone healing and to evaluate the effectiveness of different fixation methods.

One of the key findings of this study is the identification of a non-union rate of 7% following plate osteosynthesis, which is consistent with existing literature indicating that approximately 5% of distal femur fractures fail to heal adequately, irrespective of the treatment modality employed (V. Neumann-Langen et al., 2023). This statistic underscores the complexity of managing such fractures and the necessity for careful consideration of surgical techniques, particularly in patients with risk factors for instability or varus collapse.

The article emphasizes the advantages of dual plating constructs, which have been shown to possess greater mechanical strength compared to alternative fixation methods. This insight is particularly relevant for patients exhibiting risk factors that may predispose them to complications such as non-union (V. Neumann-Langen et al., 2023). Additionally, the authors highlight that the management of open type C3 distal femur fractures using a condylar locking plate in conjunction with an antibiotic-impregnated collagen sheet yields improved functional and radiological outcomes, suggesting a potential best practice for this specific fracture type.

Moreover, the authors discuss the consideration of megaprothesis implantation in elderly patients over 85 years of age, reporting favorable functional and radiological outcomes in a case series of 11 patients. This finding is particularly significant, as it suggests an alternative approach for a demographic that may be at higher risk for complications following traditional fixation methods (V. Neumann-Langen et al., 2023).

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The study also notes that screw fixation demonstrated superior knee function outcomes compared to plate osteosynthesis, with a notable extension deficit observed in 10% of patients following the latter method. This highlights the importance of selecting the appropriate surgical technique based on individual patient needs and the specific characteristics of the fracture.

The article titled "Surgical Treatments for Femoral Shaft Fractures: A Narrative Review" by (Fu & Zheng, 2023) offers a comprehensive overview of current practices and considerations in the surgical management of femoral shaft fractures. The authors emphasize the importance of timely intervention, noting that surgery should ideally occur within 48 hours of injury to optimize patient outcomes. This assertion is critical as delayed treatment can lead to complications and poorer prognoses, reinforcing the necessity for prompt surgical planning in clinical settings.

A significant focus of the article is on the technique of intramedullary nailing, which is a common surgical approach for these fractures. (Fu & Zheng, 2023) highlight the importance of assessing femoral rotation during this procedure to prevent malrotation, a complication that can lead to functional impairment and the need for additional surgical interventions. This critical evaluation of rotational alignment underscores the need for meticulous attention to detail during surgical procedures.

The article also discusses the choice between antegrade and retrograde nailing, indicating that both methods yield similar clinical outcomes. This finding suggests that the decision-making process may often be influenced by physician preference rather than a distinct superiority of one technique over the other. This aspect of the review prompts a reflection on the role of surgeon experience and comfort with specific techniques in influencing surgical outcomes.

Furthermore, the authors touch upon the management of complex femoral shaft fractures, emphasizing the necessity of approximating bony fragments to ensure optimal healing and alignment. This insight is particularly relevant for orthopedic

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

surgeons dealing with intricate cases where fracture configuration may complicate standard treatment protocols.

Another critical area addressed is venous thromboembolism (VTE) prophylaxis. The authors note that while aspirin or factor Xa inhibitors are not routinely approved for use in traumatic settings, ensuring adequate VTE prophylaxis remains vital, especially for patients with risk factors such as a history of VTE or coagulopathy. This highlights the multifaceted nature of managing femoral shaft fractures, where addressing potential complications is as crucial as the surgical intervention itself.

CONCLUSION

The management of femur fractures has evolved significantly, with a focus on optimizing patient outcomes through various treatment modalities. The literature indicates a shift towards surgical interventions, particularly in cases of complex fractures, where traditional nonoperative methods have shown limitations.

(Boopathi, 2011) emphasized the challenges associated with nonoperative methods for managing compound supracondylar fractures, noting complications such as knee stiffness and delayed mobilization (Boopathi, 2011). This study underscored the necessity for surgical stabilization, advocating for anatomic reduction and early mobilization to enhance long-term recovery outcomes.

(V. Neumann-Langen et al., 2023) further explored the surgical management of distal lower leg fractures, revealing that while intramedullary nailing is a preferred technique due to its effectiveness, it is not without significant postoperative complications (V. Neumann-Langen et al., 2023). This highlights the importance of careful surgical planning and technique selection to mitigate risks.

In 2018, (Jackson et al., 2018) reviewed subtrochanteric proximal femur fractures, advocating for intramedullary fixation as the gold standard due to its advantages in reducing operative time and improving functional outcomes (Jackson et al., 2018).

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Their insights into classification systems further emphasize the need for tailored treatment based on specific fracture characteristics.

(Fischer et al., 2021) reinforced these findings by focusing on the elderly population, where intramedullary nailing was highlighted as the preferred method for intertrochanteric and subtrochanteric fractures (Fischer et al., 2021). The biomechanical advantages of intramedullary devices were noted to contribute to reduced soft tissue damage and enhanced recovery, aligning with the trend towards surgical intervention in this demographic.

A systematic review by (F. Lodde et al., 2022) evaluated double plating for femoral fractures, indicating that this technique may offer superior mechanical stability compared to other methods (F. Lodde et al., 2022). This systematic approach provided a broader understanding of surgical effectiveness and complications.

(Sharma et al., 2022) addressed the timing of surgical intervention for neck of femur fractures, linking early management to better functional outcomes and reduced morbidity (Sharma et al., 2022). Their findings underscore the public health implications of timely treatment protocols in managing femoral neck fractures.

(V. Neumann-Langen et al., 2023) reinforced the importance of surgical techniques in achieving favorable outcomes in distal femoral fractures, noting that dual plating constructs were particularly beneficial for patients at risk of non-union (V. Neumann-Langen et al., 2023). Their exploration of innovative approaches, such as megaprothesis in elderly patients, suggests potential advancements in treatment strategies.

Lastly, (Fu & Zheng, 2023) discussed the importance of timely intervention and careful intraoperative techniques to prevent complications like malrotation in femoral shaft fractures (Fu & Zheng, 2023). Their insights into comprehensive preoperative assessments and postoperative care highlight the multifaceted nature of femur fracture management.

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

In conclusion, the literature indicates a clear trend towards surgical management for femur fractures, particularly in complex cases and among the elderly. The findings across studies emphasize the significance of individualized treatment plans, timely intervention, and the careful selection of surgical techniques to optimize patient outcomes and minimize complications.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Boopathi, K., 2011. Prospective Outcome Analysis of Various Methods of Management of Compound Supracondylar Fractures of Femur. [PDF]
2. V. Neumann, M., C. Strohm, P., Reising, K., Zwingmann, J., O. Hammer, T., & P. Suedkamp, N., 2016. Complications after surgical management of distal lower leg fractures. ncbi.nlm.nih.gov
3. Jackson, C., Tanios, M., & Ebraheim, N., 2018. Management of Subtrochanteric Proximal Femur Fractures: A Review of Recent Literature. ncbi.nlm.nih.gov
4. Fischer, H., Maleitzke, T., Eder, C., Ahmad, S., Stöckle, U., & F. Braun, K., 2021. Management of proximal femur fractures in the elderly: current concepts and treatment options. ncbi.nlm.nih.gov
5. F. Lodde, M., J. Raschke, M., Stolberg-Stolberg, J., Everding, J., Rosslenbroich, S., & C. Katthagen, J., 2022. Union rates and functional outcome of double plating of the femur: systematic review of the literature. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

6. Sharma, A., H Shanthappa, A., & Agarawal, S., 2022. Is Emergency Surgery an Indicator of Good Functional Outcomes in Neck of Femur Fractures Among Adults: A Prospective Clinical Study. ncbi.nlm.nih.gov
7. V. Neumann-Langen, M., Sontheimer, V., H. Borchert, G., Izadpanah, K., Schmal, H., & J. Kubosch, E., 2023. Outcome Evaluation of Distal Femoral Fractures Following Surgical Management: A Retrospective Cohort Study. ncbi.nlm.nih.gov
8. Fu, B. S. & Zheng, Z. H., 2023. Surgical Treatments for Femoral Shaft Fractures: A Narrative Review. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Orthopaedic and Traumatology
Copyright © International Medical Journal Corp. Ltd. All rights reserved