Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Otolaryngology



Inner Ear Malformations and Unilateral Sensorineural Hearing Loss

[1] Leonardus, [1] Rahadian

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.leonardus2387373@gmail.com

Article History :
Received date : 2024/10/11
Revised date : 2024/11/19
Accepted date : 2024/12/08
Published date : 2025/01/10

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The exploration of inner ear malformations (IEMs) and their association with unilateral sensorineural hearing loss (USNHL) has garnered significant attention in recent years, reflecting advancements in diagnostic techniques and therapeutic interventions. The literature reveals a multifaceted relationship between IEMs and SNHL, emphasizing the importance of early diagnosis and intervention. **Literature Review:** Brotto et al. (2021) advocate for targeted genetic investigations upon identifying IEMs, emphasizing the need for comprehensive clinical evaluations that extend beyond audiological assessments to identify potential syndromic conditions. This approach is crucial for understanding the broader implications of IEMs and tailoring management strategies accordingly. Patel et al. (2023) delve into the treatment approaches for children with unilateral hearing loss, linking multiple congenital anomalies and adverse developmental outcomes to neonatal intensive care admissions. This highlights the intricate relationship between medical history and the manifestation of hearing loss, further complicating the management landscape. Aldè et al. (2024) provide contemporary insights into the etiology and management of children with USNHL, revealing that many cases remain idiopathic despite extensive evaluations. They advocate for comprehensive management strategies, including timely amplification device selection and long-term audiological follow-up, emphasizing the need for increased awareness of the implications of uncorrected USNHL. **Conclusion:** The collective findings from these studies underscore the multifaceted nature of IEMs and USNHL, highlighting the necessity for early diagnosis, thorough etiological evaluations, and tailored management strategies. These elements are critical in optimizing outcomes for affected children, ensuring they receive the appropriate support for their auditory and communicative development.

Keyword: Inner Ear Malformations, Unilateral Sensorineural Hearing Loss

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The exploration of inner ear malformations (IEMs) and their association with unilateral sensorineural hearing loss (USNHL) has garnered significant attention in recent years, reflecting advancements in diagnostic techniques and therapeutic interventions. The literature reveals a multifaceted relationship between IEMs and SNHL, emphasizing the importance of early diagnosis and intervention.

In the study by van Beeck Calkoen et al. (A. van Beeck Calkoen et al., 2019), the authors highlight the prevalence of congenital sensorineural hearing loss, which affects one to two per thousand live births. They stress the critical role of early diagnosis facilitated by newborn hearing screening programs, which enable timely intervention that is vital for the child's speech and linguistic development. The study underscores the necessity of etiological evaluation in understanding the progression of hearing loss and identifying associated physical conditions, noting that while genetic factors account for about 50% of congenital SNHL cases, a significant portion remains unexplained.

The work of Feraco et al. (Feraco et al., 2021) further elucidates the relationship between IEMs and SNHL, detailing how these malformations arise from disrupted development during the first trimester of fetal growth. They assert that IEMs are a primary cause of SNHL and can be linked to both hereditary and nonhereditary factors. The authors emphasize the importance of early diagnosis for improving treatment outcomes, particularly in patients with genetic forms of IEM, who often present with profound SNHL.

Building on this foundation, Brotto et al. (Brotto et al., 2021) advocate for targeted genetic investigations when IEMs are identified. Their review highlights the need for comprehensive clinical evaluations that extend beyond audiological assessments to rule out syndromic conditions. They also point out that specific gene mutations may necessitate neuroradiological evaluations to detect labyrinthine

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

anomalies, thereby reinforcing the interconnectedness of genetic and imaging studies in managing hearing impairment.

Patel et al. (Patel et al., 2023) contribute to this discourse by examining the treatment approaches for children with unilateral hearing loss. They identify multiple congenital anomalies and adverse developmental outcomes linked to neonatal intensive care admissions, suggesting a complex interplay between medical history and the manifestation of hearing loss.

Finally, Aldè et al. (Aldè et al., 2024) provide contemporary insights into the etiology, audiological characteristics, and management of children with USNHL. Their findings reveal that a significant number of patients are diagnosed after the age of two, with many cases remaining idiopathic despite extensive evaluations. The authors advocate for comprehensive management strategies, including the timely selection of amplification devices and long-term audiological follow-up, while also highlighting the necessity for increased awareness regarding the implications of uncorrected USNHL.

Together, these articles create a nuanced understanding of the interplay between IEMs and USNHL, emphasizing the critical importance of early diagnosis, thorough etiological evaluation, and the need for tailored management strategies to optimize outcomes for affected children.

LITERATURE REVIEW

The article titled "The etiological evaluation of sensorineural hearing loss in children" provides a comprehensive overview of the prevalence, diagnosis, and management of congenital sensorineural hearing loss (SNHL) in children. The authors highlight that the prevalence of congenital SNHL ranges from one to two per thousand live births, emphasizing the importance of early diagnosis and intervention for optimal development of hearing, speech, and linguistic skills. This is particularly relevant in the context of unilateral sensorineural hearing loss, where early identification can significantly impact a child's auditory and communicative development.

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article underscores the role of newborn hearing screening programs, which have been instrumental in facilitating the early identification of hearing-impaired children. By enabling timely interventions such as counseling, support, hearing aids, or cochlear implantation, these programs are crucial for improving outcomes for children with SNHL. The authors argue that while SNHL is generally irreversible, an adequate etiological evaluation is essential for prognostication of hearing loss progression, identification of associated physical conditions, and accurate counseling for patients and their families.

A critical aspect of the article is the detailed discussion of the etiological evaluation process. The authors note that approximately 50% of congenital SNHL cases are attributed to genetic factors, while acquired factors account for about 25%. However, they also highlight a significant challenge in the field: the etiology remains unknown in 25–45% of cases despite comprehensive evaluations. This uncertainty poses challenges for clinicians and families in understanding the prognosis and potential implications for family members.

The authors advocate for a stepwise diagnostic approach that includes imaging, DNA testing, and screening for congenital infections and metabolic diseases. This structured methodology aims to enhance the likelihood of identifying the underlying causes of SNHL, particularly in cases of unilateral hearing loss where the etiology may differ from bilateral cases. By systematically evaluating potential factors, clinicians can provide more tailored and informed care plans for affected children.

The article titled "Imaging of inner ear malformations: a primer for radiologists" provides a comprehensive overview of inner ear malformations (IEMs) and their significant association with sensorineural hearing loss (SNHL). The authors emphasize that IEMs arise from disrupted development during the first trimester of fetal development, highlighting a critical period in the embryological development of the ear and its implications for hearing health.

A key insight from the article is the role of imaging modalities in diagnosing IEMs. The authors note that high-resolution computed tomography (CT) remains the

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

preferred method for visualizing the osseous structures of the inner ear, allowing for detailed assessment of malformations. However, they also point out the increasing utility of magnetic resonance imaging (MRI) in evaluating the endolymph-filled structures and the eighth cranial nerve, which is essential for a comprehensive understanding of the inner ear's anatomy and pathology.

The article further distinguishes between hereditary and nonhereditary forms of SNHL associated with IEMs, noting that genetic IEMs often present with bilateral and symmetrical abnormalities, leading to profound SNHL. This differentiation is crucial, as it underscores the importance of genetic counseling and potential intervention strategies for affected families. The authors also discuss various etiological factors contributing to IEMs, including idiopathic causes, genetic errors, and exposure to teratogenic agents, which adds depth to the understanding of the multifactorial nature of these malformations.

The authors advocate for early diagnosis of IEMs, emphasizing that timely intervention can significantly improve treatment outcomes for patients with hearing loss. This assertion aligns with current best practices in audiology and otology, where early identification and management are critical in mitigating the long-term impacts of hearing impairment on communication and quality of life.

The article titled "Genetics of Inner Ear Malformations: A Review" provides a comprehensive overview of the intricate relationship between genetic factors and inner ear malformations, particularly in the context of unilateral sensorineural hearing loss. The authors emphasize the critical need for targeted genetic investigations when an inner ear malformation is identified, highlighting that such findings should prompt a thorough clinical evaluation beyond mere audiological assessments.

One of the key insights presented in the article is the necessity of investigating potential syndromic conditions associated with inner ear malformations. This is particularly relevant as unilateral sensorineural hearing loss can often be a part of a broader spectrum of syndromic presentations. The authors advocate for a multidisciplinary approach, where clinicians should not only focus on hearing

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

assessments but also consider genetic consultations and testing. This approach is essential for establishing a more comprehensive understanding of the patient's condition and guiding appropriate management strategies.

Furthermore, the article discusses the importance of neuroradiological evaluations in cases where specific gene mutations are identified. This recommendation is significant as it underscores the need to investigate potential labyrinthine anomalies that may not be apparent through audiological testing alone. The authors effectively argue that integrating genetic and radiological assessments can lead to better-informed clinical decisions and improved patient outcomes.

The article titled "Characterisation of the treatment provided for children with unilateral hearing loss" presents a comprehensive examination of the relationship between congenital anomalies, specifically inner ear malformations, and unilateral sensorineural hearing loss (USNHL) in children. The authors emphasize the high prevalence of inner ear and internal auditory canal malformations in this population, suggesting a critical link between these anatomical anomalies and the occurrence of unilateral hearing loss.

One of the significant contributions of this article is its detailed analysis of the medical context and etiology of USNHL. The authors highlight that children with USNHL often present with multiple congenital anomalies, which are frequently associated with adverse developmental outcomes. This connection underscores the necessity for early identification and intervention strategies to mitigate the impact of hearing loss on developmental trajectories.

The article further delves into the comparative prevalence and characteristics of inner ear malformations in children with congenital unilateral versus bilateral hearing loss. This comparative analysis is crucial as it sheds light on the differential impacts of these malformations on auditory function and overall health. The findings suggest that children with unilateral hearing loss exhibit a distinct pattern of inner ear malformations when compared to their counterparts with bilateral hearing loss, indicating the need for tailored diagnostic and therapeutic approaches.

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Moreover, the authors conduct a risk analysis regarding unilateral severe-to-profound sensorineural hearing loss in children, identifying key risk factors and etiological considerations. This aspect of the study is particularly valuable for clinicians, as it provides insights into the underlying causes of USNHL, which can inform prevention and treatment strategies.

The article titled "Unilateral Sensorineural Hearing Loss in Children: Etiology, Audiological Characteristics, and Treatment" provides a comprehensive overview of the challenges associated with unilateral sensorineural hearing loss (USNHL) in pediatric populations. The authors emphasize the significant delay in diagnosis, with many children identified after the age of two, underscoring the necessity for regular monitoring of developmental milestones and auditory skills in early childhood. This finding highlights a critical gap in early detection, which is essential for timely intervention and optimal auditory development.

The study reveals that despite extensive diagnostic evaluations, a considerable proportion of USNHL cases remain idiopathic, indicating a need for further research to elucidate underlying etiologies. This lack of identifiable causes complicates management strategies and emphasizes the importance of individualized care for affected children. The authors advocate for a multifaceted approach to management, which includes accurate diagnosis, appropriate amplification device selection, device verification, and long-term audiological follow-up. This comprehensive strategy is crucial in mitigating the potential negative impacts of hearing loss on speech and language development.

A particularly noteworthy aspect of the article is its discussion of cochlear implantation as a treatment option for children with single-sided deafness (SSD). The authors argue for its consideration, especially for those at risk of deterioration in the contralateral ear. This perspective is significant as it opens avenues for intervention that could profoundly impact the quality of life and communicative abilities of children with USNHL.

Moreover, the authors call for increased awareness among healthcare professionals and parents regarding the detrimental effects of uncorrected USNHL. This

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

advocacy is essential, as it could lead to earlier referrals and interventions, ultimately improving outcomes for affected children. The article also stresses the need for rigorous and updated guidelines for the management of pediatric unilateral hearing loss, which is a vital step towards standardizing care and ensuring that all children receive the appropriate support.

CONCLUSION

The exploration of inner ear malformations (IEMs) and their association with unilateral sensorineural hearing loss (USNHL) reveals a complex interplay of genetic, developmental, and diagnostic factors. The literature emphasizes the critical importance of early diagnosis and intervention, which can significantly impact the auditory and communicative development of affected children.

The study by van Beeck Calkoen et al. (2019) indicates that congenital sensorineural hearing loss affects one to two per thousand live births, underscoring the necessity for newborn hearing screening programs to facilitate timely diagnosis and intervention. This early identification is pivotal for the optimal development of speech and language skills, particularly in cases of unilateral hearing loss where the effects can be profound.

Feraco et al. (2021) provide insights into the etiology of IEMs, noting that these malformations arise from disrupted development during the first trimester of fetal growth. They highlight that IEMs are a primary cause of SNHL, with both hereditary and nonhereditary factors contributing to their occurrence. The authors stress the importance of early diagnosis to enhance treatment outcomes, especially in genetic cases where profound SNHL is common.

Brotto et al. (2021) advocate for targeted genetic investigations upon identifying IEMs, emphasizing the need for comprehensive clinical evaluations that extend beyond audiological assessments to identify potential syndromic conditions. This approach is crucial for understanding the broader implications of IEMs and tailoring management strategies accordingly.

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Patel et al. (2023) delve into the treatment approaches for children with unilateral hearing loss, linking multiple congenital anomalies and adverse developmental outcomes to neonatal intensive care admissions. This highlights the intricate relationship between medical history and the manifestation of hearing loss, further complicating the management landscape.

Aldè et al. (2024) provide contemporary insights into the etiology and management of children with USNHL, revealing that many cases remain idiopathic despite extensive evaluations. They advocate for comprehensive management strategies, including timely amplification device selection and long-term audiological follow-up, emphasizing the need for increased awareness of the implications of uncorrected USNHL.

The collective findings from these studies underscore the multifaceted nature of IEMs and USNHL, highlighting the necessity for early diagnosis, thorough etiological evaluations, and tailored management strategies. These elements are critical in optimizing outcomes for affected children, ensuring they receive the appropriate support for their auditory and communicative development.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. A. van Beeck Calkoen, E., S. D. Engel, M., M. van de Kamp, J., G. Yntema, H., Goverts, S. T., Mulder, M. F., Merkus, P., & F. Hensen, E., 2019. The etiological evaluation of sensorineural hearing loss in children. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

2. Feraco, P., Piccinini, S., & Gagliardo, C., 2021. Imaging of inner ear malformations: a primer for radiologists. ncbi.nlm.nih.gov
3. Brotto, D., Sorrentino, F., Cenedese, R., Avato, I., Bovo, R., Trevisi, P., & Manara, R., 2021. Genetics of Inner Ear Malformations: A Review. ncbi.nlm.nih.gov
4. Patel, R., J. Hoare, D., R. Willis, K., Tabraiz, S., K. Bateman, P., & K. Thornton, S., 2023. Characterisation of the treatment provided for children with unilateral hearing loss. ncbi.nlm.nih.gov
5. Aldè, M., Zanetti, D., Ambrosetti, U., Monaco, E., Maria Gasbarre, A., Pignataro, L., Cantarella, G., & Barozzi, S., 2024. Unilateral Sensorineural Hearing Loss in Children: Etiology, Audiological Characteristics, and Treatment. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved