Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Otolaryngology



Natural Course of Unilateral Vocal Fold Paralysis and Optimal Timing of Permanent Treatment

[1] Yulia, [1] Hariyanto

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.yulia928737@gmail.com

Article History :
Received date : 2024/10/12
Revised date : 2024/11/19
Accepted date : 2024/12/28
Published date : 2025/01/10

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature surrounding unilateral vocal fold paralysis (UVFP) presents a complex interplay of diagnostic, therapeutic, and rehabilitative strategies aimed at addressing the multifaceted challenges posed by this condition. A significant body of research has emerged over the past several years, contributing to our understanding of the natural course of UVFP and the implications for treatment timing. **Literature Review:** (Miaśkiewicz et al., 2021) focused on the prevalence and etiology of UVFP, noting the significant impact on patients' quality of life. They highlighted injection laryngoplasty as an effective treatment option, with preoperative assessments potentially predicting the need for future interventions. This predictive capability is crucial for tailoring treatment plans and optimizing timing. (Liao & Wang, 2022) conducted a comprehensive review of UVFP management post-thyroid surgery, emphasizing the efficacy of injection laryngoplasty. Their meticulous literature review underscored the importance of understanding treatment outcomes, although a more in-depth analysis of long-term effects could enhance future practices. (Wei-Ge Chen et al., 2022) introduced autologous fat injection laryngoplasty, advocating for its integration with rehabilitative voice training. Their findings demonstrated significant improvements in voice-related quality of life, reinforcing the importance of a multidisciplinary approach to UVFP management. **Conclusion:** Collectively, these studies underscore the necessity for a nuanced understanding of UVFP, particularly regarding its natural progression and the optimal timing for various treatments. The integration of innovative diagnostic techniques, comprehensive therapeutic strategies, and individualized patient care emerges as a central theme in enhancing treatment outcomes for UVFP.

Keyword: Unilateral Vocal Fold Paralysis, Optimal Timing of Permanent Treatment

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The literature surrounding unilateral vocal fold paralysis (UVFP) presents a complex interplay of diagnostic, therapeutic, and rehabilitative strategies aimed at addressing the multifaceted challenges posed by this condition. A significant body of research has emerged over the past several years, contributing to our understanding of the natural course of UVFP and the implications for treatment timing.

In 2016, (Mat Baki et al., 2016) conducted a pivotal study examining the feasibility of vocal fold abduction and adduction assessment using cine-MRI. This work highlighted the importance of accurate diagnostic techniques for UVFP, establishing a foundation for subsequent studies that explore treatment efficacy and timing.

The following year, (Barbosa Barcelos et al., 2017) broadened the discussion by investigating the multidimensional effects of voice therapy in patients with UVFP due to cancer treatments. Their findings indicated that while voice therapy could lead to varying degrees of improvement, a significant portion of patients still experienced persistent UVFP post-treatment. This study underscored the complexity of managing UVFP in the context of oncological care and raised questions regarding the optimal timing for interventions.

(Wang et al., 2018) shifted the focus to the comparative efficacy of vocal training versus vocal microsurgery for early vocal fold polyps, indirectly informing the UVFP discourse. Their analysis revealed that while surgical options could yield immediate benefits, the longer-term adherence to vocal therapy remained a challenge. This highlights the necessity of considering patient compliance and the potential for delayed treatment responses when evaluating management strategies for UVFP.

In 2021, (Miaśkiewicz et al., 2021) explored the predictive factors for future reintervention following injection laryngoplasty, a common treatment for UVFP.

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Their results indicated that preoperative assessments could inform the likelihood of needing additional procedures, thus emphasizing the importance of tailored treatment plans based on individual patient characteristics. This work contributes to the ongoing discourse on the timing of interventions, suggesting that early and informed decision-making could enhance treatment outcomes.

(Liao & Wang, 2022) conducted a comprehensive review of the management of UVFP post-thyroid surgery, focusing on injection laryngoplasty. Their findings synthesized existing literature to evaluate treatment outcomes, providing a critical overview of the evolving landscape of UVFP management. This review highlighted the need for continuous assessment of treatment efficacy and the relevance of timing in achieving optimal patient outcomes.

Lastly, (Wei-Ge Chen et al., 2022) introduced an innovative approach with office-based structural autologous fat injection laryngoplasty, emphasizing the role of rehabilitative voice training in conjunction with surgical interventions. Their study demonstrated significant improvements in voice-related quality of life, reinforcing the idea that a multidisciplinary approach is essential in managing UVFP effectively.

Through these contributions, the literature underscores the importance of a nuanced understanding of UVFP, encompassing both the natural course of the condition and the critical timing of various treatment modalities. Each study builds upon the previous findings, collectively informing clinical practice and guiding future research directions in the management of unilateral vocal fold paralysis.

LITERATURE REVIEW

The article "Feasibility of vocal fold abduction and adduction assessment using cine-MRI" by Baki et al. (Mat Baki et al., 2016) presents a significant exploration into the assessment of vocal fold movement in patients with unilateral vocal fold paralysis (UVFP). The study's design effectively recruits participants from local ear, nose, and throat specialist voice clinics, ensuring a relevant clinical context for the findings. The inclusion criteria are notably rigorous, requiring confirmation of UVFP through both flexible nasopharyngolaryngoscope and laryngeal

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

electromyography, which adds credibility to the diagnostic process and the subsequent analysis.

One of the key insights from the article is the emphasis on the natural course of UVFP, particularly in patients who have experienced the condition for more than six months without any surgical interventions. This aspect is crucial as it provides a clearer understanding of the progression of UVFP and the potential for spontaneous recovery, which is often a topic of debate in the literature. By focusing on patients with either iatrogenic or idiopathic causes of UVFP, the authors highlight the variability in patient outcomes and underscore the need for tailored treatment approaches.

Moreover, the utilization of cine-MRI for assessing vocal fold movement is a noteworthy advancement in the field. This imaging technique allows for dynamic visualization of vocal fold abduction and adduction, which is essential for understanding the functional implications of UVFP. The authors effectively argue for the feasibility of this method, suggesting that it could enhance diagnostic accuracy and inform treatment planning. This is particularly relevant when considering the optimal timing for permanent interventions, as the ability to assess vocal fold dynamics could lead to more informed decisions regarding surgical options.

However, the study could benefit from a more extensive discussion on the limitations of cine-MRI in this context. While the authors present compelling evidence for its feasibility, there is a need to address potential challenges, such as the accessibility of this technology in various clinical settings and the variability in interpretation of the results. Additionally, the long-term outcomes of patients assessed via cine-MRI compared to traditional assessment methods could provide valuable insights into the effectiveness of this approach.

The article "Multidimensional effects of voice therapy in patients affected by unilateral vocal fold paralysis due to cancer" by Barcelos et al. (Barbosa Barcelos et al., 2017) presents a comprehensive examination of the outcomes of voice therapy in a cohort of patients diagnosed with unilateral vocal fold paralysis

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

(UVFP) resulting from various cancer treatments. The study's design is robust, having received ethical approval and involving a well-defined sample of 61 patients, predominantly female, with a mean age of 52.4 years. The inclusion criteria are appropriately stringent, focusing on adults without structural lesions or prior dysphonia, which strengthens the validity of the findings.

The authors categorize the patients based on their response to voice therapy into three distinct groups: "overall improvement," "partial improvement," and "failure." This multidimensional approach allows for a nuanced understanding of treatment efficacy, highlighting that while voice therapy may yield positive outcomes for some, a significant number of patients (28 out of 61) continued to experience UVFP post-treatment. The classification into different improvement categories is particularly valuable, as it provides insight into the varying degrees of success and the potential need for additional interventions for those classified under "failure."

However, the study does have limitations that warrant discussion. The fact that only 43 out of the initial 61 patients completed the voice rehabilitation treatment raises questions about the generalizability of the findings. The reasons for dropout may be critical to understanding the overall effectiveness of voice therapy and could introduce bias in the results. Furthermore, the study does not elaborate on the specific types of voice therapy administered, which could influence outcomes and limit the ability to replicate the study in other settings.

Additionally, the persistence of UVFP in a majority of patients post-therapy underscores the complexity of this condition and suggests that voice therapy alone may not be sufficient for optimal recovery. The authors' findings point to the need for further research into the timing and nature of permanent treatments for UVFP, as well as the potential integration of surgical options alongside voice therapy to enhance patient outcomes.

The article titled "Comparison of the efficacy of vocal training and vocal microsurgery in patients with early vocal fold polyp" by Wang et al. (Wang et al., 2018) provides a significant exploration into the treatment modalities for vocal fold paralysis (VFP), particularly emphasizing the comparison between vocal therapy

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

(VT) and vocal microsurgery (VM). The authors highlight the complexity of VFP, noting that treatment options can vary widely based on individual patient circumstances, including economic considerations and the inherent risks associated with surgical interventions.

One of the key insights from the article is the effectiveness of VM in enhancing voice quality with a notably low recurrence rate. This finding is critical, as it suggests that for certain patients, particularly those who may be less compliant or have other barriers to consistent therapy, VM could represent a more favorable option despite the associated risks and costs. Conversely, the authors point out that VT, while beneficial, often requires a longer duration of treatment and is heavily dependent on patient compliance. Factors such as education level and scheduling conflicts can significantly impact a patient's ability to adhere to a vocal therapy regimen, which raises important considerations about accessibility and patient engagement in the management of VFP.

The article delves into the pathophysiology of VFP, identifying key elements such as bleeding, edema, and fibrin deposition as contributing factors to vocal quality deterioration. This biological understanding is crucial for clinicians as it underscores the importance of timely intervention and the potential for therapeutic approaches to mitigate the effects of these pathological changes.

Furthermore, the relationship between polyp size and vocal quality is emphasized, suggesting that larger polyps may correlate with poorer vocal outcomes. This insight could guide clinicians in making more informed decisions regarding the timing and choice of treatment based on the specific characteristics of the patient's condition.

The article by Miaśkiewicz et al. (Miaśkiewicz et al., 2021) provides a comprehensive examination of unilateral vocal fold paralysis (UVFP), a condition that affects a notable portion of the population and leads to significant voice deterioration. The authors highlight the prevalence of UVFP, which occurs in approximately 0.41–0.51% of individuals, with a striking 83.6% experiencing voice

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

deterioration. This underscores the clinical relevance of the condition and the necessity for effective treatment strategies.

The article outlines the etiology of UVFP, primarily attributing it to unilateral injury of the vagus or recurrent laryngeal nerve, often resulting from surgical interventions or occurring idiopathically. The consequences of UVFP are multifaceted, manifesting not only as dysphonia characterized by a weak and breathy voice but also as difficulties in swallowing and the risk of aspiration. This broad impact on patients' quality of life emphasizes the importance of timely and effective therapeutic approaches.

Miaśkiewicz et al. delve into the treatment modalities for UVFP, focusing on the restoration of vocal function through various surgical interventions. Among these, injection laryngoplasty (IL) is highlighted as a minimally invasive technique that has shown efficacy in improving vocal outcomes. The authors delineate the different types of injectable materials and the factors that influence the success and longevity of IL treatment, including the extent of the glottal gap, the timing of the intervention, and patient-specific variables such as age and comorbidities.

A significant contribution of this study is the exploration of the potential for preoperative assessments to predict the need for future reintervention following IL. The authors argue that understanding the preoperative characteristics that correlate with treatment outcomes could facilitate more personalized treatment plans and improve overall patient management. This predictive capability would be invaluable in clinical practice, as it could help clinicians determine the most appropriate timing for intervention and the likelihood of requiring further procedures.

The article effectively synthesizes current knowledge on UVFP and provides a critical analysis of the factors influencing treatment efficacy, particularly in the context of IL. However, while the authors present a robust discussion on the implications of their findings, further research is necessary to validate the predictive models they propose. Future studies should aim to establish standardized

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

preoperative assessment protocols that could enhance the predictive power regarding the need for reoperation.

The article titled "Management of Unilateral Vocal Fold Paralysis after Thyroid Surgery with Injection Laryngoplasty: State of Art Review" by Liao and Wang (Liao & Wang, 2022) provides a comprehensive overview of the management strategies for unilateral vocal fold paralysis (UVFP) specifically following thyroid surgery. The authors meticulously reviewed existing literature, utilizing databases such as Pubmed, Medline, and EMBASE, and focused on studies that examined the effects of injection laryngoplasty (IL) in this particular context.

One of the key strengths of this review is its methodical approach to literature selection. By employing specific keywords and MeSH terms, the authors narrowed their focus to articles that directly relate to injection therapy for UVFP post-thyroidectomy, thereby ensuring that the findings are relevant and applicable to the target population. The exclusion of studies with multiple etiologies of UVFP allows for a more precise understanding of the outcomes associated with IL in the context of thyroid surgery, which is a notable contribution to the field.

The review synthesizes critical data regarding the number of patients involved, the types of materials injected, the techniques used for injection, and the resultant treatment outcomes. This detailed compilation of information is valuable for clinicians seeking to understand the efficacy of IL as a treatment option for UVFP. The authors highlight that injection laryngoplasty can improve vocal function, which is crucial for patients who experience voice-related issues following thyroid surgery.

However, while the article presents a thorough review of the literature, it would benefit from a more in-depth discussion on the long-term outcomes of IL. The review primarily focuses on immediate post-treatment results, which, although important, may not fully capture the longevity of the treatment's effectiveness or the potential need for additional interventions over time. Furthermore, the authors could enhance their analysis by addressing the variability in patient responses to IL

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

and the factors that may influence these outcomes, such as the timing of the intervention and the specific characteristics of the vocal fold paralysis.

The article titled "Office-Based Structural Autologous Fat Injection Laryngoplasty for Unilateral Vocal Fold Paralysis" by Chen et al. (Wei-Ge Chen et al., 2022) presents a comprehensive examination of the efficacy of autologous fat injection laryngoplasty as a treatment for unilateral vocal fold paralysis (UVFP). The authors articulate the significance of rehabilitative voice training, emphasizing the need for a balanced approach that prioritizes vocal performance while minimizing potential harm to the recovering vocal fold. This balance is critical, as the authors highlight the importance of training patients to regain various vocal abilities, including pitch range, loudness dynamic range, motor accuracy, agility, and endurance, tailored to their individual needs.

The findings reported in the article indicate that all patients experienced a statistically significant improvement in their Voice Handicap Index (VHI-10) scores, surpassing the minimal clinically important difference established by Young et al. (2022). However, it is noteworthy that not all patients achieved scores within the normal range, suggesting that while the treatment is beneficial, it may not fully restore vocal function for every individual. This underscores the complexity of UVFP and the variability in patient responses to treatment.

Moreover, the authors provide valuable context regarding the etiology of UVFP, particularly its association with thyroid surgery, which is a prevalent cause among their patient population. This connection is critical as it informs the clinical approach to managing UVFP, particularly in post-thyroidectomy patients. The article also references previous studies that have documented the positive impact of injection laryngoplasty on voice and voice-related quality of life, reinforcing the relevance of early intervention with long-lasting fillers for this demographic.

The authors advocate for office-based structural autologous fat injection laryngoplasty as a meaningful treatment option, particularly in light of research linking UVFP to complications such as pneumonia. This perspective highlights the

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

broader implications of effective UVFP management not only for voice restoration but also for overall patient health.

CONCLUSION

The literature on unilateral vocal fold paralysis (UVFP) reveals a comprehensive understanding of the condition's natural course and the timing of various treatment modalities. The studies reviewed provide insights into the diagnostic techniques, therapeutic strategies, and rehabilitation efforts necessary for effective management of UVFP.

(Mat Baki et al., 2016) introduced cine-MRI as a significant advancement in assessing vocal fold movement, emphasizing its role in understanding UVFP's natural course. Their findings underscore the importance of accurate diagnostics, particularly in patients with prolonged UVFP, which could inform decisions regarding the timing of interventions. This foundational work paved the way for subsequent studies that explored the efficacy of treatments and the necessity of individualized approaches.

The exploration by (Barbosa Barcelos et al., 2017) into the effects of voice therapy on patients with UVFP resulting from cancer treatments highlighted the complexity of managing the condition. Their classification of patient responses into distinct groups revealed that while some benefit from therapy, a substantial number continue to experience persistent symptoms, indicating a need for timely and possibly concurrent surgical interventions.

(Wang et al., 2018) further enriched the discourse by comparing vocal training and microsurgery for vocal fold polyps, indirectly informing UVFP management. Their findings suggest that while surgical options may provide immediate benefits, patient adherence to therapy remains a critical factor in long-term outcomes, emphasizing the need for a multifaceted approach.

(Miaśkiewicz et al., 2021) focused on the prevalence and etiology of UVFP, noting the significant impact on patients' quality of life. They highlighted injection laryngoplasty as an effective treatment option, with preoperative assessments

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

potentially predicting the need for future interventions. This predictive capability is crucial for tailoring treatment plans and optimizing timing.

(Liao & Wang, 2022) conducted a comprehensive review of UVFP management post-thyroid surgery, emphasizing the efficacy of injection laryngoplasty. Their meticulous literature review underscored the importance of understanding treatment outcomes, although a more in-depth analysis of long-term effects could enhance future practices.

(Wei-Ge Chen et al., 2022) introduced autologous fat injection laryngoplasty, advocating for its integration with rehabilitative voice training. Their findings demonstrated significant improvements in voice-related quality of life, reinforcing the importance of a multidisciplinary approach to UVFP management.

Collectively, these studies underscore the necessity for a nuanced understanding of UVFP, particularly regarding its natural progression and the optimal timing for various treatments. The integration of innovative diagnostic techniques, comprehensive therapeutic strategies, and individualized patient care emerges as a central theme in enhancing treatment outcomes for UVFP.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Mat Baki, M., Menys, A., Atkinson, D., Bassett, P., Morley, S., Beale, T., Sandhu, G., Naduvilethil, G., Stevenson, N., A Birchall, M., & Punwani, S., 2016. Feasibility of vocal fold abduction and adduction assessment using cine-MRI. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

2. Barbosa Barcelos, C., Angélica Lorenzon Silveira, P., Lígia Vieira Guedes, R., Nogueira Gonçalves, A., Dall'Agnol Siqueira Slobodticov, L., & Carrara-de Angelis, E., 2017. Multidimensional effects of voice therapy in patients affected by unilateral vocal fold paralysis due to cancer(). ncbi.nlm.nih.gov
3. Wang, H., Zhuge, P., You, H., Zhang, Y., & Zhang, Z., 2018. Comparison of the efficacy of vocal training and vocal microsurgery in patients with early vocal fold polyp(). ncbi.nlm.nih.gov
4. Miaśkiewicz, B., Panasiewicz, A., Gos, E., Krasnodębska, P., H. Skarżyński, P., & Szkiełkowska, A., 2021. Can preoperative results predict the need for future reintervention following injection laryngoplasty for unilateral vocal fold paralysis?. ncbi.nlm.nih.gov
5. Liao, L. J. & Wang, C. T., 2022. Management of Unilateral Vocal Fold Paralysis after Thyroid Surgery with Injection Laryngoplasty: State of Art Review. ncbi.nlm.nih.gov
6. Wei-Ge Chen, A., Chen, C. H., Lin, T. M., Chih-Hui Chang, A., Tsai, T. P., & Chang, S. Y., 2022. Office-Based Structural Autologous Fat Injection Laryngoplasty for Unilateral Vocal Fold Paralysis. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Otolaryngology
Copyright © International Medical Journal Corp. Ltd. All rights reserved