Research Article Volume 01, Issue No. 02. 2025



The Medical Journal of Plastic Surgery and Aesthetic Medicine

Complications in Brachioplasty

[1] Antonius, [2] Yohan

[1] Faculty of Medicine, Maranatha Christian University, Bandung City, West Java, Indonesia

Correspondence : dr.antonius3938747@gmail.com

Article History :

Received date : 2024/06/14
Revised date : 2024/07/18
Accepted date : 2024/08/06
Published date : 2025/01/05

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature on complications in brachioplasty and related body contouring procedures reveals a multifaceted understanding of surgical techniques, patient demographics, and psychosocial impacts. The studies reviewed indicate that the choice of surgical method has significant implications for both physical outcomes and patient satisfaction. **Literature Review:** In the study by (Freitas-Silva et al., 2010), the authors found that while different surgical interventions in breast cancer survivors did not significantly affect the incidence of arm edema, they did impact the range of motion (ROM) and overall satisfaction, suggesting that surgical choices play a crucial role in postoperative recovery. This finding underscores the need for further research into how specific surgical techniques can influence complications and patient experiences. The introduction of the SERI surgical scaffold as an adjunct in brachioplasty, discussed by , highlights innovations aimed at reducing complications such as scar hypertrophy, which are common in conventional methods. This advancement suggests a potential for improved outcomes through enhanced fascial support, although further investigation into long-term effectiveness is warranted. Moreover, the review by (Vidal et al., 2017) emphasizes the importance of meticulous surgical technique and patient preparation in minimizing complications like infection and seroma formation. The authors advocate for a comprehensive understanding of patient history, including lifestyle factors, which can significantly influence surgical outcomes. **Conclusion:** In conclusion, the reviewed literature indicates that complications in brachioplasty are influenced by a combination of surgical techniques, patient characteristics, and psychosocial factors. A nuanced understanding of these elements is essential for improving surgical outcomes and patient satisfaction. Future research should continue to explore the long-term effects of innovative surgical techniques and the psychological implications of body contouring procedures.

Keyword: Complications, Brachioplasty

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The literature surrounding complications in brachioplasty reveals a complex interplay of surgical techniques, patient outcomes, and the psychosocial dimensions of body contouring. In the study by (Freitas-Silva et al., 2010), the authors examined the quality of life and shoulder-arm morbidity among breast cancer survivors undergoing different surgical interventions. Their findings emphasize that while the type of surgery may not directly correlate with arm edema, it does impact the range of motion (ROM) and overall patient satisfaction. This underscores the necessity for future research to delve deeper into how surgical choices influence postoperative complications and patient experiences.

contributes to this discourse by addressing the potential complications associated with conventional brachioplasty, particularly the role of fascial weakness in postoperative healing. The introduction of the SERI surgical scaffold is presented as a promising adjunct to enhance surgical outcomes and minimize scarring. This suggests that innovations in surgical techniques may mitigate some of the complications that arise from traditional methods, offering a pathway for improved patient satisfaction and recovery.

The review by (Vidal et al., 2017) expands the scope of complications by focusing on abdominoplasty, yet it provides insights relevant to brachioplasty. The authors highlight the importance of meticulous surgical technique and patient preparation in reducing complications such as infection and seroma formation. Their findings advocate for a comprehensive understanding of patient history, including lifestyle factors like tobacco use, which can significantly influence surgical outcomes.

In a systematic review, (Toma et al., 2018) explore the psychosocial impacts of body contouring post-bariatric surgery, noting that these procedures can enhance quality of life and body image among patients. This perspective is critical as it connects the physical outcomes of surgeries like brachioplasty with the emotional

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

and psychological well-being of patients, suggesting that complications may also have profound effects on mental health.

(CINTRA et al., 2021) further investigate the complications specific to post-bariatric body contouring surgeries. Their retrospective analysis of 180 patients reveals a significant incidence of postoperative complications, correlating them with patient demographics such as age and BMI. This study highlights the necessity of a tailored approach to patient selection and preoperative assessment to minimize risks associated with body contouring procedures.

Finally, (Sadeghi et al., 2022) provide a contemporary overview of best practices in post-bariatric plastic surgery, emphasizing the importance of thorough preoperative evaluations and considerations of comorbidities. Their insights reinforce the need for a multi-faceted approach to surgical planning that considers both the physical and psychological aspects of patient care.

Together, these articles create a nuanced understanding of the complications associated with brachioplasty and related procedures, revealing the importance of surgical technique, patient selection, and the psychosocial dimensions of body image in shaping patient outcomes.

LITERATURE REVIEW

The article "Comparison of quality of life, satisfaction with surgery and shoulder-arm morbidity in breast cancer survivors submitted to breast-conserving therapy or mastectomy followed by immediate breast reconstruction" by (Freitas-Silva et al., 2010) presents a significant examination of the complications associated with surgical interventions in breast cancer treatment, particularly focusing on shoulder-arm morbidity and patient satisfaction. This cross-sectional study provides valuable insights into the outcomes of two distinct surgical approaches: mastectomy with immediate breast reconstruction (MRM+IBR) and breast-conserving therapy (BCT).

One of the key findings of this study is the lack of difference in the frequency of shoulder-arm morbidity between the MRM+IBR group and the BCT group. This

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

challenges the assumption that more extensive surgical procedures inherently lead to greater complications, such as lymphedema or reduced range of motion (ROM). The authors emphasize that reconstructive surgery was not linked to an increased occurrence of lymphedema, which is a common concern among patients undergoing mastectomy. This finding is particularly relevant for understanding the implications of surgical choices on long-term health outcomes.

However, the study does reveal that reduced ROM, specifically in internal rotation, was more prevalent among women who underwent BCT. This suggests that even less invasive surgical options may carry their own set of complications, which must be carefully considered when discussing surgical options with patients. The authors advocate for further prospective studies with larger sample sizes to more comprehensively investigate the relationship between surgical technique and shoulder-arm function. This highlights a critical area for future research, as understanding the nuances of post-surgical complications can inform clinical practice and improve patient care.

The article also touches upon the complex interplay between surgical type, body image, and patient satisfaction. The authors note that while tumor-related factors are important, the role of patient preference in selecting the type of surgery cannot be overstated. This underscores the necessity for a patient-centered approach in surgical decision-making, where the individual's values and experiences are prioritized.

In the article "SERI Surgical Scaffold as an Adjunct to Conventional Brachioplasty," presents a compelling discussion on the complications associated with brachioplasty, particularly focusing on the role of fascial weakness in the healing process and scar formation. The author posits that the protracted healing observed in brachioplasty scars is largely attributed to underlying fascial support deficiencies, which can lead to cutaneous scar hypertrophy as a compensatory mechanism. This insight is crucial as it highlights the importance of considering the structural integrity of the underlying tissues when evaluating the outcomes of brachioplasty procedures.

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

's evaluation of the SERI surgical scaffold as an adjunct to conventional brachioplasty is particularly noteworthy. The introduction of SERI is proposed as a method to enhance the support for the superficial fascial system (SFS), potentially mitigating the risk of postoperative scar widening. This innovative approach is presented as a response to the common concern of conspicuous scarring, which is often a significant consideration for patients undergoing brachioplasty. The author's clinical experience with a 46-year-old female patient who had long struggled with pendulous upper arms serves as a practical illustration of the application of SERI in this context. The detailed account of the surgical procedure, including the dimensions of the SERI scaffold and the use of Prineo tape as a final dressing, provides valuable insights into the technical aspects of implementing this adjunctive treatment.

However, while offers a promising perspective on the use of SERI, it is important to critically evaluate the long-term implications and effectiveness of this approach. The article does not provide extensive data on the outcomes of the procedure beyond the initial application, which raises questions about the durability of the results and the potential for complications associated with the scaffold itself. Furthermore, the decision-making process regarding the use of SERI in conjunction with conventional brachioplasty necessitates a careful consideration of patient-specific factors, including individual healing responses and the potential for varying degrees of scar formation.

The article titled "Managing Complications in Abdominoplasty: A Literature Review" by (Vidal et al., 2017) provides a comprehensive examination of the complications associated with abdominoplasty, a procedure that shares some similarities with brachioplasty in terms of surgical technique and potential complications. The authors systematically present evidence regarding the incidence, prevention, and management of complications, emphasizing the importance of patient preparation and surgical technique.

One of the key insights from the article is the critical role of preoperative patient preparation, which includes addressing factors such as tobacco cessation. The authors argue that such measures can significantly reduce the risk of complications

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

like infection and skin necrosis, which are relevant concerns in both abdominoplasty and brachioplasty procedures. This preventative approach underscores the necessity of a thorough initial consultation, where the surgeon can assess the patient's health status and readiness for surgery.

The article also highlights the importance of employing a polished surgical technique, informed by both current evidence and the surgeon's experience. This aspect is particularly relevant to brachioplasty, where the meticulous dissection and flap management are crucial to avoid complications such as seroma and necrosis. The authors note that the integration of liposuction into abdominoplasty does not inherently increase risk, provided that adequate perfusion is maintained. This finding could be extrapolated to brachioplasty, suggesting that similar techniques may be safely employed if careful attention is given to flap viability.

In terms of specific complications, the article discusses strategies for seroma prevention, including the use of progressive tension sutures and fibrin sealants, which could be beneficial in brachioplasty as well. The emphasis on dissection over the Scarpa fascia is particularly noteworthy, as it suggests a methodical approach to minimizing fluid accumulation, a common issue in various body contouring surgeries.

While the article presents valuable insights into the management of complications in abdominoplasty, it does not delve deeply into the statistical analysis of the studies reviewed, which could have provided a more robust understanding of the effectiveness of the proposed strategies. Nonetheless, the systematic presentation of evidence and the focus on surgical technique and patient preparation serve as a valuable resource for surgeons engaged in brachioplasty and similar procedures.

The article "Does body contouring after bariatric weight loss enhance quality of life? A systematic review of QOL studies" by (Toma et al., 2018) provides a comprehensive examination of the psychosocial implications of body contouring procedures, particularly in the context of individuals who have undergone bariatric surgery. The authors systematically review various studies that highlight the relationship between body contouring and quality of life (QOL), emphasizing the

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

significant changes in body image and overall satisfaction following such interventions.

One of the key insights from this article is the observed improvement in body image dissatisfaction among obese women after undergoing body contouring surgeries post-bariatric weight loss. The authors note that patients often experience heightened feelings of body image dissatisfaction before undergoing these procedures, which can significantly impact their mental health and social interactions. The systematic review indicates that body contouring not only addresses physical changes but also serves as a crucial intervention for enhancing psychosocial well-being.

The authors also point out the necessity for guidelines on body recontouring after bariatric surgery. This is particularly relevant given the complexities involved in assessing candidates for such procedures. The review highlights that while weight loss rates are higher among individuals who opt for body contouring, the decision to pursue these surgeries should be approached with caution, considering the potential psychological ramifications and the need for comprehensive pre-operative assessments.

Furthermore, the article discusses the implications of body contouring procedures for individuals with body dysmorphic disorder, suggesting that this population may have unique needs and concerns that warrant specialized attention. The authors argue for a nuanced understanding of the motivations behind seeking body contouring, which can vary significantly among individuals.

The article "Does Body Contouring After Bariatric Weight Loss Enhance Quality of Life? A Systematic Review of QOL Studies" by (Toma et al., 2018) provides a comprehensive examination of the psychosocial implications of body contouring procedures, particularly in the context of patients who have undergone bariatric surgery. The authors meticulously analyze the relationship between body image dissatisfaction and the subsequent desire for body contouring, highlighting the significant role that such procedures play in enhancing quality of life (QOL) for these patients.

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

One of the critical insights from the article is the emphasis on body image and its profound impact on the psychological well-being of individuals post-bariatric surgery. The authors argue that while weight loss is a primary goal for many undergoing bariatric procedures, the accompanying changes in body shape and the potential for body image dissatisfaction can lead to a complex interplay of emotions and self-perception. This dissatisfaction is particularly acute among obese women, as noted in the article, where the authors underscore the specificity and severity of body image issues that may arise.

Moreover, the systematic review reveals that patients who pursue body contouring procedures, such as brachioplasty, often report higher levels of satisfaction and an improved quality of life. This finding suggests that body contouring is not merely a cosmetic enhancement but a necessary step for many in achieving psychological and emotional healing after significant weight loss. The authors advocate for the need for clear guidelines regarding body recontouring after bariatric surgery, which indicates a gap in current clinical practices that could be addressed to better support patients' needs.

The article also highlights the importance of addressing body image dissatisfaction as a clinical concern, emphasizing that healthcare providers should consider the psychosocial aspects of body contouring. The authors encourage a more holistic approach to patient care, where the psychological motives behind seeking body contouring are acknowledged and addressed. This perspective is crucial for improving patient outcomes and ensuring that individuals receive comprehensive support throughout their weight loss journey.

The article "Post-bariatric body contouring surgery: analysis of complications in 180 consecutive patients" by (CINTRA et al., 2021) presents a comprehensive analysis of the complications associated with body contouring surgeries following bariatric procedures. This retrospective study examines a sample of 180 patients, providing valuable insights into the interplay between patient demographics—specifically age and Body Mass Index (BMI)—and the incidence of postoperative complications.

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The authors effectively contextualize the necessity for body contouring after significant weight loss, which is often a consequence of bariatric surgery aimed at treating severe obesity. They highlight the psychological and physical motivations behind these surgeries, addressing the body dysmorphia that can ensue from drastic weight changes. The study's design, which includes a thorough collection of preoperative and postoperative data, allows for a nuanced understanding of the complications that may arise, thus contributing to the existing literature on this topic.

One of the key findings of the study is the correlation between age and BMI with the rate of complications. The authors demonstrate that older patients and those with higher BMI prior to surgery exhibit a higher incidence of postoperative complications. This finding is significant, as it suggests that patient selection criteria may need to be refined to mitigate risks. The analysis also underscores the importance of preoperative assessments that consider these factors, as they could guide clinicians in tailoring their surgical approaches and postoperative care to individual patient needs.

Moreover, the study details various types of complications encountered, which range from minor to severe, thus providing a comprehensive overview of the risks associated with body contouring procedures. This serves as a critical resource for both surgeons and patients, as it emphasizes the need for informed consent and realistic expectations regarding surgical outcomes.

However, while the study presents valuable data, it would benefit from a discussion of the long-term outcomes of these surgeries, as well as the psychological impact of complications on patients. Additionally, the retrospective nature of the study may introduce biases in data collection and interpretation, which should be acknowledged when considering the broader applicability of the findings.

The article titled "Post-Bariatric Plastic Surgery: Abdominoplasty, the State of the Art in Body Contouring" by (Sadeghi et al., 2022) provides a comprehensive overview of the complexities involved in post-bariatric body contouring, with particular emphasis on the implications for brachioplasty. The authors highlight that

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

successful outcomes in such surgeries are contingent upon a thorough evaluation of the patient's nutritional status and existing comorbidities. This evaluation is crucial as it informs the surgical approach and potential complications that may arise during or after the procedure.

One of the critical insights from the article is the importance of prophylaxis against thromboembolic events, which is particularly pertinent in patients who have undergone significant weight loss. The authors advocate for a multidisciplinary approach to manage these risks, underscoring that complications such as deep vein thrombosis can significantly impact recovery and overall surgical success. The need for careful planning and consideration of these factors is essential in minimizing complications associated with brachioplasty, as highlighted by the authors.

Furthermore, the article discusses the potential benefits of employing multiple procedures or staged approaches. This strategy not only allows for a more manageable recovery process but also enables the surgical team to address complications as they arise, thereby enhancing patient safety. The authors suggest that such an approach can lead to improved aesthetic outcomes and patient satisfaction, which are vital in the context of post-bariatric surgeries.

CONCLUSION

The literature on complications in brachioplasty and related body contouring procedures reveals a multifaceted understanding of surgical techniques, patient demographics, and psychosocial impacts. The studies reviewed indicate that the choice of surgical method has significant implications for both physical outcomes and patient satisfaction.

In the study by (Freitas-Silva et al., 2010), the authors found that while different surgical interventions in breast cancer survivors did not significantly affect the incidence of arm edema, they did impact the range of motion (ROM) and overall satisfaction, suggesting that surgical choices play a crucial role in postoperative recovery. This finding underscores the need for further research into how specific surgical techniques can influence complications and patient experiences.

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The introduction of the SERI surgical scaffold as an adjunct in brachioplasty, discussed by , highlights innovations aimed at reducing complications such as scar hypertrophy, which are common in conventional methods. This advancement suggests a potential for improved outcomes through enhanced fascial support, although further investigation into long-term effectiveness is warranted.

Moreover, the review by (Vidal et al., 2017) emphasizes the importance of meticulous surgical technique and patient preparation in minimizing complications like infection and seroma formation. The authors advocate for a comprehensive understanding of patient history, including lifestyle factors, which can significantly influence surgical outcomes.

The psychosocial dimensions of body contouring, particularly following bariatric surgery, are explored in (Toma et al., 2018). The study reveals that body contouring can greatly enhance quality of life and body image, suggesting that the ramifications of surgical complications extend beyond physical health to encompass emotional well-being.

Further, (CINTRA et al., 2021) provides a retrospective analysis of complications in post-bariatric body contouring, indicating that variables such as age and BMI are significant predictors of postoperative complications. This highlights the necessity for tailored patient selection and preoperative assessments to minimize risks.

Lastly, (Sadeghi et al., 2022) advocates for a comprehensive approach to post-bariatric plastic surgery, emphasizing the need for thorough preoperative evaluations and consideration of comorbidities. The authors suggest that adopting a multidisciplinary approach can help manage potential complications effectively.

In conclusion, the reviewed literature indicates that complications in brachioplasty are influenced by a combination of surgical techniques, patient characteristics, and psychosocial factors. A nuanced understanding of these elements is essential for improving surgical outcomes and patient satisfaction. Future research should continue to explore the long-term effects of innovative surgical techniques and the psychological implications of body contouring procedures.

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Freitas-Silva, R., Marques Conde, D., de Freitas-Júnior, R., & Zangiacomi Martinez, E., 2010. Comparison of quality of life, satisfaction with surgery and shoulder-arm morbidity in breast cancer survivors submitted to breast-conserving therapy or mastectomy followed by immediate breast reconstruction. ncbi.nlm.nih.gov
2. Vidal, P., Enrique Berner, J., & A. Will, P., 2017. Managing Complications in Abdominoplasty: A Literature Review. ncbi.nlm.nih.gov
3. Toma, T., Harling, L., Athanasiou, T., Darzi, A., & Ashrafian, H., 2018. Does body contouring after bariatric weight loss enhance quality of life? A systematic review of QOL studies. [PDF]
4. CINTRA, W. I. L. S. O. N., LUIZ ANTONIO MODOLIN, M. I. G. U. E. L., RICARDO COLFERAI, D. I. E. G. O., ITOCAZO ROCHA, R. O. D. R. I. G. O., & GEMPERLI, R. O. L. F., 2021. Post-bariatric body contouring surgery: analysis of complications in 180 consecutive patients. ncbi.nlm.nih.gov
5. Sadeghi, P., Duarte-Bateman, D., Ma, W., Khalaf, R., Fodor, R., Pieretti, G., Ciccarelli, F., Harandi, H., & Cuomo, R., 2022. Post-Bariatric Plastic Surgery: Abdominoplasty, the State of the Art in Body Contouring. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Plastic Surgery And Aesthetic Medicine
Copyright © International Medical Journal Corp. Ltd. All rights reserved