Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Radiology and Oncology Radiation



Diagnostic Modalities for Acute Compartment Syndrome of the Extremities

[1] Annisa, [1] Eka

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.annisa9384829@gmail.com

Article History :

Received date : 2024/07/16
Revised date : 2024/08/14
Accepted date : 2024/09/08
Published date : 2025/01/10

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: Acute compartment syndrome (ACS) is a critical condition characterized by increased pressure within a closed fascial compartment, leading to compromised blood flow and potential tissue necrosis. **Literature Review:** The complexities surrounding the diagnosis of ACS are addressed by (Guo et al., 2019), who emphasize the reliance on clinical assessments and the limitations of subjective criteria, such as the "5Ps": pain, pallor, paresthesias, paralysis, and pulselessness. They caution that these indicators can lead to unnecessary surgical interventions when not supported by objective pressure measurements. This concern is echoed by (Aleisawi et al., 2021), who assert that pain disproportionate to the injury is the most sensitive indicator of impending ACS, calling for improved diagnostic accuracy. The integration of intracompartmental pressure (ICP) measurements is recognized as essential, as discussed by (Shiraz et al., 2022), who advocate for combining clinical judgment with objective measures to enhance diagnostic reliability. The limitations of imaging modalities are also noted, with (A. Al-Khafaji & Riemer, 2022) highlighting that while ultrasound can rule out other conditions, it is not definitive for ACS diagnosis, and MRI, despite its superior diagnostic capability, can introduce delays in treatment. Emerging technologies, such as continuous compartment pressure monitoring, are proposed by (Schupbach et al., 2024) to improve the timeliness of surgical intervention, addressing the shortcomings of traditional diagnostic approaches. This aligns with the comprehensive review by (L. Davidson et al., 2024), which emphasizes the need for evidence-based practices that integrate both clinical assessments and objective measurements. **Conclusion:** In conclusion, the literature reflects a consensus on the necessity for improved diagnostic strategies that combine clinical acumen with objective measures, particularly ICP monitoring, to enhance the identification and management of ACS. The evolution of thought surrounding ACS diagnosis underscores the critical importance of timely intervention to mitigate severe complications, thus emphasizing the need for ongoing advancements in diagnostic modalities and clinical practices.

Keyword: Diagnostic Modalities, Acute Compartment Syndrome of the Extremities

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

Acute compartment syndrome (ACS) is a critical condition characterized by increased pressure within a closed fascial compartment, leading to compromised blood flow and potential tissue necrosis. The literature on ACS has evolved significantly over the past decade, focusing on the nuances of diagnosis and management. The review begins with the foundational insights provided by (Tapiwa Mabvuure et al., 2012), who emphasize the urgency of early diagnosis and intervention. Their work highlights that timely fasciotomy can markedly improve patient outcomes and reduce the risk of complications, underscoring the need for clinicians to remain vigilant in identifying this surgical emergency.

(Guo et al., 2019) further elaborate on the complexities of diagnosing ACS, noting the reliance on clinical assessments and the controversial nature of the diagnostic criteria. They introduce the "5Ps" (pain, pallor, paresthesias, paralysis, and pulselessness) as critical indicators, but caution that subjective clinical evaluations may lead to unnecessary surgical interventions. The authors also point out that many medical facilities lack the necessary equipment for pressure measurement, which is essential for accurate diagnosis, thus highlighting a significant gap in clinical practice.

(Aleisawi et al., 2021) reinforce the notion that traditional clinical signs are poor predictors of ACS, advocating for a more nuanced understanding of the condition. They assert that pain that is disproportionate to the injury is the most sensitive indicator of impending ACS, which echoes the findings of (Guo et al., 2019) regarding the inadequacy of standard clinical assessments. This perspective calls for a more rigorous approach to evaluation and diagnostic accuracy.

In a case study, (Shiraz et al., 2022) discuss the diagnostic challenges faced in acute compartment syndrome, particularly when no clear etiology is present. They emphasize that while clinical judgment plays a pivotal role, it alone is insufficient

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

for reliable diagnosis, thus necessitating the integration of intracompartmental pressure measurements and imaging techniques. This sentiment is echoed in (A. Al-Khafaji & Riemer, 2022)'s work, which discusses the limitations of imaging modalities in diagnosing ACS. They highlight that while ultrasound can provide valuable insights into muscle trauma and exclude other conditions, it is not a definitive diagnostic tool and can introduce delays in treatment.

(Schupbach et al., 2024) introduce a novel approach to diagnosing ACS through continuous compartment pressure monitoring, which has shown promise in improving the timeliness of surgical intervention. They acknowledge the subjective nature of clinical findings and the mixed results associated with standard pressure measurements, advocating for new technologies that may enhance diagnostic accuracy.

Finally, (L. Davidson et al., 2024) provide a comprehensive review of contemporary practices in diagnosing and managing ACS. They reiterate the importance of early identification to prevent irreversible damage, while also discussing the predominance of traumatic causes and the demographic disparities in incidence rates. Their review emphasizes the need for evidence-based practices and the integration of both clinical assessment and objective pressure measurements in the diagnostic process.

In summary, the literature on acute compartment syndrome reflects a growing recognition of the need for improved diagnostic strategies that combine clinical acumen with objective measures. The evolution of thought regarding the management of ACS underscores the critical importance of timely intervention to mitigate the risk of severe complications.

LITERATURE REVIEW

The article "Acute Compartment Syndrome of the Limbs: Current Concepts and Management" by (Tapiwa Mabvuure et al., 2012) provides a comprehensive overview of acute compartment syndrome (ACS), focusing on its aetiology,

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

pathophysiology, diagnosis, and management strategies. The authors underscore the critical nature of early diagnosis and intervention, which are paramount in improving patient outcomes and minimizing the risk of complications, including the potential for legal indemnity.

A key aspect of the article is the discussion of the various diagnostic modalities available for measuring intracompartmental pressures, which is essential for confirming the diagnosis of ACS. The authors detail both invasive and non-invasive techniques, highlighting that while direct measurement through a manometer is the gold standard, alternative methods such as the use of a handheld pressure monitor can also provide valuable insights. This critical evaluation of diagnostic techniques illustrates the complexity of accurately diagnosing ACS, as the condition can often be mistaken for other injuries or conditions.

Furthermore, the article emphasizes the pathophysiological mechanisms underlying ACS, which occur when increased pressure within a closed fascial compartment compromises circulation and function of the tissues within that compartment. The authors elucidate how timely recognition of the signs and symptoms, such as pain out of proportion to injury and sensory deficits, is crucial for effective management. They argue that a high index of suspicion should be maintained, especially in at-risk populations, such as those with recent fractures or significant soft tissue injuries.

Management strategies discussed in the article center around the necessity of early fasciotomy as the definitive treatment for ACS. The authors contend that prompt surgical intervention not only alleviates pressure but also significantly improves the prognosis for limb function and reduces the risk of long-term complications. The review provides a compelling argument for the integration of clinical judgment with objective pressure measurements to guide treatment decisions.

The article "Acute compartment syndrome: Cause, diagnosis, and new viewpoint" by (Guo et al., 2019) provides a comprehensive overview of the diagnostic modalities for acute compartment syndrome (ACS) of the extremities, emphasizing

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

the need for accurate assessment to prevent serious complications. The authors highlight that the diagnosis of ACS primarily relies on clinical evaluation and compartment pressure measurement, which are crucial for timely intervention.

The article delineates the clinical hallmarks of compartment syndrome, succinctly summarized as the "5Ps": pain out of proportion, pallor, paresthesias, paralysis, and pulselessness. These indicators are critical as their presence often suggests the onset of muscle necrosis, marking a critical stage of ACS. However, the authors critically evaluate the reliability of subjective clinical assessments, noting that they can lead to unnecessary fasciotomies, which carry significant risks and complications. This point underscores the necessity for objective measurement techniques in clinical practice.

(Guo et al., 2019) also discuss the limitations of current diagnostic tools. While invasive monitoring methods such as the Whitesides needle manometer and slit catheter are available, they are prone to complications such as blockage by muscle or blood clots, which can hinder accurate pressure readings. Furthermore, the STIC Monitor is introduced as a more straightforward option for measuring compartment pressure; however, the authors caution that it may yield misleading results due to potential technical variations. This critical evaluation of diagnostic modalities reflects a nuanced understanding of the challenges faced in diagnosing ACS, indicating that reliance on a single method may not suffice for accurate diagnosis.

The article by (Aleisawi et al., 2021) provides a detailed exploration of acute compartment syndrome (ACS), emphasizing its definition and the critical role of interstitial pressure within closed fascial compartments. The authors articulate that ACS leads to microvascular compromise, which is a significant concern for clinicians when assessing limb viability. They highlight the classic signs associated with ACS, including tense swelling, pallor, paresthesia, paralysis, and pulselessness. However, the authors critically note that these signs are often poor predictors of the condition, which underscores the complexity of diagnosing ACS in clinical practice.

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

A pivotal aspect of the article is the identification of pain as the most sensitive and reliable indicator of ACS. The authors emphasize that in patients who are neurologically responsive and sensate, pain that is disproportionate to the injury and unresponsive to analgesics serves as an early warning sign for potential compartment syndrome. This insight is particularly valuable for clinicians, as it shifts the focus from traditional signs that may be misleading to a more nuanced understanding of pain as a diagnostic modality.

The case reports included in the article serve to illustrate the challenges faced in diagnosing ACS accurately. The authors effectively use these cases to demonstrate how reliance on classic signs can lead to misdiagnosis or delayed treatment, which can have dire consequences for limb function. This practical application of theoretical knowledge enhances the article's relevance and provides a real-world context for the discussion.

The article "Unidentified Recurrent Acute Compartment Syndrome of the Right Upper Limb" by (Shiraz et al., 2022) provides a comprehensive examination of acute compartment syndrome (ACS), emphasizing its nature as a surgical emergency characterized by elevated pressure within a closed osteofascial compartment. The authors meticulously outline the consequences of untreated ACS, which can include ischemia, necrosis, infection, and potential amputation, highlighting the critical need for timely diagnosis and intervention.

A significant contribution of this article is its discussion on the diagnostic modalities employed for ACS. While the authors acknowledge that ACS is primarily diagnosed through clinical judgment, they underscore the limitations of this approach, particularly its inadequate diagnostic performance characteristics. This recognition is crucial, as it points to the necessity for supplementary diagnostic measures to enhance the accuracy of ACS detection. The article suggests that intracompartmental pressure (ICP) measurements and radiological scans can serve as valuable adjuncts to clinical evaluation, providing quantitative data that may aid in confirming a diagnosis of ACS.

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article also notes the common sites for ACS, primarily the anterior compartment of the leg, but extends the discussion to other anatomical locations such as the forearm, thigh, buttock, shoulder, hand, and foot. This broad perspective is beneficial for clinicians as it reinforces the concept that ACS can occur in various contexts, including those with no identifiable underlying etiology. The authors' inclusion of cases with unidentified recurrent ACS further emphasizes the complexity of the condition and the potential challenges in diagnosis.

However, while the article effectively highlights the importance of combining clinical assessment with ICP and imaging techniques, it could benefit from a more detailed exploration of the specific thresholds for ICP that indicate ACS and the types of radiological scans that are most effective in diagnosing the condition. Furthermore, the discussion could be enriched by including recent advancements in imaging technology or diagnostic algorithms that may enhance the identification of ACS.

The article titled "Spontaneous acute forearm compartment syndrome: Case report of a clinical diagnosis with limited imaging options" by (A. Al-Khafaji & Riemer, 2022) provides an insightful examination of acute compartment syndrome (ACS) and the challenges associated with its diagnosis. The authors emphasize that ACS is a surgical emergency characterized by elevated intracompartmental pressure, with a critical threshold set at > 30 mmHg, despite the accepted normal pressure being < 10 mmHg. This detail underscores the importance of understanding pressure dynamics within muscle compartments as a fundamental aspect of diagnosing ACS.

The article highlights that fractures account for a significant majority (75%) of the predisposing factors for ACS, while other causes include penetrating traumas, burns, vascular injuries, bleeding disorders, and intense athletic activities. This broad range of potential causes illustrates the complexity of ACS and the necessity for a thorough clinical assessment to ascertain the underlying etiology. The authors assert that the gold standard for diagnosing ACS relies heavily on clinical

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

symptomatology, which can often be ambiguous and lead to delays in diagnosis and subsequent treatment.

The discussion on imaging modalities is particularly noteworthy. The authors point out that while ultrasound (US) is a more accessible option for dynamic imaging of muscle trauma, its role in ACS diagnosis is limited. US can reveal signs of edema, indicated by increased echogenicity, but this finding is not specific to compartment syndrome. Furthermore, US can effectively rule out other conditions such as thrombus, muscle tears, and vascular occlusions, which may mimic ACS symptoms.

In contrast, magnetic resonance imaging (MRI) is highlighted as superior for diagnosing soft tissue pathologies, including those relevant to ACS. However, the authors caution that the time required to obtain an MRI can lead to delays in critical surgical intervention, which is vital in cases of ACS. This tension between the need for accurate diagnosis and the urgency of treatment is a significant concern, as delays can result in irreversible muscle damage.

The article also touches on the implications of delayed diagnoses, which can stem from unclear presentations of the condition or delays in patients seeking medical care. This aspect is crucial, as it emphasizes the need for heightened awareness and education regarding the signs and symptoms of ACS among both healthcare providers and patients.

The article "Continuous Compartment Pressure Monitoring Allows the Early Detection of Compartment Syndrome After Arterial Revascularization" by (Schupbach et al., 2024) provides a critical examination of the diagnostic modalities for acute compartment syndrome (CS), particularly in the context of post-arterial revascularization. The authors emphasize that while the diagnosis of acute CS is traditionally based on clinical findings, these indicators are often debated and may not consistently lead to timely intervention.

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article outlines the classical symptoms of CS, including disproportionate pain, pain on passive muscle stretch, hypoaesthesia, weakness, swelling, and tight compartments. These clinical signs, while essential, may not always be reliable, leading to potential delays in diagnosis and treatment. The authors argue that continuous compartment pressure monitoring can significantly enhance the diagnostic process. They present evidence suggesting that this method reduces the time to surgical intervention, which is crucial given that the gold standard treatment for acute CS is an emergency fasciotomy aimed at alleviating the elevated pressures within the compartment.

Moreover, the article highlights concerning statistics regarding the outcomes of fasciotomies, noting that nearly 17% of such procedures reveal some degree of myonecrosis, and 5.4% result in amputations. These figures underscore the inadequacy of current diagnostic techniques, which may lead to either delayed treatment or unnecessary surgical interventions. The authors advocate for the integration of advanced monitoring systems, specifically a micro-electrical machine systems (MEMS)-based pressure monitor, which has shown promising results in trauma patients. This technological advancement could potentially address the shortcomings of traditional diagnostic methods by providing real-time pressure data, thus facilitating earlier and more accurate diagnosis of CS.

The article "Practical Review on the Contemporary Diagnosis and Management of Compartment Syndrome" by (L. Davidson et al., 2024) provides a thorough examination of acute compartment syndrome (ACS), emphasizing the critical nature of timely diagnosis and intervention to mitigate severe morbidity associated with this condition. The authors effectively highlight that ACS is predominantly a surgical emergency resulting from elevated intracompartmental pressures, which can surpass arterial perfusion pressure, leading to ischemia and subsequent complications such as irreversible myonecrosis and nerve damage.

The article underscores that traumatic injuries account for approximately 75% of ACS cases, with the lower leg and forearm being the most frequently affected areas.

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

This statistic is particularly relevant as it points to the necessity for heightened awareness among healthcare providers, especially in emergency settings, to recognize the signs and symptoms of ACS in patients presenting with trauma. The authors also note a significant disparity in incidence rates between genders, attributing the higher prevalence in men to a combination of increased trauma exposure and a greater proportion of muscle mass relative to overall tissue, which may contribute to the development of ACS.

A critical aspect of the review is the discussion on diagnostic modalities. While the authors acknowledge that direct measurement devices can objectively identify compartment pressures, they assert that clinical diagnosis remains the gold standard. This assertion is particularly important as it emphasizes the need for clinicians to rely on a comprehensive clinical assessment, including the evaluation of symptoms such as pain out of proportion to injury, sensory deficits, and motor weakness, which are essential for timely diagnosis.

The review effectively synthesizes current evidence-based practices for the recognition and management of ACS, providing a clear framework for clinicians. However, it could benefit from a deeper exploration of the limitations of current diagnostic tools and the potential for advancements in technology to enhance diagnostic accuracy. Additionally, while the authors provide a robust overview of the condition, further discussion on post-operative management and rehabilitation following surgical intervention could enrich the understanding of long-term outcomes for patients with ACS.

CONCLUSION

The literature on acute compartment syndrome (ACS) has evolved to emphasize the critical nature of timely diagnosis and intervention, which are paramount for improving patient outcomes and minimizing complications. The foundational work by (Tapiwa Mabvuure et al., 2012) highlights the urgency of recognizing ACS as a surgical emergency, advocating for early fasciotomy to alleviate pressure and prevent irreversible tissue damage.

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The complexities surrounding the diagnosis of ACS are addressed by (Guo et al., 2019), who emphasize the reliance on clinical assessments and the limitations of subjective criteria, such as the "5Ps": pain, pallor, paresthesias, paralysis, and pulselessness. They caution that these indicators can lead to unnecessary surgical interventions when not supported by objective pressure measurements. This concern is echoed by (Aleisawi et al., 2021), who assert that pain disproportionate to the injury is the most sensitive indicator of impending ACS, calling for improved diagnostic accuracy.

The integration of intracompartmental pressure (ICP) measurements is recognized as essential, as discussed by (Shiraz et al., 2022), who advocate for combining clinical judgment with objective measures to enhance diagnostic reliability. The limitations of imaging modalities are also noted, with (A. Al-Khafaji & Riemer, 2022) highlighting that while ultrasound can rule out other conditions, it is not definitive for ACS diagnosis, and MRI, despite its superior diagnostic capability, can introduce delays in treatment.

Emerging technologies, such as continuous compartment pressure monitoring, are proposed by (Schupbach et al., 2024) to improve the timeliness of surgical intervention, addressing the shortcomings of traditional diagnostic approaches. This aligns with the comprehensive review by (L. Davidson et al., 2024), which emphasizes the need for evidence-based practices that integrate both clinical assessments and objective measurements.

In conclusion, the literature reflects a consensus on the necessity for improved diagnostic strategies that combine clinical acumen with objective measures, particularly ICP monitoring, to enhance the identification and management of ACS. The evolution of thought surrounding ACS diagnosis underscores the critical importance of timely intervention to mitigate severe complications, thus emphasizing the need for ongoing advancements in diagnostic modalities and clinical practices.

DISCLOSURE STATEMENT

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Tapiwa Mabvuure, N., Malahias, M., Hindocha, S., Khan, W., & Juma, A., 2012. Acute Compartment Syndrome of the Limbs: Current Concepts and Management. ncbi.nlm.nih.gov
2. Guo, J., Yin, Y., Jin, L., Zhang, R., Hou, Z., & Zhang, Y., 2019. Acute compartment syndrome: Cause, diagnosis, and new viewpoint. ncbi.nlm.nih.gov
3. Aleisawi, H., Alsager, G., Pasha, A., Alyousif, H., Alsarhan, H., & Surur, S., 2021. Is It Compartment Syndrome? Two Case Reports and Literature Review. ncbi.nlm.nih.gov
4. Shiraz, A., Bakhtiar, H., Farman, G., Khan, S., & Rafiq, N., 2022. Unidentified Recurrent Acute Compartment Syndrome of the Right Upper Limb. ncbi.nlm.nih.gov
5. Al-Khafaji, R. & Riemer, K., 2022. Spontaneous acute forearm compartment syndrome: Case report of a clinical diagnosis with limited imaging options. ncbi.nlm.nih.gov
6. Schupbach, D., Reindl, R., L Gill, H., S Liberman, A., & J Harvey, E., 2024. Continuous Compartment Pressure Monitoring Allows the Early Detection of Compartment Syndrome After Arterial Revascularization. ncbi.nlm.nih.gov
7. L. Davidson, A., A. Sutherland, M., C. Siska, R., & E. Janis, J., 2024. Practical Review on the Contemporary Diagnosis and Management of Compartment Syndrome. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Radiology and Oncology Radiation
Copyright © International Medical Journal Corp. Ltd. All rights reserved