Research Article Volume 01, Issue No. 02. 2025

The Medical Journal of Surgery and Anaesthesiology



Implications of Preoperative Depression for Lumbar Spine Surgery Outcomes

[1] Yulia, [1] Jessica

[1] Faculty of Medicine, Maranatha Christian University, Bandung City, West Java, Indonesia

Correspondence : dr.yulia38447@gmail.com

Article History :

Received date : 2024/07/12
Revised date : 2024/08/25
Accepted date : 2024/09/09
Published date : 2025/01/10

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature review reveals a significant connection between preoperative depression and the outcomes of lumbar spine surgery, emphasizing the necessity of integrating mental health evaluations into surgical planning. **Literature Review:** Further expanding on this topic, (L. Jackson et al., 2019) emphasizes the importance of validated screening tools for identifying patients at risk due to psychological comorbidities. Their findings indicate that individuals with pre-existing psychological disorders report higher incidences of pain post-surgery, reinforcing the idea that addressing mental health issues prior to surgery can lead to improved surgical outcomes. The bidirectional relationship between chronic pain and mood disorders is particularly noteworthy, as it suggests that effective pain management can positively influence mental health. The systematic review and meta-analysis conducted by (Ye et al., 2024) further supports these findings, demonstrating a significant association between preoperative mental disorders and adverse postoperative outcomes. The reciprocal relationship between postoperative pain and mental health underscores the complexity of managing patients undergoing lumbar spine surgery and highlights the need for a comprehensive approach to patient care. **Conclusion:** In conclusion, the reviewed literature collectively advocates for a multidisciplinary approach that prioritizes mental health alongside physical health in the context of lumbar spine surgery. By incorporating mental health assessments into preoperative evaluations, healthcare providers can identify patients at risk and implement timely interventions, ultimately leading to improved surgical outcomes and enhanced quality of life for patients.

Keyword: Implications, Preoperative Depression, Lumbar Spine Surgery Outcomes

Downloaded from The Medical Journal of Surgery and Anaesthesiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The implications of preoperative depression for lumbar spine surgery outcomes have garnered increasing attention in the medical literature, highlighting the complex interplay between mental health and surgical success. The foundational study by (Sinikallio et al., 2010) establishes a critical link between depressive symptoms during the rehabilitation period and poor outcomes following lumbar spinal stenosis surgery. This research underscores the importance of assessing both preoperative and postoperative depressive symptoms, suggesting that their predictive value can significantly inform surgical outcomes over a two-year perspective. The findings indicate that patients with elevated depressive symptoms may experience less favorable surgical results, prompting a need for early identification and intervention.

Building on this foundation, (L. Jackson et al., 2019) expand the discourse by introducing validated screening tools designed to identify patients at risk due to psychological comorbidities. Their study emphasizes that individuals with pre-existing psychological disorders, particularly anxiety and depression, report higher incidences of acute and chronic pain post-surgery. This correlation suggests that addressing these psychological issues before surgery could substantially improve surgical outcomes. Furthermore, they note a bidirectional relationship between chronic pain and mood disorders, indicating that effective management of pain can lead to improvements in mental health, thereby enhancing overall recovery.

Most recently, (Ye et al., 2024) conducted a systematic review and meta-analysis that reinforces the findings of earlier studies, specifically within the context of scoliosis surgery. Their work highlights a significant association between preoperative mental disorders, including depression and anxiety, and adverse postoperative outcomes such as increased pain, disability, and reduced quality of life. The analysis reveals a reciprocal relationship where postoperative pain exacerbates mental health issues, reinforcing the necessity for a comprehensive

Downloaded from The Medical Journal of Surgery and Anaesthesiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

approach to patient care that integrates mental health considerations. This holistic perspective calls for proactive engagement from both healthcare providers and patients in addressing mental health concerns before and after surgical interventions.

Together, these studies illustrate the profound impact of preoperative depression on surgical outcomes in lumbar spine procedures, advocating for a multidisciplinary approach that prioritizes mental health alongside physical health in surgical planning and recovery.

LITERATURE REVIEW

The article "Depressive symptoms during rehabilitation period predict poor outcome of lumbar spinal stenosis surgery: A two-year perspective" by (Sinikallio et al., 2010) provides a comprehensive analysis of how depressive symptoms can influence surgical outcomes for patients undergoing lumbar spine surgery, specifically in the context of lumbar spinal stenosis (LSS). The authors conduct a prospective clinical study that assesses both preoperative and postoperative depressive symptoms and their correlation with surgical outcomes over a two-year period.

The study's main finding is that preoperative depressive symptoms significantly predict poorer outcomes following LSS surgery. This aligns with existing literature that underscores the importance of mental health in the context of physical health interventions. The authors detail their methodology, which involves a longitudinal assessment of patients, allowing for a nuanced understanding of how depressive symptoms evolve throughout the rehabilitation process. This approach not only highlights the immediate impact of preoperative mental health on surgical success but also emphasizes the importance of ongoing psychological assessment during recovery.

One critical evaluation of the material is the emphasis placed on the predictive value of depressive symptoms, which suggests that addressing mental health prior to

Downloaded from The Medical Journal of Surgery and Anaesthesiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

surgery could be crucial in optimizing surgical outcomes. The authors argue that recognition and treatment of depressive symptoms in patients could lead to improved recovery trajectories and overall satisfaction with surgical results. This point is particularly relevant given the increasing acknowledgment of the biopsychosocial model in healthcare, which integrates psychological factors into the management of physical health conditions.

However, while the study effectively demonstrates a correlation between depressive symptoms and surgical outcomes, it could benefit from a more detailed exploration of the mechanisms underlying this relationship. For instance, the article could delve deeper into how depression may affect pain perception, adherence to rehabilitation protocols, or even the physiological responses to surgical interventions. Understanding these mechanisms could provide healthcare professionals with better tools to address mental health issues in surgical candidates.

Furthermore, the study's reliance on self-reported measures of depressive symptoms may introduce bias, as patients' perceptions of their mental health could be influenced by their physical pain or expectations of surgery. Future research could incorporate more objective measures of depression and explore the role of multidisciplinary approaches that include psychological support as part of preoperative care.

The article "Correlating Psychological Comorbidities and Outcomes After Spine Surgery" by (L. Jackson et al., 2019) presents a comprehensive examination of the relationship between psychological disorders, particularly anxiety and depression, and surgical outcomes in patients undergoing lumbar spine surgery. The authors emphasize the importance of early identification and treatment of psychological comorbidities, which can significantly influence both objective and subjective surgical outcomes.

The study highlights the development and validation of objective screening tools designed to identify patients at risk of experiencing suboptimal surgical outcomes

Downloaded from The Medical Journal of Surgery and Anaesthesiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

due to underlying psychological conditions. This proactive approach is essential, as patients with psychological disorders are shown to report higher rates of acute pain and have an increased likelihood of developing chronic pain compared to those without such disorders. The findings underscore the critical role that psychological health plays in the pain experience and recovery process following surgery.

Moreover, the article elucidates the bidirectional relationship between chronic pain and psychological disorders. Patients suffering from chronic pain not only face the risk of exacerbating their psychological conditions but may also experience improvements in mood disorders following successful surgical interventions. This phenomenon suggests that addressing psychological health prior to surgery could potentially enhance recovery outcomes and overall quality of life for patients.

The authors also delve into the implications of psychological comorbidities on postoperative complications and chronic narcotic use. Their analysis indicates that individuals with anxiety or depression may have a heightened risk for adverse surgical outcomes, which can lead to prolonged recovery times and increased dependence on pain management strategies. This highlights the necessity for healthcare providers to incorporate psychological assessments as a standard part of preoperative evaluations for lumbar spine surgery candidates.

The article titled "The Association Between Mental Disorders and Postoperative Outcomes of Scoliosis Surgery: A Systematic Review and Meta-analysis" by (Ye et al., 2024) presents a thorough examination of the implications of preoperative mental disorders, particularly depression and anxiety, on postoperative outcomes in scoliosis surgery. This systematic review and meta-analysis provide critical insights into how these mental health conditions can adversely affect surgical results, including increased pain, disability, and a decline in quality of life, which are particularly relevant for understanding outcomes in lumbar spine surgery.

The authors systematically synthesized data from multiple studies, highlighting a significant correlation between preoperative mental health status and postoperative experiences. They found that patients with preoperative depression and anxiety

Downloaded from The Medical Journal of Surgery and Anaesthesiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

reported worse outcomes post-surgery, including heightened pain levels and decreased satisfaction with surgical results. This finding is particularly important as it emphasizes the need for healthcare professionals to recognize and address mental health issues prior to surgical intervention. The article effectively argues that a holistic approach to patient care—one that encompasses both physical and mental health—is essential for improving surgical outcomes.

Moreover, the article delineates a reciprocal relationship between postoperative pain and mental health, suggesting that not only do preoperative mental disorders lead to poorer surgical outcomes, but also that the experience of postoperative pain can exacerbate existing mental health issues. This bidirectional relationship underscores the complexity of managing patients undergoing lumbar spine surgery and the necessity for integrated care strategies that include mental health assessments and interventions.

The implications for clinical practice are profound. The authors advocate for the incorporation of mental health evaluations into preoperative assessments, which could lead to timely interventions for those identified as having mental health concerns. This proactive approach may not only improve postoperative outcomes but also enhance patients' overall quality of life. Furthermore, the article encourages patients to engage more actively in their mental health care, thereby fostering a collaborative environment between healthcare providers and patients.

CONCLUSION

The literature review reveals a significant connection between preoperative depression and the outcomes of lumbar spine surgery, emphasizing the necessity of integrating mental health evaluations into surgical planning. The foundational study by (Sinikallio et al., 2010) establishes a critical link between depressive symptoms and poor surgical outcomes, highlighting the predictive value of these symptoms over a two-year period. This study suggests that early identification and intervention for depressive symptoms can enhance recovery trajectories and overall satisfaction with surgical results.

Downloaded from The Medical Journal of Surgery and Anaesthesiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Further expanding on this topic, (L. Jackson et al., 2019) emphasizes the importance of validated screening tools for identifying patients at risk due to psychological comorbidities. Their findings indicate that individuals with pre-existing psychological disorders report higher incidences of pain post-surgery, reinforcing the idea that addressing mental health issues prior to surgery can lead to improved surgical outcomes. The bidirectional relationship between chronic pain and mood disorders is particularly noteworthy, as it suggests that effective pain management can positively influence mental health.

The systematic review and meta-analysis conducted by (Ye et al., 2024) further supports these findings, demonstrating a significant association between preoperative mental disorders and adverse postoperative outcomes. The reciprocal relationship between postoperative pain and mental health underscores the complexity of managing patients undergoing lumbar spine surgery and highlights the need for a comprehensive approach to patient care.

In conclusion, the reviewed literature collectively advocates for a multidisciplinary approach that prioritizes mental health alongside physical health in the context of lumbar spine surgery. By incorporating mental health assessments into preoperative evaluations, healthcare providers can identify patients at risk and implement timely interventions, ultimately leading to improved surgical outcomes and enhanced quality of life for patients.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

Downloaded from The Medical Journal of Surgery and Anaesthesiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved

1. Sinikallio, S., M Lehto, S., Aalto, T., Airaksinen, O., Kröger, H., & Viinamäki, H., 2010. Depressive symptoms during rehabilitation period predict poor outcome of lumbar spinal stenosis surgery: A two-year perspective. ncbi.nlm.nih.gov
2. L. Jackson, K., Rumley, J., Griffith, M., Agochukwu, U., & DeVine, J., 2019. Correlating Psychological Comorbidities and Outcomes After Spine Surgery. ncbi.nlm.nih.gov
3. Ye, L., Lin, S., Lv, Y., Ge, C., & Chen, X., 2024. The Association Between Mental Disorders and Postoperative Outcomes of Scoliosis Surgery: A Systematic Review and Meta-analysis. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Surgery and Anaesthesiology
Copyright © International Medical Journal Corp. Ltd. All rights reserved