Research Article Volume 01, Issue No. 02. 2025

Medical Journal of Urology and General Surgery



Association of Obesity With Septic Complications After Major Abdominal Surgery

[1] Arief, [1] Trisna

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.arief293848@gmail.com

Article History :

Received date : 2024/10/10
Revised date : 2024/11/22
Accepted date : 2024/12/14
Published date : 2025/01/10

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature consistently illustrates that obesity is a significant risk factor for adverse surgical outcomes, particularly following major abdominal surgeries. The introduction emphasizes the multifaceted implications of increased body mass index (BMI) on surgical recovery and complication rates, highlighting a strong correlation between obesity and increased postoperative complications such as wound infections and mortality. **Literature Review:** The systematic review by (S. Almasaudi et al., 2018) further elucidates this relationship, indicating that obesity is linked to higher rates of surgical site infections and prolonged hospital stays in colorectal cancer surgeries. The authors advocate for regular monitoring of obese patients post-surgery due to their unique physiological characteristics that can lead to poorer operative outcomes. This sentiment is echoed in the findings of (John Wilson et al., 2018), which specifically address the heightened risk of deep infections in obese patients undergoing total knee arthroplasty, reinforcing the critical nature of obesity as a risk factor across various surgical contexts. Moreover, the article by (KARANTH & L. KARANTH, 2021) highlights the challenges faced by obese patients in the South Indian population, noting increased surgery durations and postoperative pain, which necessitate tailored preoperative assessments and postoperative monitoring. This need for individualized care is further emphasized in the retrospective study by (Tefera Kassahun et al., 2022), which underscores the significant complications and mortality rates associated with obesity in emergency laparotomy procedures. **Conclusion:** In conclusion, the body of literature reviewed indicates a clear and consistent pattern that obesity is a critical factor influencing surgical outcomes. The interplay between obesity, comorbidities, and surgical complications necessitates a nuanced understanding and proactive management strategies to optimize patient care and surgical success. Addressing obesity through preoperative weight management and tailored postoperative care is essential in improving surgical outcomes for this high-risk population.

Keyword: Obesity, Septic Complications, Major Abdominal Surgery

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The association between obesity and postoperative complications has garnered significant attention in recent years, particularly in the context of major abdominal surgeries. The literature reveals a consistent pattern indicating that obesity is a critical factor influencing surgical outcomes, with various studies highlighting the multifaceted implications of increased body mass index (BMI) on patient recovery and complication rates.

In 2014, (N. Pierpont et al., 2014) provided a comprehensive review emphasizing that obesity is linked to numerous comorbidities which adversely affect surgical outcomes. Their findings indicate that even modest weight loss can significantly decrease morbidity and mortality associated with various health conditions, including those pertinent to surgical recovery. The authors specifically noted that higher BMI correlates with increased rates of surgical complications, such as wound infections and mortality, underscoring the need for careful consideration of obesity in surgical settings.

Building on this foundation, (John Wilson et al., 2018) explored the specific risks associated with obesity in the context of total knee arthroplasty (TKA) patients. They highlighted that obese patients are at a heightened risk for deep infections post-surgery, reinforcing the notion that obesity not only complicates surgical procedures but also exacerbates the risk of postoperative infections. This article adds to the growing body of evidence that obesity is a significant risk factor for surgical site infections across various surgical contexts.

The systematic review by (S. Almasaudi et al., 2018) further delineated the relationship between BMI and postoperative infection rates in colorectal cancer surgeries. Their findings indicated that obesity is associated with a higher incidence of surgical site infections, prolonged hospital stays, and increased healthcare costs. The authors emphasized the need for regular monitoring of obese patients post-

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

surgery, as their unique physiological characteristics can lead to poorer operative outcomes. This study underscores the critical importance of understanding the impact of obesity on surgical recovery, particularly in high-risk populations.

(KARANTH & L. KARANTH, 2021) contributed to the discourse by examining the specific challenges faced by obese patients undergoing surgery in the South Indian population. Their research highlighted not only the increased duration of surgeries and higher rates of postoperative pain in obese individuals but also the necessity for tailored preoperative assessments and postoperative monitoring to mitigate complications. This article stresses the importance of developing targeted strategies to manage the unique needs of obese surgical patients.

Most recently, (Tefera Kassahun et al., 2022) conducted a retrospective study focusing on emergency laparotomy procedures for high-risk abdominal emergencies. Their findings reinforced previous conclusions that obesity significantly increases the likelihood of complications and mortality in surgical patients. The study highlighted that comorbid conditions associated with obesity play a crucial role in determining postoperative outcomes, thus necessitating a comprehensive approach to managing obese patients in emergency surgical contexts.

Overall, the literature consistently demonstrates that obesity is a significant risk factor for adverse surgical outcomes, particularly in major abdominal surgeries. The interplay between obesity, comorbidities, and surgical complications necessitates a nuanced understanding and proactive management strategies to optimize patient care and surgical success.

LITERATURE REVIEW

The article titled "Obesity and Surgical Wound Healing: A Current Review" by (N. Pierpont et al., 2014) provides a comprehensive examination of the impact of obesity on surgical outcomes, particularly focusing on wound healing and postoperative complications. The authors articulate that obesity is not merely a

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

singular factor but rather a complex interplay of comorbidities that exacerbate surgical risks and hinder recovery processes.

The article's main premise highlights that obesity is significantly associated with increased rates of surgical complications, including wound infections and wound separation. This assertion is supported by a plethora of studies that demonstrate a direct correlation between Body Mass Index (BMI) and adverse surgical outcomes. The authors emphasize that even modest weight loss can lead to substantial improvements in various health metrics, including a 30% reduction in diabetes-related morbidity and mortality with just a 10-kilogram weight loss. This finding underscores the importance of preoperative weight management as a potential strategy to enhance surgical recovery and outcomes in obese patients.

Moreover, the authors delve into the implications of visceral adiposity, which they argue may pose an even greater risk than BMI alone. This distinction is critical as it suggests that interventions targeting visceral fat could be pivotal in improving surgical outcomes. The article also notes that a higher BMI is linked to increased infection rates in trauma patients, reinforcing the notion that obesity creates a challenging environment for wound healing.

The authors further contextualize their findings within broader health concerns, indicating that the negative outcomes associated with obesity extend beyond surgical complications to include increased mortality rates from various cancers and cardiovascular diseases. The consistency of these findings across different demographics lends credence to the argument that obesity is a significant public health issue that requires urgent attention, particularly in surgical contexts.

Despite the robust evidence presented, the article could benefit from a more detailed exploration of potential interventions that could mitigate these risks. While the authors acknowledge the role of weight loss, the discussion lacks specific strategies or protocols that could be implemented preoperatively to optimize patient outcomes.

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Surgical site infection in overweight and obese Total Knee Arthroplasty patients" by (John Wilson et al., 2018) provides a comprehensive examination of the implications of obesity on surgical outcomes, particularly focusing on the incidence of surgical site infections (SSIs) following total knee arthroplasty (TKA). The authors present a compelling argument that obesity serves as a significant risk factor for postoperative complications, echoing broader concerns within surgical literature regarding the impact of obesity on recovery and infection rates.

The authors delve into the multifaceted relationship between obesity and surgical outcomes, emphasizing that increased body mass index (BMI) correlates with heightened risks of SSIs. They cite evidence that obese patients not only face a higher likelihood of infection but also experience prolonged recovery times and increased healthcare costs. This aligns with the findings from meta-analyses that indicate poorer outcomes in obese patients undergoing various surgical procedures, including laparoscopic colorectal surgery and cardiac surgery, as highlighted in their discussion ((John Wilson et al., 2018)).

A critical evaluation of the article reveals that while the authors effectively outline the risks associated with obesity, they could have expanded on the underlying mechanisms that contribute to these increased risks. For instance, the article mentions the role of obesity in modulating the immune response, suggesting that the inflammatory state associated with obesity may impair the body’s ability to combat infections. However, a more detailed exploration of the immunological complications of obesity would provide readers with a deeper understanding of how these factors interplay during the postoperative period.

Furthermore, the article addresses the financial implications of surgical procedures in obese patients, indicating that the increased risk of complications leads to higher healthcare costs. This aspect is particularly relevant in discussions surrounding healthcare resource allocation and the management of surgical patients. The authors could enhance their argument by incorporating statistical data on the economic burden of obesity-related complications to further substantiate their claims.

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article titled "The relationship between body mass index and short term postoperative outcomes in patients undergoing potentially curative surgery for colorectal cancer: a systematic review and meta-analysis" by (S. Almasaudi et al., 2018) provides a comprehensive analysis of how body mass index (BMI) correlates with postoperative complications, particularly infections, in colorectal cancer surgeries. This systematic review synthesizes data from various studies to elucidate the impact of obesity on surgical outcomes, highlighting the pressing issue of infection rates among obese patients.

The authors meticulously outline the relationship between obesity and the incidence of postoperative infections, noting that obesity is a significant risk factor regardless of ethnicity. The review underscores the association of postoperative site infections with extended hospital stays, elevated readmission rates, and heightened healthcare costs. This finding is critical, as it emphasizes the broader implications of obesity on healthcare systems, particularly in the context of major abdominal surgeries.

One of the key insights from the review is the distinction between types of infections, specifically the increased risk of deep or organ space infections among obese patients. This is particularly concerning, as these complications are linked to not only immediate postoperative challenges but also long-term outcomes, including recurrence rates and overall survival. The authors advocate for regular monitoring of obese patients during the postoperative period to mitigate these risks. This recommendation is well-founded, considering the evidence presented regarding the detrimental effects of excess body fat on wound healing and infection susceptibility.

Moreover, the article draws attention to the potential benefits of laparoscopic surgery over open surgery for obese individuals. The authors cite a prior meta-analysis indicating that laparoscopic approaches may reduce the risk of surgical site infections, despite a higher incidence of superficial infections. This nuanced understanding contributes to the ongoing discourse on surgical techniques and their implications for patient outcomes, particularly in the context of obesity.

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Obesity: is it an additional risk factor in analyzing surgical outcomes in the South Indian population?" by (KARANTH & L. KARANTH, 2021) provides a critical examination of the implications of obesity on surgical outcomes, particularly focusing on major abdominal surgeries within a specific demographic. The authors present compelling evidence that obesity significantly influences surgical parameters, including increased duration of surgery and heightened post-operative pain levels.

One of the key insights from the study is the observed correlation between obesity and a slight increase in 30-day readmission rates, particularly within the infection subset. This finding underscores the necessity for heightened vigilance in the management of post-operative care for obese patients. The authors argue convincingly that there is a pressing need for tailored pain management strategies and meticulous monitoring of potential complications in this population. Their emphasis on early intervention is particularly noteworthy; it highlights the critical nature of proactive measures in mitigating the risks associated with obesity during the surgical recovery process.

The study also raises important considerations regarding the broader implications of rising obesity rates globally. As these rates continue to escalate, the potential for adverse post-operative outcomes is likely to increase correspondingly. The authors advocate for the development of safe, alternative, and minimally invasive surgical techniques that can accommodate the challenges posed by increased adiposity. This perspective is crucial, as it aligns with the current trends in surgical practice that prioritize patient safety and effective recovery protocols.

Moreover, the authors recommend the establishment of comprehensive preoperative assessments for high-risk patients, which would facilitate better postoperative treatment and complication control. This approach is essential, as it not only aims to enhance patient outcomes but also seeks to optimize resource utilization within healthcare systems.

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article titled "The impact of obesity on surgical outcomes in patients undergoing emergency laparotomy for high-risk abdominal emergencies," authored by (Tefera Kassahun et al., 2022), provides a comprehensive analysis of the implications of obesity on postoperative outcomes in a specific surgical context. The study employs a retrospective single-center design, which allows for a focused examination of the relationship between obesity and adverse surgical outcomes in patients undergoing emergency laparotomy.

The authors highlight that patients with obesity exhibit significantly higher rates of complications and mortality compared to their nonobese counterparts. This finding aligns with existing literature that underscores the detrimental impact of obesity on surgical outcomes. The study meticulously documents that nearly all postoperative-specific complications were more prevalent among obese patients, indicating a pervasive influence of obesity on surgical recovery.

A critical aspect of the article is its exploration of the comorbidity rates associated with obesity. The authors report that patients with obesity not only face higher complication rates but also have a significantly increased prevalence of comorbid conditions, particularly cardiovascular and renal diseases. This association suggests that the increased complication rates observed in obese patients may be partially attributable to these underlying health issues. The study effectively emphasizes the importance of considering comorbidities when evaluating surgical risks in obese patients, thus contributing to a nuanced understanding of the challenges faced in surgical settings.

Moreover, the article discusses the implications of prolonged mechanical ventilation and extended hospital stays for patients with obesity. The authors note that these factors further exacerbate the risk of developing complications, which can ultimately lead to increased mortality rates. This insight is particularly valuable, as it highlights the need for targeted interventions and careful management strategies for obese patients undergoing high-risk abdominal surgeries.

CONCLUSION

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The literature consistently illustrates that obesity is a significant risk factor for adverse surgical outcomes, particularly following major abdominal surgeries. The introduction emphasizes the multifaceted implications of increased body mass index (BMI) on surgical recovery and complication rates, highlighting a strong correlation between obesity and increased postoperative complications such as wound infections and mortality (N. Pierpont et al., 2014).

The systematic review by (S. Almasaudi et al., 2018) further elucidates this relationship, indicating that obesity is linked to higher rates of surgical site infections and prolonged hospital stays in colorectal cancer surgeries. The authors advocate for regular monitoring of obese patients post-surgery due to their unique physiological characteristics that can lead to poorer operative outcomes. This sentiment is echoed in the findings of (John Wilson et al., 2018), which specifically address the heightened risk of deep infections in obese patients undergoing total knee arthroplasty, reinforcing the critical nature of obesity as a risk factor across various surgical contexts.

Moreover, the article by (KARANTH & L. KARANTH, 2021) highlights the challenges faced by obese patients in the South Indian population, noting increased surgery durations and postoperative pain, which necessitate tailored preoperative assessments and postoperative monitoring. This need for individualized care is further emphasized in the retrospective study by (Tefera Kassahun et al., 2022), which underscores the significant complications and mortality rates associated with obesity in emergency laparotomy procedures.

In conclusion, the body of literature reviewed indicates a clear and consistent pattern that obesity is a critical factor influencing surgical outcomes. The interplay between obesity, comorbidities, and surgical complications necessitates a nuanced understanding and proactive management strategies to optimize patient care and surgical success. Addressing obesity through preoperative weight management and tailored postoperative care is essential in improving surgical outcomes for this high-risk population.

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. N. Pierpont, Y., Phuong Dinh, T., Emerick Salas, R., L. Johnson, E., G. Wright, T., C. Robson, M., & G. Payne, W., 2014. Obesity and Surgical Wound Healing: A Current Review. ncbi.nlm.nih.gov
2. John Wilson, C., Georgiou, K., Oburu, E., Theodoulou, A., H Deakin, A., & Krishnan, J., 2018. Surgical site infection in overweight and obese Total Knee Arthroplasty patients. [PDF]
3. S. Almasaudi, A., T. McSorley, S., A. Edwards, C., & C. McMillan, D., 2018. The relationship between body mass index and short term postoperative outcomes in patients undergoing potentially curative surgery for colorectal cancer: a systematic review and meta-analysis. [PDF]
4. KARANTH, D. I. V. Y. A. & L. KARANTH, V. E. E. N. A., 2021. Obesity: is it an additional risk factor in analyzing surgical outcomes in the South Indian population?. ncbi.nlm.nih.gov
5. Tefera Kassahun, W., Mehdorn, M., & Babel, J., 2022. The impact of obesity on surgical outcomes in patients undergoing emergency laparotomy for high-risk abdominal emergencies. ncbi.nlm.nih.gov

Downloaded from The Medical Journal of Urology and General Surgery
Copyright © International Medical Journal Corp. Ltd. All rights reserved