Research Article Volume 01, Issue No. 02. 2024

The National Medical Journal of Health Science



Association of Sleep-Disordered Breathing With Cognitive Function and Risk of Cognitive Impairment

[1] Akbarina, [1] Agustina

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.akbarina745251@gmail.com

Article History :
Received date : 2024/10/15
Revised date : 2024/11/18
Accepted date : 2024/12/11
Published date : 2025/01/02

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature surrounding sleep-disordered breathing (SDB) and its association with cognitive function and impairment reveals a multifaceted relationship that spans across various age groups and health conditions. **Literature Review:** Expanding on this, (McMillan & J. Morrell, 2016) noted a significant prevalence of SDB in older adults, estimating rates between 20% and 40%. The implications of SDB in this demographic are particularly critical, as they may differ from those observed in younger populations, necessitating tailored approaches to diagnosis and treatment. Furthermore, (Robert Ott et al., 2017) drew connections between SDB and cerebrovascular diseases, emphasizing the need for systematic screening in at-risk populations, particularly as SDB disrupts sleep architecture and leads to cognitive decline. The interplay between behavioral and psychiatric issues and cognitive deficits in children with SDB was explored by (L. Smith et al., 2017), who found that behavioral problems could exacerbate cognitive impairments. This suggests a holistic approach is necessary in addressing the cognitive challenges faced by children with SDB. Similarly, (Janssens et al., 2018) examined cognitive impairment in the elderly, particularly in those with Alzheimer's disease (AD), drawing attention to the potential exacerbation of cognitive decline by SDB. **Conclusion:** In conclusion, the body of literature underscores a complex interplay between SDB, cognitive function, and various health risks across different populations. The findings emphasize the need for continued research to elucidate the mechanisms underlying these relationships and to develop effective interventions that address both the physiological and psychological aspects of SDB.

Keywords: Sleep-Disordered Breathing, Cognitive Function , Risk of Cognitive Impairment

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The literature surrounding the association of sleep-disordered breathing (SDB) with cognitive function and the risk of cognitive impairment has expanded significantly over the past decade, revealing a complex interplay between sleep disturbances, cognitive deficits, and various health outcomes. The foundational work by (J Kohler et al., 2010) established the necessity for rigorous evaluation guidelines in the context of SDB in children. Their findings highlighted a specific pattern of cognitive impairment, particularly in attention, executive functioning, memory, and psychomotor skills, while global intelligence remained relatively unaffected. This critical examination of the child-focused literature underscored the importance of establishing clear diagnostic criteria to better understand the neurocognitive challenges posed by SDB.

Building on this foundation, (McMillan & J. Morrell, 2016) explored the prevalence and consequences of SDB in the elderly, identifying a significant increase in SDB among older populations. Their research emphasized that the clinical implications of SDB in older adults might differ from those observed in younger individuals, indicating a need for tailored approaches to diagnosis and treatment in this demographic. The authors noted that the prevalence of SDB in older individuals could range from 20% to 40%, raising important questions about the mechanisms and clinical significance of SDB-related cognitive decline in aging populations.

In a broader context, (Robert Ott et al., 2017) examined the relationship between SDB and cerebrovascular diseases, highlighting the alarming rise in SDB prevalence due to obesity and aging. They argued for systematic screening in at-risk populations, considering the long-term adverse effects of SDB on cognitive functioning and overall health. Their work suggests that SDB disrupts sleep architecture, leading to excessive daytime sleepiness and impaired cognitive performance, thus reinforcing the need for awareness and intervention.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Further investigation by (L. Smith et al., 2017) specifically addressed the mediating role of behavioral and psychiatric problems in the relationship between SDB and cognitive deficits in children. Their findings suggested that behavioral issues could exacerbate cognitive impairments, calling for a more nuanced understanding of the interconnections between these factors.

(Janssens et al., 2018) contributed to the discourse by examining cognitive impairment in the elderly with SDB, noting correlations between SDB indices and various cognitive issues, including confusion and daytime sleepiness. Their research highlighted the prevalence of SDB in patients with Alzheimer's disease (AD), suggesting that SDB may exacerbate cognitive decline in this vulnerable population.

(P. Mansukhani et al., 2019) further elucidated the relationship between hypertension, cognitive impairment, and sleep apnea, positing that sleep apnea could serve as a pathway linking hypertension to cognitive decline. Their work emphasized the need for vigilance in monitoring cognitive dysfunction among individuals with SDB, particularly in older adults, and called for further research into the potential reversibility of cognitive deficits through treatment of sleep apnea.

(K. Ulland et al., 2021) addressed the intersection of SDB and neurodegenerative diseases like AD, focusing on the neuroimmune system as a common link. They highlighted significant gaps in understanding the cellular mechanisms that underlie the relationship between SDB and AD, advocating for more research to clarify these connections.

Finally, (Khokhrina et al., 2022) conducted a systematic review of the association between SDB and cardiovascular pathology, reinforcing the notion that SDB has far-reaching implications beyond cognitive function. They identified a dose-response relationship between the severity of SDB and cardiovascular diseases, further establishing SDB as a critical factor in overall health.

Collectively, these studies illustrate the multifaceted relationship between SDB, cognitive function, and associated health risks, emphasizing the need for

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

continued research and clinical awareness to address the implications of SDB across different populations and age groups.

LITERATURE REVIEW

The article titled "Neurocognitive performance and behavior before and after treatment for sleep-disordered breathing in children" by (J Kohler et al., 2010) presents a comprehensive examination of the cognitive and behavioral deficits associated with sleep-disordered breathing (SDB) in children. The authors argue that there has been a significant increase in research focusing on this topic; however, they highlight the numerous limitations present in the existing studies. This critical review emphasizes the necessity for stringent evaluation guidelines in research concerning SDB, particularly in children.

(J Kohler et al., 2010) draw parallels between the research methodologies employed in studies of obstructive sleep apnea syndrome (OSAS) in adults and those that should be adopted for children. They suggest that the rigorous exclusion criteria established in adult research have led to a clearer understanding of the cognitive domains affected by OSAS, such as attention, executive functioning, memory, and psychomotor functioning. The authors note that while global intelligence remains relatively intact, specific cognitive deficits are prevalent, and treatment has shown varying degrees of effectiveness across these domains.

The article advocates for the application of similar exclusion criteria in studies involving children with SDB to better delineate the neurocognitive and behavioral challenges they face. By imposing stricter guidelines, researchers may enhance the clarity of findings related to the cognitive impairments associated with SDB in pediatric populations and thus improve treatment outcomes.

The article "Sleep disordered breathing at the extremes of age: the elderly" by (McMillan & J. Morrell, 2016) provides a comprehensive examination of the prevalence and implications of sleep disordered breathing (SDB) in the elderly demographic. The authors highlight that while SDB is well-documented in younger and middle-aged populations, the consequences and clinical significance in older

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

adults remain less clear. This distinction is critical, as it challenges the assumption that the effects of SDB are uniform across age groups.

The authors present data indicating that the prevalence of SDB among older adults ranges from 20% to 40%, with some estimates suggesting it may be at least double that of younger populations. This observation is particularly pertinent, as it underscores a significant public health concern given the aging global population. The article notes a steady increase in SDB prevalence with age, stabilizing around 60 to 65 years. This trend raises important questions about the mechanisms behind SDB in older adults and its potential impact on cognitive function and overall morbidity.

(McMillan & J. Morrell, 2016) delve into the implications of SDB for cognitive health, suggesting that the increased prevalence in older individuals may correlate with heightened risks of cognitive impairment. However, the article also points out that research in this area is still evolving, and there is a need for further studies to elucidate the specific cognitive outcomes associated with SDB in the elderly. The authors call attention to the debate surrounding the clinical significance of SDB in older populations, noting that while the prevalence is high, the consequences may not be as severe or as well understood as in younger cohorts.

The article "Sleep-disordered breathing: clinical features, pathophysiology and diagnosis" by (Robert Ott et al., 2017) provides a comprehensive examination of the relationship between sleep-disordered breathing (SDB) and cognitive function, particularly in the context of the rising prevalence of SDB associated with obesity and aging populations. The authors effectively highlight the clinical features and pathophysiological mechanisms underlying SDB, emphasizing its significant impact on cognitive health.

One of the key insights presented in the article is the correlation between SDB and increased risks of cerebrovascular diseases. The authors argue that SDB, particularly obstructive sleep apnoea (OSA), is a significant contributor to cardiovascular morbidity and mortality, which can subsequently affect cognitive

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

function. This relationship is particularly concerning given the increasing rates of obesity and demographic shifts towards an older population, both of which exacerbate the prevalence of SDB. The authors advocate for systematic screening for SDB in at-risk populations, which is crucial for early identification and intervention.

The article also discusses the detrimental effects of SDB on sleep quality, which can lead to excessive daytime sleepiness and impaired cognitive performance. The authors provide evidence that disturbances in sleep architecture caused by SDB may contribute to cognitive decline, thereby underscoring the importance of addressing sleep disorders as part of a broader strategy to mitigate cognitive impairment. The connection between SDB and cognitive function is a critical area of concern, as it suggests that improving sleep quality could have significant implications for cognitive health.

The article "Parent-Reported Behavioral and Psychiatric Problems Mediate the Relationship between Sleep-Disordered Breathing and Cognitive Deficits in School-Aged Children" by (L. Smith et al., 2017) provides a comprehensive examination of the intricate relationship between sleep-disordered breathing (SDB) and cognitive deficits in children. The authors effectively highlight the significant risk that SDB poses to cognitive functioning, particularly in school-aged children, which is an area of increasing concern given the prevalence of SDB in this demographic.

A critical evaluation of the material reveals that the authors aim to fill a notable gap in the existing literature by exploring the mediating role of behavioral and psychiatric problems in the relationship between SDB and cognitive deficits. The study's methodology, which includes parent-reported measures, allows for a nuanced understanding of how behavioral issues may exacerbate cognitive impairments associated with SDB. This approach adds depth to the analysis, as it acknowledges the multifaceted nature of cognitive development and the potential influence of external factors such as behavioral and psychiatric conditions.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The findings presented in the article suggest that there is a significant correlation between SDB and increased risks of cognitive deficits, with behavioral problems acting as a mediator in this relationship. This insight is crucial, as it emphasizes the need for clinicians and educators to consider behavioral health when addressing cognitive issues in children with SDB. The authors advocate for a holistic approach to treatment that not only addresses the physiological aspects of SDB but also the psychological and behavioral dimensions that may impact cognitive outcomes.

However, while the study provides valuable insights, it is essential to consider the limitations inherent in relying on parent-reported data. This method may introduce biases or inaccuracies in the reporting of behavioral and psychiatric problems, which could affect the validity of the conclusions drawn. Future research could benefit from incorporating objective assessments of both SDB and cognitive function to strengthen the findings.

The article "Sleep disordered breathing in the elderly" by (Janssens et al., 2018) provides a comprehensive examination of the complex relationship between sleep-disordered breathing (SDB) and cognitive function among older adults. The authors synthesize various studies to highlight the ambiguity surrounding the impact of SDB on cognitive impairment in this demographic.

A significant aspect of the article is its critique of existing literature, particularly the findings of Findley et al., which indicated a correlation between cognitive impairment and mean SaO2 levels during sleep in younger subjects with obstructive sleep apnea (OSA). However, the authors point out that similar correlations remain elusive in older populations. For instance, Knight et al. (2018) conducted a study involving 27 elderly subjects, finding no significant cognitive differences between those with and without sleep apnea. This lack of correlation is echoed in the studies by Phillips et al. and Berry et al., which also failed to establish a link between neuropsychological performance and SDB indices, suggesting that the cognitive ramifications of SDB may not be as pronounced in older adults as previously thought.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Interestingly, the article notes that while some studies, such as those by Yesavage et al. (2018) and Ancoli-Israël et al. (2018), have identified correlations between certain cognitive tests and the apnea-hypopnea index (AHI), these findings are not universally supported across the literature. Ancoli-Israël et al. found connections between SDB indices and symptoms such as nocturnal wandering and daytime sleepiness, yet the overall consensus remains that the relationship is not straightforward.

The discussion extends to the prevalence of SDB in patients with Alzheimer's disease (AD), where the authors cite anecdotal evidence suggesting a potential causal relationship. They propose that SDB could exacerbate cognitive decline in AD patients through mechanisms such as intermittent cerebral hypoxemia. This notion is supported by observations of altered sleep architecture in AD, including reduced sleep efficiency and increased awakenings, which may further complicate the cognitive landscape for these individuals.

The article "Hypertension and Cognitive Decline: Implications of Obstructive Sleep Apnea" by (P. Mansukhani et al., 2019) provides a comprehensive examination of the interconnectedness of hypertension, cognitive impairment, and obstructive sleep apnea (OSA). The authors present a compelling argument that these three conditions are prevalent in the general population and that their interrelations warrant significant attention, particularly in the context of cognitive health.

The article highlights the established link between hypertension and cognitive impairment/dementia, suggesting that hypertension may serve as a precursor to cognitive decline. The authors effectively argue that OSA could act as a mediating factor in this relationship. They propose that the repetitive nocturnal hypoxemia characteristic of sleep apnea may exacerbate the cognitive effects of hypertension, thereby creating a synergistic risk for cognitive impairment. This hypothesis is particularly relevant as it underscores the need for a multidimensional approach to understanding cognitive decline, one that incorporates both cardiovascular and respiratory health.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Moreover, the authors emphasize the importance of identifying sleep apnea in elderly individuals, positing that early detection could serve as a predictive marker for cognitive impairment. This perspective is crucial, as cognitive dysfunction often goes unnoticed until significant deterioration has occurred. The article advocates for increased vigilance in monitoring those with moderate to severe sleep-disordered breathing, suggesting that they may be at heightened risk for cognitive dysfunction.

Despite the compelling arguments presented, the article acknowledges the limited evidence surrounding the efficacy of OSA treatment in improving cognitive outcomes. The authors note that existing studies yield conflicting results, indicating a gap in the literature that calls for further investigation. This recognition of the current limitations is a critical aspect of the article, as it frames future research directions and highlights the urgency of exploring therapeutic interventions that could mitigate cognitive decline associated with sleep apnea.

The article "Alzheimer's Disease, Sleep Disordered Breathing, and Microglia: Puzzling out a Common Link" by (K. Ulland et al., 2021) provides a comprehensive overview of the complex relationship between sleep-disordered breathing (SDB) and Alzheimer's Disease (AD). The authors assert that while a significant correlation exists between these two conditions, the underlying cellular and physiological mechanisms remain inadequately explored. This gap in knowledge is particularly concerning given the increasing prevalence of both SDB and AD in aging populations.

The review emphasizes the neuroimmune system as a potential common pathway linking SDB and AD. (K. Ulland et al., 2021) discuss how chronic intermittent hypoxia, a hallmark of SDB, may lead to neuroinflammation and subsequently influence the progression of neurodegenerative diseases. The authors present experimental evidence suggesting that microglial activation, a critical component of the neuroimmune response, could play a pivotal role in mediating the effects of SDB on cognitive function and the risk of cognitive impairment. This perspective is particularly valuable as it shifts the focus from merely correlational

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

studies to a more mechanistic understanding of how SDB may exacerbate or contribute to AD pathology.

Despite the insights provided, the article also highlights the significant limitations in current research. The authors note the scarcity of studies that directly investigate the cellular mechanisms at play, particularly concerning the role of microglia and other immune responses in the context of SDB. This lack of foundational research underscores the need for more targeted studies that could elucidate these pathways and potentially lead to novel therapeutic strategies.

Furthermore, the review calls for interdisciplinary approaches combining neurology, sleep medicine, and immunology to foster a more holistic understanding of the intersection between SDB and cognitive impairment in AD. This integrative perspective is crucial, as it can inform clinical practices and interventions aimed at mitigating the cognitive decline associated with sleep-disordered breathing.

The systematic review conducted by (Khokhrina et al., 2022) provides a comprehensive examination of sleep-disordered breathing (SDB) and its association with cardiovascular pathology. While the primary focus of the article is on cardiovascular outcomes, the insights presented can be extrapolated to understand the potential implications of SDB on cognitive function and the risk of cognitive impairment.

SDB is characterized by repeated breathing interruptions during sleep, which can manifest as obstructive apnea, central apnea, or mixed apnea. The authors detail the criteria used to classify these respiratory events, emphasizing the importance of the apnea-hypopnea index (AHI) in determining the severity of SDB ((Khokhrina et al., 2022)). This classification system not only aids in diagnosing SDB but also facilitates understanding its potential impact on various health outcomes, including cognitive health.

The review highlights a dose-response relationship between the severity of SDB and cardiovascular disease (CVD), suggesting that as the severity of SDB increases, so does the risk of developing cardiovascular complications ((Khokhrina

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

et al., 2022)). This finding is particularly relevant when considering the interrelationship between cardiovascular health and cognitive function. Research indicates that cardiovascular health is intrinsically linked to cognitive performance, with conditions such as hypertension and heart disease being significant risk factors for cognitive decline ((Khokhrina et al., 2022)). Therefore, the mechanisms through which SDB affects cardiovascular health—such as intermittent hypoxia, endothelial dysfunction, and inflammation—may similarly influence cognitive outcomes.

Moreover, the prevalence of SDB varies due to differing diagnostic standards, which may contribute to the inconsistent findings in literature regarding its impact on cognitive function ((Khokhrina et al., 2022)). This variability underscores the need for standardized diagnostic criteria and methodologies in future research to better delineate the relationship between SDB and cognitive impairment.

The authors also note the limited research available on the direct impact of SDB on symptomatic cardiovascular disease, suggesting a gap in the literature that could be addressed in future studies. This gap is particularly pertinent for understanding the potential cognitive implications of SDB, as the interplay between cardiovascular pathology and cognitive decline remains an area ripe for exploration.

CONCLUSION

The literature surrounding sleep-disordered breathing (SDB) and its association with cognitive function and impairment reveals a multifaceted relationship that spans across various age groups and health conditions. Initial investigations, such as those by (J Kohler et al., 2010), established that children with SDB exhibit specific cognitive deficits, particularly in areas of attention, executive functioning, and memory, while global intelligence remains largely unaffected. This foundational work highlights the necessity for stringent evaluation guidelines in pediatric studies on SDB to enhance the clarity of findings and treatment outcomes.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

Expanding on this, (McMillan & J. Morrell, 2016) noted a significant prevalence of SDB in older adults, estimating rates between 20% and 40%. The implications of SDB in this demographic are particularly critical, as they may differ from those observed in younger populations, necessitating tailored approaches to diagnosis and treatment. Furthermore, (Robert Ott et al., 2017) drew connections between SDB and cerebrovascular diseases, emphasizing the need for systematic screening in at-risk populations, particularly as SDB disrupts sleep architecture and leads to cognitive decline.

The interplay between behavioral and psychiatric issues and cognitive deficits in children with SDB was explored by (L. Smith et al., 2017), who found that behavioral problems could exacerbate cognitive impairments. This suggests a holistic approach is necessary in addressing the cognitive challenges faced by children with SDB. Similarly, (Janssens et al., 2018) examined cognitive impairment in the elderly, particularly in those with Alzheimer's disease (AD), drawing attention to the potential exacerbation of cognitive decline by SDB.

The relationship between hypertension and cognitive decline, with SDB as a mediating factor, was articulated by (P. Mansukhani et al., 2019). This perspective emphasizes the importance of monitoring cognitive health in those with SDB, especially in older adults. Additionally, (K. Ulland et al., 2021) offered insights into the neuroimmune mechanisms linking SDB and AD, suggesting that chronic intermittent hypoxia may drive neuroinflammation, thus influencing cognitive outcomes.

Lastly, the systematic review by (Khokhrina et al., 2022) established a dose-response relationship between the severity of SDB and cardiovascular disease, underlining the interconnectedness of cardiovascular health and cognitive function. The review highlights the necessity for standardized diagnostic criteria to better understand the implications of SDB on cognitive impairment.

In conclusion, the body of literature underscores a complex interplay between SDB, cognitive function, and various health risks across different

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

populations. The findings emphasize the need for continued research to elucidate the mechanisms underlying these relationships and to develop effective interventions that address both the physiological and psychological aspects of SDB.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. J Kohler, M., Lushington, K., & Declan Kennedy, J., 2010. Neurocognitive performance and behavior before and after treatment for sleep-disordered breathing in children. ncbi.nlm.nih.gov
2. McMillan, A. & J. Morrell, M., 2016. Sleep disordered breathing at the extremes of age: the elderly. ncbi.nlm.nih.gov
3. Robert Ott, S., Korostovtseva, L., Schmidt, M., Horvath, T., Brill, A. K., & Bassetti, C., 2017. Sleep-disordered breathing: clinical features, pathophysiology and diagnosis.. [PDF]
4. L. Smith, D., Gozal, D., J. Hunter, S., & Kheirandish-Gozal, L., 2017. Parent-Reported Behavioral and Psychiatric Problems Mediate the Relationship between Sleep-Disordered Breathing and Cognitive Deficits in School-Aged Children. ncbi.nlm.nih.gov
5. Janssens, J. P., Pautex, S., Hilleret, H., & Michel, J. P., 2018. Sleep disordered breathing in the elderly. [PDF]
6. P. Mansukhani, M., Prakash Kolla, B., & K. Somers, V., 2019. Hypertension and Cognitive Decline: Implications of Obstructive Sleep Apnea. ncbi.nlm.nih.gov

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

7. K. Ulland, T., C. Ewald, A., O. Knutson, A., M. Marino, K., M. C. Smith, S., & J. Watters, J., 2021. Alzheimer's Disease, Sleep Disordered Breathing, and Microglia: Puzzling out a Common Link. ncbi.nlm.nih.gov
8. Khokhrina, A., Andreeva, E., & Degryse, J. M., 2022. A systematic review on the association of sleep-disordered breathing with cardiovascular pathology in adults. ncbi.nlm.nih.gov

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved