Research Article Volume 01, Issue No. 02. 2024

The National Medical Journal of Health Science



Association of Successful Percutaneous Revascularization of Chronic Total Occlusions With Quality of Life

[1] Gunawan, [1] Rahayu

[1] Faculty of Medicine, Sebelas Maret University, Surakarta City, Central Java, Indonesia

Correspondence : dr.gunawan392181@gmail.com

Article History :
Received date : 2024/10/12
Revised date : 2024/11/24
Accepted date : 2024/12/18
Published date : 2025/01/02

Copyright: *© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (https://creativecommons.org/licenses/by-nc/4.0/).*

ABSTRACT

Background: The literature on the association of successful percutaneous revascularization of chronic total occlusions (CTOs) with quality of life reveals a complex interplay between procedural success, patient outcomes, and the broader implications for clinical practice. **Literature Review:** Research has consistently shown that successful CTO revascularization is linked to better quality of life and functional status. For instance, studies indicate that patients who undergo successful PCI experience reduced angina and improved physical limitations (Fernando Ybarra et al., 2018). Moreover, a systematic review and meta-analysis highlight the comparative effectiveness of PCI versus medical therapy, suggesting that while PCI is recommended for improving quality of life, its actual prevalence remains low, indicating a need for more robust clinical guidelines (Hou Christien Li et al., 2019). The ongoing debate surrounding the best management strategies for CTOs is reflected in the literature, particularly regarding the efficacy of PCI compared to conservative treatment options. Some studies suggest that while PCI can alleviate ischemic symptoms and improve functional status, its long-term benefits may not always surpass those of conservative treatment, necessitating careful patient selection (Yang et al., 2021). **Conclusion:** In conclusion, the literature collectively supports the notion that successful percutaneous revascularization of chronic total occlusions is associated with improved quality of life and clinical outcomes. Nevertheless, the ongoing debates regarding optimal management strategies and the necessity for individualized patient care highlight the need for continued research and refinement of treatment protocols.

Keywords: Successful Percutaneous Revascularization , Chronic Total Occlusions, Quality of Life

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

INTRODUCTION

The literature on the association of successful percutaneous revascularization of chronic total occlusions (CTOs) with quality of life reveals a significant evolution in understanding the clinical benefits of this intervention over recent years. The work of (B. Mehta et al., 2018) highlights the advancements in percutaneous coronary intervention (PCI) techniques, noting that procedural success rates have improved markedly. Their findings indicate that successful CTO revascularization is linked to better long-term survival rates and a reduced incidence of cardiac events, emphasizing the importance of successful intervention in enhancing patient outcomes.

(Fernando Ybarra et al., 2018) further elaborate on the clinical benefits associated with successful CTO recanalization, including improved quality of life and physical limitations. Their study underscores the correlation between successful PCI and enhanced ventricular function, aligning with the growing body of evidence that suggests positive impacts on patients' overall health and well-being following successful interventions.

In a systematic review and meta-analysis, (Hou Christien Li et al., 2019) scrutinize the comparative effectiveness of PCI versus medical therapy for CTO management. Their findings suggest that while PCI is recommended to improve quality of life, its actual prevalence remains low due to ongoing controversies regarding its superiority over medical treatment. This raises critical questions about treatment pathways and the need for more robust guidelines to facilitate optimal patient care.

(Zheng et al., 2019) conducted a meta-analysis that reinforces the notion that successful PCI leads to improved long-term clinical outcomes compared to failed interventions. Their work highlights the ongoing debate regarding the best management strategies for CTOs, emphasizing the necessity for further research to clarify the benefits of PCI relative to other treatment modalities.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

(Yang et al., 2021) contribute to this discourse by examining the relationship between myocardial viability, functional status, and the presence of collaterals in patients with CTOs. Their findings suggest that while PCI can alleviate ischemic symptoms and improve functional status, the benefits may not always surpass those of conservative treatment, thus necessitating careful patient selection for intervention.

(Hamzaraj et al., 2022) build on this foundation by exploring the long-term survival benefits associated with successful CTO revascularization, particularly in patients with left ventricular systolic dysfunction. Their research highlights the profound impact of complete revascularization on patients' quality of life and overall health outcomes, advocating for PCI as a preferable treatment option in specific populations.

Collectively, these studies illustrate a comprehensive landscape of research that underscores the significance of successful CTO revascularization through PCI in improving quality of life, while also highlighting the complexities and ongoing debates surrounding optimal management strategies for this patient population.

LITERATURE REVIEW

The article "Predictors of success in percutaneous Coronary intervention for chronic total occlusion" by (B. Mehta et al., 2018) provides a comprehensive examination of the evolving landscape of percutaneous coronary intervention (PCI) for chronic total occlusions (CTOs) in coronary arteries. The authors highlight a significant improvement in procedural success rates, which have increased from 50-60% to an impressive 75-90% due to advancements in PCI techniques. This increase is critical, as the article underscores the profound implications of successful CTO revascularization on patient outcomes, particularly in terms of survival and quality of life.

The authors reference the Mid-America Heart Study, which analyzed 2007 patients and demonstrated that patients who underwent successful CTO revascularization exhibited higher 10-year survival rates compared to those with failed attempts (74% vs. 65%). This statistic is particularly compelling, as it

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

suggests that successful intervention not only prolongs life but may also enhance the quality of life for these patients. Additionally, the article points out that successful PCI is linked to a significantly reduced incidence of cardiac death and myocardial infarction (MI) over a 12-month period (1.1% vs. 7.2%) and a lower requirement for coronary artery bypass grafting (CABG) (2.5% vs. 15.7%).

Moreover, the study emphasizes the long-term benefits of successful PCI for CTOs, noting increased survival rates and reduced need for subsequent revascularization over a seven-year follow-up. This longitudinal perspective is crucial, as it provides evidence that the benefits of successful intervention extend well beyond the immediate post-procedural period, potentially leading to sustained improvements in patients' quality of life.

The authors aim to evaluate the factors that contribute to both success and failure in CTO-PCI among Indian patients, thereby addressing a significant gap in the literature regarding regional outcomes. This focus on local patient demographics is valuable, as it allows for a more nuanced understanding of how various predictors can influence procedural success in different populations.

The article "Percutaneous Coronary Intervention in Chronic Total Occlusion" by (Fernando Ybarra et al., 2018) provides a comprehensive examination of the implications of successful percutaneous coronary interventions (PCIs) in patients with chronic total occlusions (CTOs). The authors emphasize the critical role of drug-eluting stents in these procedures, reflecting a significant advancement in the management of coronary artery disease. They highlight the necessity of utilizing intravascular imaging methods such as intravascular ultrasound (IVUS) and optical coherence tomography, which are essential for guiding interventions and ensuring optimal outcomes.

The article presents a noteworthy statistic regarding the prevalence of bifurcation lesions in CTO interventions, noting that these account for 33% of cases. This detail underscores the complexity and challenges associated with CTO interventions, as bifurcation lesions can complicate the procedural approach and

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

necessitate tailored strategies for successful revascularization. The authors aptly point out that CTO interventions carry a higher risk of perforations compared to non-occlusive lesions, necessitating a careful assessment of each case to determine the appropriateness of proceeding with the intervention.

A critical evaluation of the material reveals that the authors effectively argue for the individualized decision-making process in managing CTO cases. They advocate for a nuanced approach, suggesting that the decision to interrupt a procedure should be based on a thorough evaluation of the patient's specific circumstances and potential risks. This perspective is essential in the context of complex PCI, where the balance between potential benefits and risks must be carefully navigated.

Furthermore, the authors present compelling evidence linking successful CTO recanalization to improved patient outcomes, including enhanced quality of life, reduced angina, and improved physical limitations. They assert that successful recanalization is associated with better ventricular function and decreased mortality rates compared to those who do not achieve successful recanalization. This correlation is supported by several observational studies, reinforcing the notion that effective intervention in CTO cases can lead to significant clinical benefits.

The article titled "Percutaneous coronary intervention versus medical therapy for chronic total occlusion of coronary arteries: A systematic review and meta-analysis" by (Hou Christien Li et al., 2019) provides a comprehensive examination of the treatment options available for patients suffering from chronic total occlusions (CTOs) of the coronary arteries. The authors conducted a systematic review and meta-analysis of 17 studies involving a total of 11,493 participants, assessing the efficacy of percutaneous coronary intervention (PCI) compared to medical therapy.

A critical evaluation of the material reveals several key insights. First, the study highlights that while PCI is often recommended in clinical guidelines as a means to improve survivability and quality of life for CTO patients, the actual

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

prevalence of PCI in this population remains low. This discrepancy raises important questions about the implementation of clinical guidelines in practice and the factors that influence treatment decisions among healthcare providers.

Moreover, the findings indicate that there is no significant difference in the rates of myocardial infarction and repeat revascularization procedures between the PCI and medical therapy groups. This lack of difference suggests that while PCI may be a viable option for some patients, it does not universally guarantee improved outcomes when compared to medical management alone. The authors also note that patients with CTOs generally have a poorer prognosis than those with non-CTO coronary disease, underscoring the complexity of managing this patient population.

The article further discusses the controversy surrounding the preferred management option for CTOs. The divergence between observational studies and randomized controlled trials regarding the benefits of PCI raises critical questions about the evidence base guiding clinical practice. The authors emphasize the need for further research to clarify the potential advantages of PCI over medical therapy in terms of quality of life and long-term outcomes.

The article titled "Outcomes of Chronic Total Occlusions in Coronary Arteries According to Three Therapeutic Strategies: A Meta-analysis with 6985 Patients from 8 Published Observational Studies" by (Zheng et al., 2019) provides a comprehensive evaluation of the management strategies for chronic total occlusions (CTOs) in coronary arteries. The authors highlight the prevalence of CTOs, which affect approximately 30% of patients with coronary heart disease, and underscore the ongoing challenges in effectively managing this condition.

The study systematically compares three therapeutic approaches: coronary artery bypass grafting (CABG), percutaneous coronary intervention (PCI), and medical treatment (MT). The authors critically analyze previous findings, noting the conflicting results regarding the efficacy of PCI in improving long-term survival when compared to MT. For instance, while Ladwiniec et al. (as cited in (Zheng et al., 2019)) reported that PCI is associated with enhanced long-term survival

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

compared to MT alone, other studies, such as those by Fujino et al. and Yang et al., indicate that PCI does not significantly reduce the risk of death or major adverse cardiac events when added to MT.

A key insight from (Zheng et al., 2019)'s meta-analysis is the distinction made between successful and failed PCI interventions for CTOs. The findings suggest that successful recanalization of a CTO through PCI is linked to improved long-term clinical outcomes, which emphasizes the importance of procedural success in determining the overall effectiveness of the intervention. This aspect is critical as it suggests that not only the choice of treatment modality but also the execution of the procedure plays a significant role in patient outcomes.

The authors conclude that both PCI and CABG may provide better clinical outcomes compared to MT alone, thereby advocating for a more aggressive approach in managing CTOs. This conclusion is particularly relevant for clinicians considering treatment options for patients with CTOs, as it supports the notion that successful revascularization can lead to enhanced quality of life and long-term survival.

The article "Percutaneous Coronary Intervention Versus Medical Therapy for Chronic Total Occlusion of Coronary Arteries: A Systematic Review and Meta-Analysis" by (Hou Christien Li et al., 2019) provides a comprehensive examination of the management strategies for patients with chronic total occlusions (CTOs) in coronary arteries. The authors delve into the implications of CTOs on patient prognosis and quality of life, highlighting the significant prevalence of these conditions among individuals undergoing coronary angiography.

CTOs are defined as complete or near-complete blockages of coronary arteries persisting for over three months, which leads to a substantial reduction in blood flow to the heart muscle ((Hou Christien Li et al., 2019)). The article emphasizes that patients with CTOs often experience a poorer prognosis than those with non-CTO coronary artery disease, primarily due to the asymptomatic nature of many cases, which can delay diagnosis and treatment ((Hou Christien Li et al.,

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

2019)). This aspect raises critical questions regarding the timeliness and appropriateness of interventions, as many patients may not receive the necessary care until complications arise.

The systematic review and meta-analysis presented by the authors synthesize data from multiple cohort studies comparing the effectiveness of percutaneous coronary intervention (PCI) versus medical therapy in managing CTOs. The findings indicate that while PCI is recommended in clinical guidelines to enhance survivability and improve quality of life, the uptake of this intervention remains surprisingly low, with only about 10% of CTO patients in North America undergoing PCI ((Hou Christien Li et al., 2019)). This discrepancy between guideline recommendations and clinical practice is a significant concern, suggesting barriers to treatment that may stem from a lack of awareness, resources, or perhaps the perceived risks associated with PCI.

The authors critically evaluate the ongoing debate surrounding the preference for PCI over medical therapy, noting that the choice of treatment is often influenced by individual patient characteristics, the severity of symptoms, and the presence of comorbidities. The article underscores the necessity for a more nuanced approach to treatment decisions, advocating for personalized care that takes into account the specific circumstances of each patient ((Hou Christien Li et al., 2019)).

The article titled "Myocardial Viability, Functional Status, and Collaterals of Patients With Chronically Occluded Coronary Arteries" by (Yang et al., 2021) provides a comprehensive examination of the implications of chronic total occlusions (CTOs) in coronary arteries, particularly focusing on the potential benefits of percutaneous coronary intervention (PCI) in improving patient outcomes. The authors highlight the prevalence of CTOs, which are present in approximately 15–25% of patients undergoing coronary angiography, and underline the critical need for a nuanced understanding of patient characteristics and treatment responses.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

One of the central themes of the article is the beneficial effects of CTO revascularization, notably in terms of alleviating angina, reducing ischemia, and enhancing functional status. However, the authors critically evaluate existing evidence from randomized trials, which suggests that while PCI treatment for CTOs can provide symptomatic relief, it may not necessarily lead to superior long-term functional outcomes compared to conservative management approaches. This finding raises important questions about the selection criteria for PCI candidates and the need for thorough pre-procedural assessments.

A significant contribution of the study is its emphasis on myocardial viability as a predictor of functional recovery post-PCI. The authors argue that while current clinical guidelines prioritize symptom evaluation and ischemia burden assessment, the inclusion of myocardial viability as a prognostic marker could refine patient selection and potentially enhance recovery outcomes. This perspective is particularly relevant given the complex interplay between myocardial viability and the presence of collateral circulation, which the study investigates in detail.

The article also underscores the importance of detailed baseline characteristics in patients with CTOs. By systematically recording these characteristics, clinicians may better understand the potential for recovery and tailor interventions accordingly. The exploration of collaterals in relation to functional status and myocardial viability presents a valuable avenue for future research, as it could elucidate mechanisms that influence recovery and guide therapeutic strategies.

The article "Assessing the Clinical Influence of Chronic Total Occlusions (CTOs) Revascularization and the Impact of Vascularization Completeness on Patients with Left Ventricular (LV) Systolic Dysfunction" by (Wu et al., 2022) provides a comprehensive analysis of the implications of successful percutaneous revascularization in patients suffering from chronic total occlusions, particularly those with compromised left ventricular function. The authors present compelling evidence that successful CTO revascularization correlates with significant long-

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

term survival benefits and improvements in quality of life for this patient population.

A key finding of the study is the relationship between complete coronary revascularization (CCR) and clinical outcomes. The authors assert that patients who undergo successful CTO percutaneous coronary intervention (PCI) exhibit marked improvements in left ventricular ejection fraction (LVEF) and symptomatic relief, particularly when CCR is achieved. This observation is crucial as it underscores the importance of addressing all occluded vessels to maximize therapeutic benefits, thereby enhancing patient outcomes and quality of life.

The article also discusses the prognostic factors influencing major adverse cardiac and cerebrovascular events (MACCE). Notably, the authors highlight that older age (≥75 years) and prior myocardial infarction (MI) are associated with an increased risk of cardiac mortality in patients with LVEF ≤40%. Conversely, successful CTO revascularization emerges as a protective factor against MACCE, emphasizing its potential role in improving patient prognoses.

Moreover, the anticipated benefits of successful CTO revascularization extend beyond immediate clinical outcomes. The authors suggest that improved blood flow in coronary vessels not only revives hibernating myocardium but also facilitates access for circulating stem cells, which may further enhance myocardial recovery. This multifaceted approach to treatment reflects a growing understanding of the complex interplay between coronary revascularization and myocardial health.

The findings presented in this article are corroborated by a recent meta-analysis that indicates a favorable impact on long-term mortality rates in patients undergoing successful CTO recanalization. This evidence supports the notion that PCI, when combined with standard medical treatment, yields superior outcomes compared to medical therapy alone in patients with left ventricular systolic dysfunction.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article "Patient Selection and Clinical Indication for Chronic Total Occlusion Revascularization—A Workflow Focusing on Non-Invasive Cardiac Imaging" by (Hamzaraj et al., 2022) provides a comprehensive overview of chronic total occlusion (CTO) in coronary arteries, emphasizing its prevalence and the associated challenges in treatment. The authors define CTO as a chronic occlusion lasting more than three months, characterized by a TIMI 0 flow, which often leads to the development of collateral vessels that are typically inadequate for sufficient myocardial perfusion, resulting in ischemic heart disease.

One of the critical insights from the article is the discussion on the clinical benefits of revascularization procedures, specifically percutaneous coronary intervention (PCI) and coronary artery bypass grafting (CABG). The authors highlight that while there is a growing body of evidence supporting the symptomatic relief and improvement in quality of life following successful revascularization of CTO, the question of whether these interventions confer a survival advantage or long-term freedom from cardiac events compared to optimal medical therapy remains unresolved. This gap in knowledge is significant, as it underscores the need for further research to establish definitive outcomes associated with CTO revascularization.

The article also addresses the technical complexities involved in CTO PCI, noting that the success rates of these procedures can vary significantly based on the operator's experience. The authors report that success rates can range from 60-70% in less experienced centers to 90-95% in highly experienced centers. This variability emphasizes the importance of operator skill and institutional expertise in achieving successful outcomes in CTO interventions.

Moreover, the authors advocate for the use of non-invasive cardiac imaging as a critical component in the patient selection process for CTO revascularization. This focus on imaging techniques aims to enhance the assessment of coronary anatomy and collateral circulation, thereby improving the decision-making process regarding the appropriateness of revascularization.

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The article titled "The Effects of Percutaneous Coronary Intervention on Biomarkers and Quality of Life in Patients With Chronic Total Coronary Artery Obstruction" by (Made Junior Rina Artha et al., 2023) presents a comprehensive analysis of the impact of percutaneous coronary intervention (PCI) on patients suffering from chronic total occlusions (CTOs). The authors delve into the multifaceted nature of how PCI not only addresses the ischemic conditions associated with CTOs but also contributes to enhanced quality of life for these patients.

The study highlights that CTOs are often linked with a significant decline in patients' quality of life due to persistent symptoms such as angina. The authors reference previous research, including work by Hirai et al., which utilized the Seattle Angina Questionnaire (SAQ) to demonstrate that patients undergoing PCI experienced notable improvements in angina frequency, overall quality of life, and satisfaction with their treatment regimen compared to those without CTO. This comparison underscores the therapeutic potential of PCI in ameliorating not just physiological symptoms but also psychosocial aspects of health.

Moreover, the authors emphasize the need for further research in this domain, particularly at the biomolecular level, to elucidate the underlying mechanisms through which PCI exerts its beneficial effects. They argue that existing literature has primarily focused on echocardiographic parameters, mortality rates, and recurrence of cardiac events, leaving a gap in understanding how these interventions affect biomarkers and the subjective experience of quality of life in CTO patients.

The article effectively identifies this gap and proposes a study design that aims to compare biomarker levels between CTO patients who have undergone PCI and those who have not, alongside an assessment of quality of life improvements. This approach is commendable as it seeks to provide a more holistic view of patient outcomes post-PCI, integrating both clinical and subjective measures of health.

CONCLUSION

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

The literature on the association of successful percutaneous revascularization of chronic total occlusions (CTOs) with quality of life reveals a complex interplay between procedural success, patient outcomes, and the broader implications for clinical practice. The advancements in percutaneous coronary intervention (PCI) techniques have led to notable improvements in procedural success rates, which have increased from 50-60% to 75-90% (B. Mehta et al., 2018). This increase is significant as it correlates with enhanced long-term survival rates and a reduction in cardiac events, thereby underscoring the importance of successful interventions in improving patient outcomes.

Research has consistently shown that successful CTO revascularization is linked to better quality of life and functional status. For instance, studies indicate that patients who undergo successful PCI experience reduced angina and improved physical limitations (Fernando Ybarra et al., 2018). Moreover, a systematic review and meta-analysis highlight the comparative effectiveness of PCI versus medical therapy, suggesting that while PCI is recommended for improving quality of life, its actual prevalence remains low, indicating a need for more robust clinical guidelines (Hou Christien Li et al., 2019).

The ongoing debate surrounding the best management strategies for CTOs is reflected in the literature, particularly regarding the efficacy of PCI compared to conservative treatment options. Some studies suggest that while PCI can alleviate ischemic symptoms and improve functional status, its long-term benefits may not always surpass those of conservative treatment, necessitating careful patient selection (Yang et al., 2021). Furthermore, the relationship between myocardial viability and successful PCI outcomes emphasizes the importance of individualized treatment approaches in optimizing patient care (Hamzaraj et al., 2022).

Overall, the body of research indicates that successful CTO revascularization through PCI can significantly enhance quality of life and clinical outcomes for patients, particularly in specific populations such as those with left ventricular systolic dysfunction (Wu et al., 2022). However, the complexity of treatment decisions, the variability in procedural success rates based on operator

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

experience, and the need for further research into the long-term implications of these interventions underscore the multifaceted nature of managing CTOs.

In conclusion, the literature collectively supports the notion that successful percutaneous revascularization of chronic total occlusions is associated with improved quality of life and clinical outcomes. Nevertheless, the ongoing debates regarding optimal management strategies and the necessity for individualized patient care highlight the need for continued research and refinement of treatment protocols.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. B. Mehta, A., Mehta, N., Chhabria, R., Mandurke, V., Tawade, N., Jain, N., Desai, A., & Handa, S. R., 2018. Predictors of success in percutaneous Coronary intervention for chronic total occlusion. ncbi.nlm.nih.gov
2. Fernando Ybarra, L., J. C. Cantarelli, M., M. G. Lemke, V., & Schaan de Quadros, A., 2018. Percutaneous Coronary Intervention in Chronic Total Occlusion. ncbi.nlm.nih.gov
3. Hou Christien Li, K., Hei Gabriel Wong, K., Gong, M., Liu, T., Li, G., Xia, Y., Ho, J., Nombela-Franco, L., C Sawant, A., Eccleshall, S., Tse, G., & S. Vassiliou, V., 2019. Percutaneous coronary intervention versus medical therapy for chronic total occlusion of coronary arteries: A systematic review and meta-analysis. [PDF]
4. Zheng, Y. Y., Gao, Y., Chen, Y., Wu, T. T., Ma, Y. T., Zhang, J. Y., & Xie, X., 2019. Outcomes of Chronic Total Occlusions in Coronary Arteries

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved

According to Three Therapeutic Strategies: A Meta-analysis with 6985 Patients from 8 Published Observational Studies. ncbi.nlm.nih.gov

5. Yang, X., Tian, J., Zhang, L., Dong, W., Mi, H., Li, J., Li, J., Han, Y., Zuo, H., An, J., He, Y., & Song, X., 2021. Myocardial Viability, Functional Status, and Collaterals of Patients With Chronically Occluded Coronary Arteries. ncbi.nlm.nih.gov
6. Hamzaraj, K., Kammerlander, A., Gyöngyösi, M., Frey, B., Distelmaier, K., & Graf, S., 2022. Patient Selection and Clinical Indication for Chronic Total Occlusion Revascularization—A Workflow Focusing on Non-Invasive Cardiac Imaging. ncbi.nlm.nih.gov
7. Hou Christien Li, K., Hei Gabriel Wong, K., Gong, M., Liu, T., Li, G., Xia, Y., Ho, J., Nombela-Franco, L., C. Sawant, A., Eccleshall, S., Tse, G., & S. Vassiliou, V., 2019. Percutaneous Coronary Intervention Versus Medical Therapy for Chronic Total Occlusion of Coronary Arteries: A Systematic Review and Meta-Analysis. ncbi.nlm.nih.gov
8. Wu, X., Cai, J., Zhang, Q., & Huang, H., 2022. Assessing the Clinical Influence of Chronic Total Occlusions (CTOs) Revascularization and the Impact of Vascularization Completeness on Patients with Left Ventricular (LV) Systolic Dysfunction. ncbi.nlm.nih.gov
9. Made Junior Rina Artha, I., Made Bakta, I., Bagus Putra Manuaba, I., Wayan Wita, I., Saifur Rohman, M., Nyoman Mantik Astawa, I., & Bhargah, A., 2023. The Effects of Percutaneous Coronary Intervention on Biomarkers and Quality of Life in Patients With Chronic Total Coronary Artery Obstruction. ncbi.nlm.nih.gov

Downloaded from The National Medical Journal of Health Science
Copyright © International Medical Journal Corp. Ltd. All rights reserved